

Building evidence for conservation globally

Journal of Threatened Taxa

10.11609/jott.2026.18.6.29003-29158

www.threatenedtaxa.org

26 June 2026 (Online & Print)

18(6): 29003-29158

ISSN 0974-7907 (Online)

ISSN 0974-7893 (Print)



Open Access





ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

Publisher
Wildlife Information Liaison Development Society
www.wild.zooreach.org

Host
Zoo Outreach Organization
www.zooreach.org

Srivari Illam, No. 61, Karthik Nagar, 10th Street, Saravanampatti, Coimbatore, Tamil Nadu 641035, India
Registered Office: 3A2 Varadarajulu Nagar, FCI Road, Ganapathy, Coimbatore, Tamil Nadu 641006, India
Ph: +91 9385339863 | www.threatenedtaxa.org
Email: sanjay@threatenedtaxa.org

EDITORS

Founder & Chief Editor

Dr. Sanjay Molur

Wildlife Information Liaison Development (WILD) Society & Zoo Outreach Organization (ZOO),
Coimbatore, Tamil Nadu 641006, India

Assistant Editor

Dr. Chaithra Shree J., WILD/ZOO, Coimbatore, Tamil Nadu 641006, India

Managing Editor

Mr. B. Ravichandran, WILD/ZOO, Coimbatore, Tamil Nadu 641006, India

Associate Editors

Dr. Mandar Paingankar, Government Science College Gadchiroli, Maharashtra 442605, India

Dr. Ulrike Streicher, Wildlife Veterinarian, Eugene, Oregon, USA

Ms. Priyanka Iyer, ZOO/WILD, Coimbatore, Tamil Nadu 641006, India

Board of Editors

Dr. Russel Mittermeier

Executive Vice Chair, Conservation International, Arlington, Virginia 22202, USA

Prof. Mewa Singh Ph.D., FASC, FNA, FNASC, FNAPsy

Ramanna Fellow and Life-Long Distinguished Professor, Biopsychology Laboratory, and
Institute of Excellence, University of Mysore, Mysuru, Karnataka 570006, India; Honorary
Professor, Jawaharlal Nehru Centre for Advanced Scientific Research, Bangalore; and Adjunct
Professor, National Institute of Advanced Studies, Bangalore

Stephen D. Nash

Scientific Illustrator, Conservation International, Dept. of Anatomical Sciences, Health Sciences
Center, T-8, Room 045, Stony Brook University, Stony Brook, NY 11794-8081, USA

Dr. Fred Pluthero

Toronto, Canada

Dr. Priya Davidar

Sigur Nature Trust, Chadapatti, Mavinhalla PO, Nilgiris, Tamil Nadu 643223, India

Dr. John Fellowes

Honorary Assistant Professor, The Kadoorie Institute, 8/F, T.T. Tsui Building, The University of
Hong Kong, Pokfulam Road, Hong Kong

Prof. Dr. Mirco Solé

Universidade Estadual de Santa Cruz, Departamento de Ciências Biológicas, Vice-coordenador
do Programa de Pós-Graduação em Zoologia, Rodovia Ilhéus/Itabuna, Km 16 (45662-000)
Salobrinho, Ilhéus - Bahia - Brasil

Dr. Rajeev Raghavan

Professor of Taxonomy, Kerala University of Fisheries & Ocean Studies, Kochi, Kerala, India

English Editors

Mrs. Mira Bhojwani, Pune, India

Dr. Fred Pluthero, Toronto, Canada

Copy Editors

Ms. Usha Madgunaki, Zooreach, Coimbatore, India

Ms. Trisa Bhattacharjee, Zooreach, Coimbatore, India

Ms. Paloma Noronha, Daman & Diu, India

Web Development

Mrs. Latha G. Ravikumar, ZOO/WILD, Coimbatore, India

Typesetting

Mrs. Radhika, Zooreach, Coimbatore, India

Mrs. Geetha, Zooreach, Coimbatore, India

Fundraising/Communications

Mrs. Payal B. Molur, Coimbatore, India

Subject Editors 2021–2023

Fungi

Dr. B. Shivaraju, Bengaluru, Karnataka, India

Dr. R.K. Verma, Tropical Forest Research Institute, Jabalpur, India

Dr. Vatsavaya S. Raju, Kakatiya University, Warangal, Andhra Pradesh, India

Dr. M. Krishnappa, Jnana Sahyadri, Kuvempu University, Shimoga, Karnataka, India

Dr. K.R. Sridhar, Mangalore University, Mangalagangothri, Mangalore, Karnataka, India

Dr. Gunjan Biswas, Vidyasagar University, Midnapore, West Bengal, India

Dr. Kiran Ramchandra Ranadive, Annasaheb Magar Mahavidyalaya, Maharashtra, India

Plants

Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India

Dr. N.P. Balakrishnan, Ret. Joint Director, BSI, Coimbatore, India

Dr. Shonil Bhagwat, Open University and University of Oxford, UK

Prof. D.J. Bhat, Retd. Professor, Goa University, Goa, India

Dr. Ferdinando Boero, Università del Salento, Lecce, Italy

Dr. Dale R. Calder, Royal Ontario Museum, Toronto, Ontario, Canada

Dr. Cleofas Cervancia, Univ. of Philippines Los Baños College Laguna, Philippines

Dr. F.B. Vincent Florens, University of Mauritius, Mauritius

Dr. Merlin Franco, Curtin University, Malaysia

Dr. V. Irudayaraj, St. Xavier's College, Palayamkottai, Tamil Nadu, India

Dr. B.S. Kholia, Botanical Survey of India, Gangtok, Sikkim, India

Dr. Pankaj Kumar, Department of Plant and Soil Science, Texas Tech University, Lubbock, Texas, USA

Dr. V. Sampath Kumar, Botanical Survey of India, Howrah, West Bengal, India

Dr. A.J. Solomon Raju, Andhra University, Visakhapatnam, India

Dr. Vijayasankar Raman, University of Mississippi, USA

Dr. B. Ravi Prasad Rao, Sri Krishnadevaraya University, Anantpur, India

Dr. K. Ravikumar, FRLHT, Bengaluru, Karnataka, India

Dr. Aparna Watve, Pune, Maharashtra, India

Dr. Qiang Liu, Xishuangbanna Tropical Botanical Garden, Yunnan, China

Dr. Noor Azhar Mohamed Shazili, Universiti Malaysia Terengganu, Kuala Terengganu, Malaysia

Dr. M.K. Vasudeva Rao, Shiv Ranjani Housing Society, Pune, Maharashtra, India

Prof. A.J. Solomon Raju, Andhra University, Visakhapatnam, India

Dr. Mandar Datar, Agharkar Research Institute, Pune, Maharashtra, India

Dr. M.K. Janarthanam, Goa University, Goa, India

Dr. K. Karthigeeyan, Botanical Survey of India, India

Dr. Errol Vela, University of Montpellier, Montpellier, France

Dr. P. Lakshminarasimhan, Botanical Survey of India, Howrah, India

Dr. Larry R. Noblick, Montgomery Botanical Center, Miami, USA

Dr. K. Haridasan, Pallavur, Palakkad District, Kerala, India

Dr. Analinda Manila-Fajard, University of the Philippines Los Baños, Laguna, Philippines

Dr. P.A. Sinu, Central University of Kerala, Kasaragod, Kerala, India

Dr. Afroz Alam, Banasthali Vidyapith (accredited A grade by NAAC), Rajasthan, India

Dr. K.P. Rajesh, Zamorin's Guruvayurappan College, GA College PO, Kozhikode, Kerala, India

Dr. David E. Boufford, Harvard University Herbaria, Cambridge, MA 02138-2020, USA

Dr. Ritesh Kumar Choudhary, Agharkar Research Institute, Pune, Maharashtra, India

Dr. A.G. Pandurangan, Thiruvananthapuram, Kerala, India

Dr. Navendu Page, Wildlife Institute of India, Chandrabani, Dehradun, Uttarakhand, India

Dr. Kannan C.S. Warriar, Institute of Forest Genetics and Tree Breeding, Tamil Nadu, India

Invertebrates

Dr. R.K. Avasthi, Rohtak University, Haryana, India

Dr. D.B. Bastawade, Maharashtra, India

Dr. Partha Pratim Bhattacharjee, Tripura University, Suryamaninagar, India

Dr. Kailash Chandra, Zoological Survey of India, Jabalpur, Madhya Pradesh, India

Dr. Ansie Dippenaar-Schoeman, University of Pretoria, Queenswood, South Africa

Dr. Rory Dow, National Museum of Natural History Naturalis, The Netherlands

Dr. Brian Fisher, California Academy of Sciences, USA

Dr. Richard Gallon, Llandudno, North Wales, LL30 1UP

Dr. Hemant V. Ghate, Modern College, Pune, India

Dr. M. Monwar Hossain, Jahangirnagar University, Dhaka, Bangladesh

For Focus, Scope, Aims, and Policies, visit https://threatenedtaxa.org/index.php/JoTT/aims_scope

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions>

For Policies against Scientific Misconduct, visit https://threatenedtaxa.org/index.php/JoTT/policies_various

continued on the back inside cover

Cover: Nile Crocodile *Crocodylus niloticus* regulating body temperature on a warm day. Digital art on Procreate by © Aakanksha Komanduri.



An evasive naticid surfaces in India: first confirmed report of *Gennaeosinum perobliquum* (Dautzenberg & Fischer, 1907) (Gastropoda: Naticidae)

Aparna Mishra¹ , Sanjaya Dalai² , Roberto Ardovini³ , N.V. Subba Rao⁴  & Dipti Raut⁵ 

^{1,2,5} Centre of Excellence in Environment and Public Health, Environmental Science Laboratory, Department of Zoology, Ravenshaw University, Cuttack, Odisha 753003, India.

³ Department of Information Technology, ITIS FERMI ROMA Technical Institute, Rome 00135, Italy.

⁴ Molluscan Section, Zoological Survey of India, New Alipore, Kolkata, West Bengal 700053, India.

¹aparnamishra839@gmail.com, ²s.dalai1996@gmail.com, ³roberto_ardovini@fastwebnet.it, ⁴subbarao_nalluri@hotmail.com, ⁵diptiraut@ravenshawuniversity.ac.in (corresponding author)

Abstract: This study provides the first record of the naticid species *Gennaeosinum perobliquum* in Indian waters. Previously documented from Vietnam, Indonesia, and South Africa, *G. perobliquum* was collected during a benthic survey, from the intertidal zone of Chandipur, Balasore, Odisha, on the eastern coast of India. A clear description of the shell morphology is provided, along with detailed morphometric and imaging analyses. By comparing *G. perobliquum* with related naticid species found in India, this work aims to refine taxonomic understanding and contribute to knowledge of naticid biodiversity in the region. This newly documented record not only expands the known geographic distribution of *G. perobliquum* but also underscores the importance of continued benthic faunal surveys for further exploration of molluscan biodiversity in the Bay of Bengal and adjacent regions.

Keywords: Bay of Bengal, benthos, Chandipur, first record, geographic distribution, Indian waters, intertidal, mollusc biodiversity, moon shells, Odisha.

Abbreviation: TSL—Total Shell Length | WB—Width of the last whorl | HB—Height of the last whorl | SL—Spire Length | CL—Columella Length | UL—Length of the Umbilical callus | (AW)_a—Aperture Width towards anterior side | (AW)_p—Aperture Width towards posterior side | dd—Dead specimen | RZEVBU—Ravenshaw University Zoology Environmental Science Budhabalanga Estuary | ZSI M—Zoological Survey of India Mollusca.

Editor: Han Raven, Naturalis Biodiversity Center, Leiden, Netherlands.

Date of publication: 26 June 2026 (online & print)

Citation: Mishra, A., S. Dalai, R. Ardovini, N.V.S. Rao & D. Raut (2026). An evasive naticid surfaces in India: first confirmed report of *Gennaeosinum perobliquum* (Dautzenberg & Fischer, 1907) (Gastropoda: Naticidae). *Journal of Threatened Taxa* 18(6): 29120–29126. <https://doi.org/10.11609/jott.10328.18.6.29120-29126>

Copyright: © Mishra et al. 2026. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by providing adequate credit to the author(s) and the source of publication.

Funding: The first author AM acknowledges the financial research support provided by the Council of Scientific & Industrial Research, New Delhi, India (File no. 09/1036(24086)/2025EMR-1 dated 13/10/2025).

Competing interests: The authors declare no competing interests.

Author details: APARNA MISHRA: a Ph.D research scholar with research interest in benthic ecology and focus on molluscan taxonomy. SANJAYA DALAI: a Ph.D research scholar with research interest in benthic ecology and focus on molluscan taxonomy. ROBERTO ARDOVINI: professor at ITIS Fermi Roma (Technical Institute of Information Technology, Rome), specializing in malacology and conchology, with a focus on the taxonomy and description of marine gastropods and bivalves. His work includes descriptions of new species and contributions to global shell inventories. N.V. SUBBA RAO: scientist and former head of the Molluscan Section, Zoological Survey of India. Expertise in the taxonomy, systematics, and faunal documentation of Indian molluscs across marine, freshwater, and terrestrial habitats. Landmark works are foundational reference material for Indian malacology. DIPTI RAUT: professor of Zoology at Ravenshaw University, Cuttack, Odisha, India. Research focuses on macrobenthic community ecology. Work integrates environmental monitoring with marine biodiversity documentation.

Author contribution: Conceptualization: Dipti Raut, N. V. Subba Rao; Methodology: Dipti Raut, Aparna Mishra, Sanjaya Dalai; Data Collection: Sanjaya Dalai, Aparna Mishra, Roberto Ardovini; Data Analysis: Aparna Mishra, Sanjaya Dalai, Roberto Ardovini; Writing Original Draft: Aparna Mishra, Sanjaya Dalai; Writing Review and Editing: Dipti Raut, N. V. Subba Rao, Roberto Ardovini; Supervision: Dipti Raut

Acknowledgments: The first author AM acknowledging the financial research support provided by the Council of Scientific & Industrial Research, New Delhi, India (File no. 09/1036(24086)/2025EMR-1 dated 13/10/2025). The authors are thankful to the Department of Zoology, Ravenshaw University, Cuttack, Odisha, India, for the laboratory facilities. The anonymous reviewers' comments greatly improved the manuscript.



INTRODUCTION

Members of the cosmopolitan family Naticidae within the phylum Mollusca and class Gastropoda inhabit a wide range of environments, from intertidal to the deep sea. Commonly called moon shells, these oval-shaped predatory gastropods are known for their distinctive drilling behavior, which they use to prey on various bivalves and gastropods. They typically dwell in sandy and muddy substrates, where they remain hidden (Huelsken et al. 2008). The Naticidae family includes four subfamilies, 38 genera, and about 260–270 species (Kabat 1996; WoRMS 2025).

The genus *Gennaeosinum* Iredale, 1929, comprises three valid species. Kilburn (1976) classified it as a subgenus of *Eunaticina*; *Gennaeosinum* is consistently characterized by a thick funicle, a feature absent in all *Eunaticina* species. Although 36 Naticidae species across 12 genera have been documented in India, no *Gennaeosinum* species has been reported from the region (Tripathy et al. 2024). This study documents the first record of *Gennaeosinum perobliquum* (Dautzenberg & Fischer, 1907) in the Bay of Bengal, expanding its known distribution. *Gennaeosinum perobliquum* was described initially from “Ben-Son” (probably Binh Son, south of Da Nang) in Vietnam (Dautzenberg & Fischer 1907) and later recorded from Durban, Kwazulu-Natal, South Africa (Kilburn 1976) and Indonesia (Bakker et al. 2026). This species is often misidentified as *Eunaticina papilla* (Gmelin, 1791) due to similar spiral lirae patterns. As detailed here, *G. perobliquum* can be reliably distinguished from *Eunaticina* species occurring in India.

A recent study (Mishra et al. 2024) identified the naticid *Sinum laevigatum* (Lamarck, 1822) along Odisha’s coast, underscoring the need for further research on Naticidae in this region. Consequently, extensive surveys were conducted along Odisha’s entire coastline, with particular focus on the intertidal zone of Chandipur, where specimens were collected.

This study not only adds *G. perobliquum* to the list of species known in Indian waters but also highlights the rich malacofaunal diversity present along the largely unexplored eastern coast of India. It provides a thorough examination of the specimen, including the holotype description and comparative analyses with closely related species, aiming to mitigate future taxonomic ambiguities. Enhanced photographic documentation and detailed morphometric measurements are included to enrich the understanding of shell sculpture and address existing gaps in the literature.

MATERIALS AND METHODS

A survey was conducted in the Chandipur intertidal region (21.455° N, 87.045° E) (Figure 1) during September 2025. Samples were randomly collected during low tide by hand-picking and stored; dead specimens were placed in ziplock pouches, and live specimens were preserved in 4% formalin. Dead specimens were carefully washed, air-dried, and sorted. Sediment samples were obtained through manual excavation using a hand shovel for the purpose of analyzing soil texture. Morphometric measurements were obtained using a dial caliper (Safeseed, China; resolution 0.1 mm, accuracy ± 0.2 mm), while shell surface structures were examined under a stereo microscope (Leica E24W, Germany). Specimen photographs were captured with a Nikon Z 30 camera (MC 50 mm / 2.8 f). Following vouchering, specimens were deposited in the Environmental Science Laboratory, Department of Zoology, Ravenshaw University, Cuttack, Odisha, India.

Morphological identification was performed based on the holotype description (Dautzenberg & Fischer 1907), supplemented by additional information from Kilburn (1976). A critical comparison was made with other naticids reported from India, as documented by Rao (2003). The taxonomic status was verified through the World Register of Marine Species (WoRMS 2025).

RESULT

Systematics

Family: Naticidae Guilding, 1834

Subfamily: Sininae Woodring, 1928

Genus: *Gennaeosinum* Iredale, 1929

Type species: *Gennaeosinum peleum* Iredale, 1929 (type by original designation)

Gennaeosinum perobliquum (Dautzenberg & H. Fischer, 1907) (Image 1A–H)

Chresonomy

Sigaretus (Eunaticina) perobliquus Dautzenberg & H. Fischer, 1907: p. 178, pl. 5, figures 4–5.

Eunaticina papilla (non Gmelin); Kensley, 1973: 93, fig. 323.

Eunaticina (Gennaeosinum) perobliqua (Dautzenberg & Fischer, 1907) - Kilburn 1976; p. 872, figure. 22 (c).

Eunaticina (Gennaeosinum) perobliqua (Dautzenberg & Fischer, 1907) - Bakker H, Gill A, Creuwels J (2026). Naturalis Biodiversity Center (NL) - Mollusca. Naturalis Biodiversity Center. Occurrence dataset <https://doi.org/10.21203/rs.3.rs-1234567/v1>




Figure 1. Map showing the sampling location of the Chandipur intertidal region, Odisha, eastern coast of India.

org/10.15468/yefvnc accessed via GBIF.org on 2026-05-05. <https://www.gbif.org/occurrence/2444340122>.

Type Locality: “Ben-Son” (probably Binh Son, south of Da Nang) in northern Vietnam

IUCN status: Not Evaluated.

Material examined: Two specimens (dd), Accession number: RZEB18 (TSL 13.5 mm, WB 10.2 mm); RZEB19 (TSL 13.9 mm, WB 10.5 mm) were examined from the Chandipur intertidal, Balasore, Odisha, eastern coast of India (21.455° N, 87.045° E), coll. Sanjaya Dalai & Aparna Mishra.

Additional material studied: Figured type material of *Sigaretus (Eunaticina) perobliquus* Dautzenberg & Fischer, 1907 (images examined from Dautzenberg & Fischer 1907; p. 178, pl. 5, figure 4–5).

Topotype of *Eunaticina papilla* (Gmelin, 1791) (Zoological Survey of India, Registration number ZSI M 22150/4).

Specimen of *Eunaticina linnaeana* (Récluz, 1843) (Zoological Survey of India, Kolkata, Registration number M25925/5).

Description: Shell solid, white, moderately elongated

oval; aperture strongly prosocline. Low, depressed spire with blunt tip; deep suture. Protoconch slightly convex, smooth, glossy white with brown tinge; two whorls. Transition to teleoconch not well defined; feeble spiral ribs present. Teleoconch with two whorls, last whorl highly convex much larger than previous whorls. Teleoconch surface initially smooth, later marked by strong arc-like growth lines; faint spiral ribs becoming distinct and unevenly distributed on last whorl. Around 9–10 spiral ribs on penultimate whorl, separated by shallow, broad interspaces. Ribs more elevated below suture, flattening with lower interspaces on last whorl. Spiral ribs become denser towards base; growth lines become stronger towards outer lip. Spiral threads continue into inner side of umbilical groove. Aperture semicircular, wider anteriorly. Inner lip almost straight, slight concavity towards parietal region; anterior region curved. Thick parietal callus extends over umbilicus, forming a funicle that covers umbilicus backwardly. Funicle divided by shallow groove. Wide open anteriorly, deep umbilical groove. Operculum corneous.

Remarks: A thick callus pad covers the umbilical




Figure 2. Distribution of *Gemmaeosinum perobliquum* in the Indo-West Pacific region (Black star: type locality; Red star: Present report; Blue circles: Natal and Indonesia).

Table 1. Morphometrics of *Gemmaeosinum perobliquum* collected from Odisha.

| Specimen | TSL | WB | HB | SL | CL | UL | (AW) _s | (AW) _p |
|----------|------|------|-----|-----|------|-----|-------------------|-------------------|
| RZEB18 | 13.5 | 10.2 | 7.8 | 3.8 | 11.3 | 4.0 | 6.8 | 3.9 |
| RZEB19 | 13.9 | 10.5 | 8.9 | 3.9 | 11.0 | 3.8 | 7.4 | 4.3 |

groove and the funicle, which has a shallow groove that divides it into two. These are the key identification features that confirm the first report of the genus *Gemmaeosinum* and the species *G. perobliquum* from India. The multi-spiral protoconch with two whorls signifies a planktonic veliger larval stage (Vendetti, 2007). This developmental form implies improved dispersal abilities; therefore a large distribution area is to be expected.

Distribution: From Vietnam (Holotype) through Indonesia (GBIF) and India to South Africa (Kilburn, 1976) (Figure 2).

Locality within India: Chandipur Intertidal, Balasore, Odisha, east coast of India, present study (Figure 2).

Habitat: The species has repeatedly been recorded

from shallow water. Chandipur’s intertidal stretch shows diverse sedimentary conditions, with a silty mudflat in the middle region. The upper intertidal region has a coarse, sandy substratum, encompassing high diversity.

DISCUSSION

Identification of Naticidae can be challenging because of their highly cryptic morphology and very minor interspecific variation (Sharma et al. 2021). In India, three genera of the Sininae subfamily are represented: *Eunaticina* P. Fischer, 1885, *Sigatica* O. Meyer & Aldrich, 1886, *Sinum* Röding, 1798. The genus *Gemmaeosinum* herein reported is the fourth,




Image 1. *Gemmaeosinum perobliquum*. RZEB18 (A–D): A—dorsal view | B—side view | C—ventral view | D—apical view. RZEB19 (E–H): E—dorsal view | F—ventral view | G—protoconch | H—callus pad. © Aparna Mishra and Sanjaya Dalai.

represented by a single species. The aforementioned three genera are unique in several ways. *Eunaticina* differs from *Sinum* by its auricular, more depressed shape and wider aperture. *Gennaeosinum* is similar to *Eunaticina* and was treated as its subgenus by Kilburn (1976). *Gennaeosinum* is differentiated from *Eunaticina* by a thick callus pad in the parietal region, which is absent in *Eunaticina*. *Gennaeosinum* includes three species: *G. intercisum* Iredale, 1931, *G. peleum* Iredale, 1929, and *G. perobliquum* (Dautzenberg & Fischer, 1907). Kilburn included several other species, namely *G. kraussi* (E.A. Smith, 1902) and *G. linguifera* Thiele, 1925, but now *G. kraussi* is accepted as *Eunaticina kraussi* whereas *G. linguifera* is accepted as *Natica Linguifera*. This work confirmed the first report of the *Gennaeosinum* genus and the species *G. perobliquum* from Indian marine waters. Previously, this species was found in the western and central Indo-Pacific regions. It now extends its range into the Bay of Bengal. There are few descriptions or images of the species besides the original description, which has excellent figures. Kilburn (1976) did not provide an illustration. The Indian specimens align with the original description and Kilburn's description of the features of the South African specimen.

The species can be confused with two *Eunaticina* species well known from the Indo-West Pacific: *E. papilla* (Gmelin, 1791) and *E. linnaeana* (Récluz, 1843). Kilburn (1976) compared *G. perobliquum* with *E. linnaeana*, but the specimens are more similar to *E. papilla*. *Eunaticina linnaeana* is larger and broadly ovate, while *G. perobliquum* is smaller and moderately ovate. Both have low spires, but the conical shape distinguishes the latter. *Eunaticina linnaeana* has fine spiral grooves, whereas *G. perobliquum* is ornamented with well-developed, flat spiral ribs. *Eunaticina linnaeana* has an aperture that is wider, elongated, and larger, giving it a distinct look compared to the semicircular opening in *G. perobliquum*. *Eunaticina linnaeana* has frequently been misidentified as *E. papilla*, leading to similar confusion with *G. perobliquum*. *E. papilla* has an exerted spire, shallow suture, and ovate aperture. *G. perobliquum* resembles *E. papilla* in size, spiral ribs, and moderately large aperture. Distinctively, *G. perobliquum* stands out from both *Eunaticina* species because it features a thick callus pad in the parietal region, divided into two by a shallow groove, a feature always absent in *Eunaticina*.

Various factors influence the dispersal of naticid species in Indian marine waters. Oceanic currents and ballast water from ships play a major role (Ruiz et al. 2000; Cowen & Sponaugle 2009; Treml et al. 2015). The Bay of Bengal, located in the northern Indian Ocean,

shares environmental and habitat characteristics with the central and western Indo-Pacific marine regions. These include high sea surface temperatures, a variety of coastal and open-water habitats, and rich biodiversity, all characteristic of tropical Indo-West Pacific ecosystems (Spalding et al. 2007). Climate change and rising sea temperatures are likely to shift the ecological niches of these species. This could extend their distribution beyond previously established ranges (Wallingford & Sorte 2022; Lin et al. 2024). Changes in species distribution may be hard to establish due to a lack of comprehensive surveys and challenges in species identification, especially given the cryptic nature of *Gennaeosinum perobliquum* (D'Souza & Shenoy 2023). Research on the Naticidae family in India has been limited, particularly along the eastern coast. This highlights the need for further exploration, including the collection of live specimens and the establishment of a DNA database to elucidate intraspecific and interspecific relationships among Indian naticid species.

CONCLUSION

The discovery of *G. perobliquum* underscores significant connections among molluscan fauna in the Indo-West Pacific and highlights the urgent need for a comprehensive taxonomic review of naticid species in India. This study also confirms the limited exploration of malacofaunal diversity along the eastern coast. Additionally, this record expands the regional species list and deepens the understanding of biogeographical patterns in the Indo-West Pacific. These findings are crucial for developing effective conservation strategies and for protecting biodiversity in the region.

REFERENCES

- Bakker, H., A. Gill & J. Creuwels (2026). Naturalis Biodiversity Center (NL) — Mollusca. Naturalis Biodiversity Center. Occurrence dataset <https://doi.org/10.15468/yefvnk>. Accessed on 05.x.2026. <https://www.gbif.org/occurrence/2444340122>.
- Cowen, R.K. & S. Sponaugle (2009). Larval dispersal and marine population connectivity. *Annual Review of Marine Science*, 1: 443–466.
- D'Souza, S.L. & K.B. Shenoy (2023). Marine molluscs of India—a review on their diversity and distribution. *Journal of Coastal Conservation* 27(6): 67. <https://doi.org/10.1007/s11852-023-00998-0>
- Dautzenberg, P. & H. Fischer (1907). Contribution à la faune malacologique de l'Indo-Chine. *Journal de Conchyliologie* 54(3): 145–226. <https://biodiversitylibrary.org/page/16298507>
- Huelsken, T., C. Marek, S. Schreiber, I. Schmidt & M. Hollmann (2008). The Naticidae (Mollusca: Gastropoda) of Giglio Island (Tuscany,

- Italy): Shell characters, live animals, and a molecular analysis of egg masses. *Zootaxa* 1770(1): 1–140. <https://doi.org/10.11646/zootaxa.1770.1.1>
- Kabat, A.R. (1996).** Biogeography of the genera of Naticidae (Gastropoda) in the Indo-Pacific. *American Malacological Bulletin* 12(1): 29–35.
- Kensley, B. (1973).** Sea-shells of Southern Africa. Gastropods. Maskew Miller, Cape Town, 225 pp.
- Kilburn, R.N. (1976).** A revision of the Naticidae of Southern Africa and Moçambique (Mollusca). *Annals of the Natal Museum* 22(3): 829–884.
- Lin, X.N., C.Y. Ma, L.S. Hu, M.L. Liao, L.X. Ma, P.R. Teske, A. Hoffmann & Y.W. Dong (2024).** Genomics-informed range predictions under global warming reveal reduced adaptive diversity whilst buffering range shifts for a marine snail. *Global Change Biology* 30(11): e17571. <https://doi.org/10.1111/gcb.17571>
- Mishra, A., S. Dalai, R. Behera, N.V.S. Rao & D. Raut (2024).** Rare Moon Shell *Sinum laevigatum* (Gastropoda: Naticidae): a new record from coastal waters of peninsular India. *Thalassas* 40: 653–657. <https://doi.org/10.1007/s41208-023-00652-4>
- Rao N.V.S. (2003).** Indian Seashells (Part-I): Polyplacophora and Gastropoda. Records of Zoological Survey of India Occasional Paper 192: 1–416.
- Ruiz, G.M., J.T. Carlton, E.D. Grosholz & A.H. Hines (2000).** Invasion of coastal marine communities in North America: apparent patterns, processes, and biases. *Annual Review of Ecology and Systematics* 31: 481–531.
- Sharma, N., S. Mondal, S.S. Das, K. Bose & S. Saha (2021).** Morphological conservatism of the family Naticidae (Gastropoda) through time: potential causes and consequences. *Paleobiology* 47(3): 487–502. <https://doi.org/10.1017/pab.2020.62>
- Spalding, M.D., H.E. Fox, G.R. Allen, N. Davidson, Z.A. Ferdaña, M. Finlayson, B.S. Halpern, M.A. Jorge, A. Lombana, S.A. Lourie, K.D. Martin, E. McManus, J. Molnar, C.A. Recchia & J. Robertson (2007).** Marine ecoregions of the world: a bioregionalization of coastal and shelf Areas. *BioScience* 57(7): 573–583. <https://doi.org/10.1641/B570707>
- Tremblay, E.A., J.J. Roberts, Y. Chao, P.N. Halpin, H.P. Possingham & C. Riginos (2015).** Reproductive output and duration of the pelagic larval stage determine population connectivity in the Indo-Pacific. *Biological Conservation* 187: 249–257.
- Tripathy, B., P.C. Tudu, A. Mukhopadhyay, S.K. Sajan, C.R. Sreeraj, S.D. Gurumayum, A. Ghosh, R. Sultana, J. Purushothaman & A. Siddique (2024).** Checklist of Fauna of India: Mollusca. Version 1.0. Zoological Survey of India. <https://doi.org/10.26515/Fauna/1/2023/Mollusca>
- Vendetti, J.E. (2007).** Protoconch comparative morphology in extinct and extant buccinid gastropods and its utility in paleobiogeography, systematics, and inferring larval mode. *Malacologist* 48: 1–5.
- Wallingford, P.D. & C.J.B. Sorte (2022).** Dynamic species interactions associated with the range-shifting marine gastropod *Mexacanthina lugubris*. *Oecologia* 198: 749–761. <https://doi.org/10.1007/s00442-022-05128-5>
- WoRMS Editorial Board (2025).** World Register of Marine Species. Available from <https://www.marinespecies.org> at VLIZ. Accessed on 10.x.2025. <https://doi.org/10.14284/170>



Mr. Jatishwor Singh Irungbam, Biology Centre CAS, Branišovská, Czech Republic.
Dr. Ian J. Kitching, Natural History Museum, Cromwell Road, UK
Dr. George Mathew, Kerala Forest Research Institute, Peechi, India
Dr. John Noyes, Natural History Museum, London, UK
Dr. Albert G. Orr, Griffith University, Nathan, Australia
Dr. Sameer Padhye, Katholieke Universiteit Leuven, Belgium
Dr. Nancy van der Poorten, Toronto, Canada
Dr. Kareen Schnabel, NIWA, Wellington, New Zealand
Dr. R.M. Sharma, (Retd.) Scientist, Zoological Survey of India, Pune, India
Dr. Manju Siliwal, WILD, Coimbatore, Tamil Nadu, India
Dr. G.P. Sinha, Botanical Survey of India, Allahabad, India
Dr. K.A. Subramanian, Zoological Survey of India, New Alipore, Kolkata, India
Dr. P.M. Sureshan, Zoological Survey of India, Kozhikode, Kerala, India
Dr. R. Varatharajan, Manipur University, Imphal, Manipur, India
Dr. Eduard Vives, Museu de Ciències Naturals de Barcelona, Terrassa, Spain
Dr. James Young, Hong Kong Lepidopterists' Society, Hong Kong
Dr. R. Sundararaj, Institute of Wood Science & Technology, Bengaluru, India
Dr. M. Nithyanandan, Environmental Department, La Ala Al Kuwait Real Estate. Co. K.S.C., Kuwait
Dr. Himender Bharti, Punjabi University, Punjab, India
Mr. Purnendu Roy, London, UK
Mr. Saito Motoki, The Butterfly Society of Japan, Tokyo, Japan
Dr. Sanjay Sondhi, TITLI TRUST, Kalpavriksh, Dehradun, India
Dr. Nguyen Thi Phuong Lien, Vietnam Academy of Science and Technology, Hanoi, Vietnam
Dr. Nitin Kulkarni, Tropical Research Institute, Jabalpur, India
Dr. Robin Wen Jiang Ngiam, National Parks Board, Singapore
Dr. Lionel Monod, Natural History Museum of Geneva, Genève, Switzerland.
Dr. Asheesh Shivam, Nehru Gram Bharti University, Allahabad, India
Dr. Rosana Moreira da Rocha, Universidade Federal do Paraná, Curitiba, Brasil
Dr. Kurt R. Arnold, North Dakota State University, Saxony, Germany
Dr. James M. Carpenter, American Museum of Natural History, New York, USA
Dr. David M. Claborn, Missouri State University, Springfield, USA
Dr. Kareen Schnabel, Marine Biologist, Wellington, New Zealand
Dr. Amazonas Chagas Júnior, Universidade Federal de Mato Grosso, Cuiabá, Brasil
Mr. Monsoon Jyoti Gogoi, Assam University, Silchar, Assam, India
Dr. Heo Chong Chin, Universiti Teknologi MARA (UiTM), Selangor, Malaysia
Dr. R.J. Shiel, University of Adelaide, SA 5005, Australia
Dr. Siddharth Kulkarni, The George Washington University, Washington, USA
Dr. Priyadarsanan Dharma Rajan, ATREE, Bengaluru, India
Dr. Phil Alderslade, CSIRO Marine And Atmospheric Research, Hobart, Australia
Dr. John E.N. Veron, Coral Reef Research, Townsville, Australia
Dr. Daniel Whitmore, State Museum of Natural History Stuttgart, Rosenstein, Germany.
Dr. Yu-Feng Hsu, National Taiwan Normal University, Taipei City, Taiwan
Dr. Keith V. Wolfe, Antioch, California, USA
Dr. Siddharth Kulkarni, The Hormiga Lab, The George Washington University, Washington, D.C., USA
Dr. Tomas Ditrich, Faculty of Education, University of South Bohemia in Ceske Budejovice, Czech Republic
Dr. Mihaly Foldvari, Natural History Museum, University of Oslo, Norway
Dr. V.P. Uniyal, Wildlife Institute of India, Dehradun, Uttarakhand 248001, India
Dr. John T.D. Caleb, Zoological Survey of India, Kolkata, West Bengal, India
Dr. Priyadarsanan Dharma Rajan, Ashoka Trust for Research in Ecology and the Environment (ATREE), Royal Enclave, Bangalore, Karnataka, India

Fishes

Dr. Topiltzin Contreras MacBeath, Universidad Autónoma del estado de Morelos, México
Dr. Heok Hee Ng, National University of Singapore, Science Drive, Singapore
Dr. Rajeesh Raghavan, St. Albert's College, Kochi, Kerala, India
Dr. Robert D. Sluka, Chiltern Gateway Project, A Rocha UK, Southall, Middlesex, UK
Dr. E. Vivekanandan, Central Marine Fisheries Research Institute, Chennai, India
Dr. Davor Zanella, University of Zagreb, Zagreb, Croatia
Dr. A. Biju Kumar, University of Kerala, Thiruvananthapuram, Kerala, India
Dr. Akhilesh K.V., ICAR-Central Marine Fisheries Research Institute, Mumbai Research Centre, Mumbai, Maharashtra, India
Dr. J.A. Johnson, Wildlife Institute of India, Dehradun, Uttarakhand, India
Dr. R. Ravinesh, Gujarat Institute of Desert Ecology, Gujarat, India

Amphibians

Dr. Sushil K. Dutta, Indian Institute of Science, Bengaluru, Karnataka, India
Dr. Annemarie Ohler, Muséum national d'Histoire naturelle, Paris, France

Reptiles

Dr. Gernot Vogel, Heidelberg, Germany
Dr. Raju Vyasa, Vadodara, Gujarat, India
Dr. Pritpal S. Soorae, Environment Agency, Abu Dhabi, UAE.
Prof. Dr. Wayne J. Fuller, Near East University, Mersin, Turkey
Prof. Chandrashekhar U. Rivonker, Goa University, Taleigao Plateau, Goa, India
Dr. S.R. Ganesh, Kalinga Foundation, Agumbe, India.
Dr. Himansu Sekhar Das, Terrestrial & Marine Biodiversity, Abu Dhabi, UAE

Journal of Threatened Taxa is indexed/abstracted in Bibliography of Systematic Mycology, Biological Abstracts, BIOSIS Previews, CAB Abstracts, EBSCO, Google Scholar, Index Copernicus, Index Fungorum, JournalSeek, National Academy of Agricultural Sciences, NewJour, OCLC WorldCat, SCOPUS, Stanford University Libraries, Virtual Library of Biology, Zoological Records.

NAAS rating (India) 5.64

Birds

Dr. Hem Sagar Baral, Charles Sturt University, NSW Australia
Mr. H. Byju, Coimbatore, Tamil Nadu, India
Dr. Chris Bowden, Royal Society for the Protection of Birds, Sandy, UK
Dr. Priya Davidar, Pondicherry University, Kalapet, Puducherry, India
Dr. J.W. Duckworth, IUCN SSC, Bath, UK
Dr. Rajah Jayapal, SACON, Coimbatore, Tamil Nadu, India
Dr. Rajiv S. Kalsi, M.L.N. College, Yamuna Nagar, Haryana, India
Dr. V. Santharam, Rishi Valley Education Centre, Chittoor Dt., Andhra Pradesh, India
Dr. S. Balachandran, Bombay Natural History Society, Mumbai, India
Mr. J. Praveen, Bengaluru, India
Dr. C. Srinivasulu, Osmania University, Hyderabad, India
Dr. K.S. Gopi Sundar, International Crane Foundation, Baraboo, USA
Dr. Gombobaatar Sundev, Professor of Ornithology, Ulaanbaatar, Mongolia
Prof. Reuven Yosef, International Birding & Research Centre, Eilat, Israel
Dr. Taej Mundkur, Wetlands International, Wageningen, The Netherlands
Dr. Carol Inskipp, Bishop Auckland Co., Durham, UK
Dr. Tim Inskipp, Bishop Auckland Co., Durham, UK
Dr. V. Gokula, National College, Tiruchirappalli, Tamil Nadu, India
Dr. Arkady Lelej, Russian Academy of Sciences, Vladivostok, Russia
Dr. Simon Dowell, Science Director, Chester Zoo, UK
Dr. Mário Gabriel Santiago dos Santos, Universidade de Trás-os-Montes e Alto Douro, Quinta de Prados, Vila Real, Portugal
Dr. Grant Connette, Smithsonian Institution, Royal, VA, USA
Dr. P.A. Azeez, Coimbatore, Tamil Nadu, India

Mammals

Dr. Giovanni Amori, CNR - Institute of Ecosystem Studies, Rome, Italy
Dr. Anwaruddin Chowdhury, Guwahati, India
Dr. David Mallon, Zoological Society of London, UK
Dr. Shomita Mukherjee, SACON, Coimbatore, Tamil Nadu, India
Dr. Angie Appel, Wild Cat Network, Germany
Dr. P.O. Nameer, Kerala Agricultural University, Thrissur, Kerala, India
Dr. Ian Redmond, UNEP Convention on Migratory Species, Lansdown, UK
Dr. Heidi S. Riddle, Riddle's Elephant and Wildlife Sanctuary, Arkansas, USA
Dr. Karin Schwartz, George Mason University, Fairfax, Virginia.
Dr. Lala A.K. Singh, Bhubaneswar, Orissa, India
Dr. Mewa Singh, Mysore University, Mysore, India
Dr. Paul Racey, University of Exeter, Devon, UK
Dr. Honnavalli N. Kumara, SACON, Anaikatty P.O., Coimbatore, Tamil Nadu, India
Dr. Nishith Dharaiya, HNG University, Patan, Gujarat, India
Dr. Spartaco Gippoliti, Socio Onorario Società Italiana per la Storia della Fauna "Giuseppe Altobello", Rome, Italy
Dr. Justus Joshua, Green Future Foundation, Tiruchirappalli, Tamil Nadu, India
Dr. H. Raghuram, Sri S. Ramasamy Naidu Memorial College, Virudhunagar, Tamil Nadu, India
Dr. Paul Bates, Harison Institute, Kent, UK
Dr. Jim Sanderson, Small Wild Cat Conservation Foundation, Hartford, USA
Dr. Dan Challender, University of Kent, Canterbury, UK
Dr. David Mallon, Manchester Metropolitan University, Derbyshire, UK
Dr. Brian L. Cypher, California State University-Stanislaus, Bakersfield, CA
Dr. S.S. Talmale, Zoological Survey of India, Pune, Maharashtra, India
Prof. Karan Bahadur Shah, Budhanilakantha Municipality, Kathmandu, Nepal
Dr. Susan Cheyne, Borneo Nature Foundation International, Palangkaraja, Indonesia
Dr. Hemanta Kafley, Wildlife Sciences, Tarleton State University, Texas, USA

Other Disciplines

Dr. Aniruddha Belsare, Columbia MO 65203, USA (Veterinary)
Dr. Mandar S. Paingankar, University of Pune, Pune, Maharashtra, India (Molecular)
Dr. Jack Tordoff, Critical Ecosystem Partnership Fund, Arlington, USA (Communities)
Dr. Ulrike Streicher, University of Oregon, Eugene, USA (Veterinary)
Dr. Hari Balasubramanian, EcoAdvisors, Nova Scotia, Canada (Communities)
Dr. Rayanna Hellem Santos Bezerra, Universidade Federal de Sergipe, São Cristóvão, Brazil
Dr. Jamie R. Wood, Landcare Research, Canterbury, New Zealand
Dr. Wendy Collinson-Jonker, Endangered Wildlife Trust, Gauteng, South Africa
Dr. Rajeshkumar G. Jani, Anand Agricultural University, Anand, Gujarat, India
Dr. O.N. Tiwari, Senior Scientist, ICAR-Indian Agricultural Research Institute (IARI), New Delhi, India
Dr. L.D. Singla, Guru Angad Dev Veterinary and Animal Sciences University, Ludhiana, India
Dr. Rupika S. Rajakaruna, University of Peradeniya, Peradeniya, Sri Lanka
Dr. Bahar Baviskar, Wild-CER, Nagpur, Maharashtra 440013, India

Reviewers 2021–2023

Due to paucity of space, the list of reviewers for 2021–2023 is available online.

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Print copies of the Journal are available at cost. Write to:
The Managing Editor, JoTT,
c/o Wildlife Information Liaison Development Society,
3A2 Varadarajulu Nagar, FCI Road, Ganapathy, Coimbatore,
Tamil Nadu 641006, India
ravi@threatenedtaxa.org & ravi@zooreach.org



www.threatenedtaxa.org

OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

June 2026 | Vol. 18 | No. 6 | Pages: 29003–29158

Date of Publication: 26 June 2026 (Online & Print)

DOI: 10.11609/jott.2026.18.6.29003-29158

Articles

Asiatic Elephant conservation as a driver of forest carbon stock stabilization and avoided degradation in India

– Tarun Kathula & Tanu Jindal, Pp. 29003–29009

Genetic polymorphism of Dhofar Toad *Firouzophrynus dhufarensis* (Parker, 1931) (Amphibia: Bufonidae) across central Saudi Arabia

– Rawan Al-Shehri, Mohammed F. Albeshri & Ehab Eid, Pp. 29010–29019

Fish diversity in selected urban, suburban, and rural wetlands of Vellore District, Tamil Nadu, India

– Annie Pushpa Isaac, Sherrie Jesulyn David, Deepak Samuel Vijay Kumar & Nirmal Magadalenal Nathaniel, Pp. 29020–29035

Macrofungal species richness, composition, distribution, and ecological preference along the elevation gradient in Agasthyamala Biosphere Reserve, southern Western Ghats, India

– Kurunna Kandy Akshaya, Arumugam Karthikeyan, Arunachalam Rajasekaran, Binai Nagarajan & Cheravengat Kunhikannan, Pp. 29036–29051

Efficacy of 5% neem seed kernel extract against ectoparasites in six captive wildlife species at Rajiv Gandhi Zoological Park, India

– S.B. Kendre, P.D. Pawar, R.V. Jadhav, U.M. Tumlam, A.Y. Doiphode, V.G. Nimbalkar, P.K. Bhangale, V.C. Priyal & S.M. Meshram, Pp. 29052–29066

A comparative web-traffic analysis of three renowned wildlife conservation organisations - International Union for Conservation of Nature (IUCN), Wildlife Conservation Society (WCS) and World Wide Fund for Nature (WWF)

– Saswat Pati & V. Vijay Kumar, Pp. 29067–29078

Communications

First photographic record of the Himalayan Red Panda *Ailurus fulgens* (Mammalia: Carnivora: Ailuridae) in Yordi Rabe Supse Wildlife Sanctuary, Arunachal Pradesh, India

– Yomto Mayi, Shantabala Devi Gurumayum & Salvador Lyngdoh, Pp. 29079–29084

Lotus *Nelumbo* cultivations of Beehama Ganderbal offer novel habitats for diversity and seasonal variation of wetland birds

– Sheikh Tanveer Salam, Fayaz Ahmad Ahanter & Showkat Ahmad Wani, Pp. 29085–29092

Mating behavior of the Oaxacan Oak Anole *Anolis quercorum* (Squamata: Sauria: Anolidae) on a shade coffee plantation in Sierra Madre del Sur of Oaxaca, Mexico

– Jesús García Grajales, Alejandra Buenrostro Silva, Gibran Aldair Amador Larios, Diana Andrea Nieves Rocha & Ixil Pineda Ibarra, Pp. 29093–29097

Evidence for the local extirpation of the Dehradun Stream Frog *Amolops chakrataensis* Ray, 1992 from the type locality, Chakrata in western Himalaya, India, and associated threats: a call for urgent conservation action

– Vishal Kumar Prasad, Kumudani Bala Gautam, Devendra Singh, Amit Badola, Abhilasha Shrivastava, K.P. Dinesh & Amaël Borzée, Pp. 29098–29105

First record of the genus *Berlandina* Dalmas, 1922 (Araneae: Gnaphosidae) from India, with notes on *B. plumalis* (O. Pickard-Cambridge, 1872) and its synonymy

– Subhash I. Parmar, Dhruv A. Prajapati & Pranav J. Pandya, Pp. 29106–29113

First record of leucosiid crab *Lyphira perplexa* Galil, 2009 (Decapoda: Brachyura: Leucosiidae) from the eastern coast of India in West Bengal

– Prabir Sahoo, Sagar Samanta, Avik Bhanja, Manas Das & Pijush Payra, Pp. 29114–29119

An evasive naticid surfaces in India: first confirmed report of *Gennaosinum perobliquum* (Dautzenberg & Fischer, 1907) (Gastropoda: Naticidae)

– Aparna Mishra, Sanjaya Dalai, Roberto Ardovali, N.V. Subba Rao & Dipti Raut, Pp. 29120–29126

Review

Taxonomic reassessment of *Ompok hypophthalmus* (Bleeker, 1846) (Actinopterygii: Siluriformes: Siluridae) in Indonesia with global implications

– Dinesh Nalage, Tejswini Sontakke, Ashwini Biradar, Vidya Pradhan & P.S. Kudnar, Pp. 29127–29132

Short Communications

Recent sighting of Black Baza *Aviceda leuphotes* Dumont, 1820 (Aves: Accipitriformes: Accipitridae) in Nandhaur Wildlife Sanctuary, Uttarakhand, India

– Prashant Kumar, Inder Singh Rautela, Chandan Kumar, Pawan Koranga, Deepak Dharmashktu & Kundan Kumar, Pp. 29133–20137

Rapid increase in artificial-substrate nesting by White-bellied Sea Eagle *Haliaeetus leucogaster* (Gmelin, 1788) (Aves: Accipitriformes: Accipitridae) in Tamil Nadu, India

– H. Byju, N. Raveendran & H. Maitreyi, Pp. 29138–29142

Notes

A photographic record of the Chinese Pangolin *Manis pentadactyla* (Linnaeus, 1758) (Mammalia: Pholidota: Manidae) from Pakyong District, Sikkim, India

– Prashanti Pradhan, Jampal Dorjee Bhutia, Prem Kumar Chhetri & Bharat Kumar Pradhan, Pp. 29143–29145

First camera-trap records of three wild carnivores from Corbett Tiger Reserve, India

– Mridula, Kamakshi S. Tanwar, Anurag Nashirabadkar, Sudip Banerjee, Anindita Bidisha Chatterjee, Shikha Bisht & Yadvendra V. Jhala, Pp. 29146–29149

Photographic record of the Eastern Bronzeback Tree Snake *Dendrelaphis cf. proarchos* (Wall, 1909) from Dudhwa Tiger Reserve, Uttar Pradesh, India

– Vipin Kapoor Sainy, Aqsa Jaseem, Rohit Ravi, Apoorva Gupta, H. Raja Mohan, R. Jagadeesh & Kirti Chaudhary, Pp. 29150–29153

***Rhododendron pendulum* (Ericaceae) from Singalila National Park: an addition to the flora of West Bengal, India**

– Sulaxana Baraily & Projjwal Chandra Lama, Pp. 29154–29158

Publisher & Host



Threatened Taxa