

MONOGRAPH

A key to the bats (Mammalia: Chiroptera) of South Asia

C. Srinivasulu, Paul A. Racey & Shahroukh Mistry



Journal of Threatened Taxa

Journal of Threatened Taxa

ISSN 0974-7907 (online) | 0974-7893 (print)

Monograph

Date of publication (online): 05 July 2010

Date of publication (print): 05 July 2010

ISBN 978-81-902319-0-9 (online) | 978-81-902319-1-3 (print)

Editor: Csorba Gabor**Manuscript details:**

Ms # o2352

Received 28 November 2009

Final revised received 14 April 2010

Finally accepted 19 April 2010

Citation: Srinivasulu, C., P.A. Racey & S. Mistry (2010). A key to the bats (Mammalia: Chiroptera) of South Asia. *Journal of Threatened Taxa* 2(7): 1001-1076.**Copyright:** © C. Srinivasulu, Paul A. Racey & Shahroukh Mistry 2010. Creative Commons Attribution 3.0 Unported License. JoTT allows unrestricted use of this article in any medium for non-profit purposes, reproduction and distribution by providing adequate credit to the authors and the source of publication.**Author Details****DR. C. SRINIVASULU** is an Assistant Professor of Zoology at University College of Science, Osmania University, India. He heads the research laboratory at Osmania University that focuses on biodiversity inventorying, conservation, ecology and animal taxonomy (including bats) with special reference to Eastern Ghats and Godavari River basin in Andhra Pradesh.**PROFESSOR PAUL A. RACEY** is a Regius Professor of Natural History (Emeritus) at University of Aberdeen, UK and a Visiting Professor at University of Exeter, UK. He is a renowned bat specialist and is Co-Chair, Bat Specialist Group of IUCN's Species Survival Commission and Vice-Chairman, Fauna and Flora International.**DR. SHAHROUKH MISTRY** is at the Biological Sciences Department, Butte College, California, USA and is a member of Scientific Advisory Board of Bat Conservation International, Austin, Texas, USA and Chair, Board of Directors of North American Society of Bat Researchers.**Author Contributions**

C. Srinivasulu did the ground work to prepare the key. C. Srinivasulu and Paul Racey worked on microchiroptera key, while C. Srinivasulu and Shahroukh Mistry worked on megachiroptera key. All the authors contributed equally in refining and finalizing the key.



A key to the bats (Mammalia: Chiroptera) of South Asia

C. Srinivasulu¹, Paul A. Racey² & Shahroukh Mistry^{3,4}

¹ Wildlife Biology Section, Department of Zoology, Osmania University, Hyderabad, Andhra Pradesh 500007, India

² Centre for Ecology and Conservation, School of BioSciences, University of Exeter, Tremough Campus, Penryn, TR10 9EZ, United Kingdom

³ Biology Department, Butte College, 3536 Butte Campus Drive, Oroville, CA 95965, USA

⁴ Biological Sciences, California State University, Chico, CA 95929, USA

Email: ¹ c.srinivasulu@osmania.ac.in, ²p.racey@abdn.ac.uk, ³mistrysh@butte.edu

Abstract: A checklist and dichotomous key to 128 species of bats known from South Asia including Afghanistan, India, Pakistan, Nepal, Bhutan, Bangladesh, Sri Lanka and Maldives is provided. Character matrices for families, genera and species are also included. This article also briefly reviews their distribution (both physiographic and country-wise), status and main identification characters.

Keywords: Checklist, Chiroptera, dichotomous key, diversity, Mammalia, South Asia

Table of Contents

| | |
|---|------|
| Acknowledgements | 1006 |
| Introduction | 1007 |
| Scope of this article | 1007 |
| Taxonomic composition and endemism of the bats of South Asia | 1007 |
| Characters used for identification of bats | 1007 |
| Brief notes on Bats of South Asia | 1011 |
| Family Pteropodidae | 1011 |
| Brief descriptions of the genera of family Pteropodidae present in South Asia | 1011 |
| Family Rhinopomatidae | 1013 |
| Brief description of the genus of the family Rhinopomatidae present in South Asia | 1013 |
| Family Emballonuridae | 1013 |
| Brief descriptions of the genera of the family Emballonuridae present in South Asia | 1013 |
| Family Megadermatidae | 1013 |
| Brief description of the genus of the family Megadermatidae present in South Asia | 1013 |
| Family Rhinolophidae | 1014 |
| Brief description of the genus of the family Rhinolophidae present in South Asia | 1014 |
| Family Hipposideridae | 1014 |
| Brief descriptions of the genera of the family Hipposideridae present in South Asia | 1014 |
| Family Molossidae | 1015 |
| Brief descriptions of the genera of the family Molossidae present in South Asia | 1015 |
| Family Vespertilionidae | 1015 |

| | |
|--|------|
| Brief descriptions of the genera of the family Vespertilionidae present in South Asia | 1016 |
| Family Miniopteridae | 1018 |
| References | 1019 |
| Appendix I. Some important literature consulted to prepare the key | 1019 |
| Table 1. Classical and modern classification of bats | 1020 |
| Table 2. Checklist of bats of South Asia | 1021 |
| Key to the Suborders and the Families | 1023 |
| Keys to the Genera and Species of Bats of South Asia | 1024 |
| Key 1, Family Pteropodidae (13 species) | 1026 |
| Key 2 Family Megadermatidae (2 species) | 1026 |
| Key 3, Family Rhinopomatidae (3 species) | 1026 |
| Key 4, Family Rhinolophidae (20 species) | 1027 |
| Key 5, Family Hipposideridae (15 species) | 1029 |
| Key 6, For subfamilies of the family Vespertilionidae (63 species) | 1031 |
| Key 6A, Subfamily Murininae (8 species) | 1032 |
| Key 6B, Subfamily Kerivoulinae (3 species) | 1033 |
| Key 6C, Subfamily Myotinae (14 species) | 1033 |
| Key 6D, For tribes of the subfamily Vespertilioninae (38 species) | 1035 |
| Key 6D i, Tribe Plecotini (5 species) | 1036 |
| Key 6D ii, Tribe Nycticeiini (4 species) | 1037 |
| Key 6D iii, Tribe Eptesicini (9 species) | 1038 |
| Key 6D iv, Tribe Pipistrellini (12 species) | 1039 |
| Key 6D v, Tribe Vespertilionini (8 species) | 1041 |
| Key 7, Family Miniopteridae (3 species) | 1042 |
| Key 8, Family Emballonuridae (6 species) | 1043 |
| Key 9, Family Molossidae (4 species) | 1043 |
| Table 3. Diagnostic morphological characters of eight genera of the family Pteropodidae present in South Asia | 1044 |
| Table 3.1. Diagnostic morphological characters of the two species of the genus <i>Rousettus</i> Gray, 1821 present in South Asia | 1045 |
| Table 3.2. Diagnostic morphological characters of four species of the genus <i>Pteropus</i> Brisson, 1762 present in South Asia | 1045 |
| Table 3.3. Diagnostic morphological characters of the two species of the genus <i>Cynopterus</i> Cuvier, F., 1824 present in South Asia ... | 1046 |
| Table 3.4. Diagnostic morphological characters of the species of the genera <i>Megaerops</i> Peters, 1865; <i>Latidens</i> Thonlongya, 1972; <i>Sphaerias</i> Miller, 1906; <i>Eonycteris</i> Dobson, 1873; and <i>Macroglossus</i> Cuvier, F., 1824 present in South Asia | 1046 |
| Table 4. Diagnostic morphological characters of the genera of the families Rhinopomatidae, Emballonuridae, Megadermatidae and Rhinolophidae present in South Asia | 1047 |
| Table 4.1. Diagnostic morphological characters of the three species of the genus <i>Rhinopoma</i> E. Geoffroy, 1818 present in South Asia | 1048 |

| | |
|--|------|
| Table 4.2. Diagnostic morphological characters of five species of the genera <i>Taphozous</i> E. Geoffroy, 1818 and <i>Saccoaimus</i> Lesson, 1842 present in South Asia | 1049 |
| Table 4.3. Diagnostic morphological characters of the two species of the genus <i>Megaderma</i> E. Geoffroy, 1810 present in South Asia | 1048 |
| Table 4.4. Diagnostic morphological characters of twenty species of the genus <i>Rhinolophus</i> Lacépède, 1799 present in South Asia .. | 1050 |
| Table 5. Diagnostic morphological characters of the genera of the family Hipposideridae present in South Asia | 1054 |
| Table 5.1. Diagnostic morphological characters of twelve species of the genus <i>Hipposideros</i> Gray, 1831 present in South Asia... | 1055 |
| Table 5.2. Diagnostic morphological characters of one species each of the genera <i>Triaenops</i> Dobson, 1871; <i>Asellia</i> Gray, 1838; and <i>Ceolops</i> Blyth, 1848 present in South Asia | 1057 |
| Table 6. Diagnostic morphological characters of the genera of the family Molossidae present in South Asia | 1058 |
| Table 6.1. Diagnostic morphological characters of the species of the genera <i>Tadarida</i> Rafinesque, 1814, <i>Chaerephon</i> Dobson, 1874 and <i>Otomops</i> Thomas, 1913 present in South Asia | 1058 |
| Table 7. Diagnostic morphological characters of the genera of the family Vespertilionidae present in South Asia | 1059 |
| Table 7.1. Diagnostic morphological characters of the species belonging to the genera <i>Harpiocephalus</i> , <i>Harpiola</i> and <i>Murina</i> present in South Asia | 1062 |
| Table 7.2. Diagnostic morphological characters of the species of the genus <i>Kerivoula</i> present in South Asia | 1063 |
| Table 7.3. Diagnostic morphological characters of the species belonging to the genus <i>Myotis</i> Kaup, 1829 present in South Asia | 1064 |
| Table 7.4. Diagnostic morphological characters of the species belonging to the genera <i>Plecotus</i> , <i>Barbastella</i> and <i>Otonycteris</i> present in South Asia | 1066 |
| Table 7.5. Diagnostic morphological characters of the species belonging to the genera <i>Scotoecus</i> , <i>Scotomanes</i> and <i>Scotophilus</i> present in South Asia | 1066 |
| Table 7.6. Diagnostic morphological characters of the species belonging to the genus <i>Arielulus</i> , <i>Hesperoptenus</i> and <i>Eptesicus</i> present in South Asia | 1067 |
| Table 7.7. Diagnostic morphological characters of the species belonging to the genera <i>Nyctalus</i> , <i>Scotozous</i> and <i>Pipistrellus</i> present in South Asia | 1068 |
| Table 7.8. Diagnostic morphological characters of species of the genera <i>Tylonycteris</i> , <i>Ia</i> , <i>Falsistrellus</i> , <i>Vespertilio</i> , <i>Philetor</i> and <i>Hypsugo</i> present in South Asia | 1069 |
| Table 8. Diagnostic morphological characters of one genus of the family Miniopteridae present in South Asia | 1070 |
| Table 8.1. Diagnostic morphological characters of species of the genera <i>Miniopterus</i> Bonaparte, 1837 present in South Asia | 1071 |

ACKNOWLEDGEMENTS

We heartily acknowledge all the bat field researchers and taxonomists from the region and elsewhere whose published materials we relied upon to prepare this key. Many thanks are due to Dr. Paul J.J. Bates and Malcolm Pearch, Harrison Zoological Institute, Sevenoaks, UK; Dr. Y.P. Sinha, Retired Scientist, Gangetic Plains Research Station, Zoological Survey of India, Patna; Dr. A.M. Hutson, IUCN/SSC/Chiroptera Specialist Group, East Sussex, UK; and Dr. M.S. Pradhan, Retired Scientist, Western Ghats Regional Station, Zoological Survey of India, Pune for their encouragement, advice, support, comments and supply of important publications. We thank Dr. Ramakrishna, Director, Zoological Survey of India, Kolkata for permission to study the chiropteran collection and Dr. S.S. Saha, Dr. T.P. Bhattacharyya and Mr. M.K. Ghosh for helping during the specimen studies. Dr. G. Marimuthu and Dr. K. Sripathi of Madurai Kamaraj University, their other colleagues and students deserve special thanks for their support during our visits to Madurai, especially during the South Asian Chiroptera CAMP Workshop. We also thank Dr. A. Madhavan, India; Dr. J.C. Daniel, Bombay Natural History Society, India; Dr. Manoj Muni, India; Dr. Wipula Yapa, University of Colombo, Sri Lanka; Mr. K.M. Swe, University of Yangon, Myanmar; Dr. T.K. Shrestha, Tribhuvan University, Nepal, and all other bat researchers, especially the CCINSA members, for sharing their knowledge. We thank Dr. Wilma Jogenuri and Dr. Priya Raman, University of Arizona, USA for helping with references; Ms. Sally Walker and Mr. Sanjay Molur, Zoo Outreach Organization, Coimbatore for encouragement; Ms. Binu Priya, Ms. Padma Priya, and Ms. J. Sheela of Zoo Outreach Organization who never once minded scores of untimely and numerous requests for references and other information. We thank Dr. Bhargavi Srinivasulu and Aditya Srinivasulu for help in the preparation of the manuscript and initial checks and Dr. Neil Furey, FFI Cambodia for last minute tweaking. We thank Dr. G. Csorba, Head, Mammal Collection, Department of Zoology, Hungarian Natural History Museum, Budapest for his very valuable suggestions and inputs. We are also grateful to field researchers in South Asia who tested the key from 2002 to 2009 and responded with positive comments to incorporate amendments. The first author acknowledges the post doctoral research grant by Council for Scientific and Industrial Research, New Delhi to work on fruit bat taxonomy in India.



INTRODUCTION

Of the rich diversity of vertebrate fauna, bats are unique in being the only group of mammals that, like birds, have sustained flight. One of the 26 mammalian orders, the Chiroptera includes 1117 species of bats world over in rather two unequal suborders - the Megachiroptera (consisting 186 species of Old World fruit bats in one family) and the Microchiroptera (consisting 931 species in 17 families) (Mickleburgh et al. 1992; Koopman 1993; Srinivasulu & Srinivasulu 2001; Hutson et al. 2001; Mickleburgh et al. 2002; Simmons 2005). Recent molecular phylogenetic studies challenged this traditional subdivision and proposed that the bats be subdivided into two new suborders (Table 1), Yinpterochiroptera (includes the families Pteropodidae, Rhinolophidae, Megadermatidae and Rhinopomatidae) and Yangochiroptera (includes all the remaining families) (Teeling et al. 2005). However, until the new suborders are widely accepted, we prefer to retain the traditional subdivision.

Bats are widely distributed and have been recorded throughout the world excepting the Antarctic and a few Oceanic Islands (Mickleburgh et al. 2002). Some of the bat families are widespread and are recorded from both the Old World and the New World. Others are restricted in their range and are recorded either only from the Old World or the New World. Of the 18 families of bats, eight families (Pteropodidae, Rhinopomatidae, Nycteridae, Megadermatidae, Rhinolophidae, Hipposideridae, Myzopodidae and Mystacinidae) are restricted to the Old World; six families (Noctilionidae, Phyllostomidae, Desmodontidae, Natalidae, Furipteridae and Thyropteridae) are restricted to the New World; and three families (Emballonuridae, Molossidae and Vespertilionidae) are found both in the Old and New Worlds (Mickleburgh et al. 2002; Simmons 2005).

Scope of this article

Although work on taxonomy and systematics of the region's bat diversity has been considerable (recently reviewed by Bates & Harrison (1997)), there exists no published material that could help bat researchers in easy identification of bats both in the field and in the laboratory or museum. The necessity of identifying living bats in the field and preserved specimens prompted the preparation of this key. During the Conservation Assessment and Management Plan (CAMP) Workshop on South Asian Bats, held at Madurai in southern India in January 2002, the need for such a key for the South Asian region was felt to be indispensable. Hence, after deliberation and seeking the opinions from fellow bat researchers, the work on the preparation of the key for field identification of all the known species from Afghanistan, India, Pakistan, Nepal, Bhutan, Bangladesh, Sri Lanka and Maldives was

undertaken by the first author. We consulted published literature in the process of preparation of this work (Appendix I).

The key, designed for use with a vernier caliper or a millimeter scale and a hand lens, is modified from Blanford (1888-1891), Corbet & Hill (1992), and Bates & Harrison (1997). Numerous other publications, listed in the Reference section, were also consulted. As some authorities suggest that keys may lead to some confusion, the present work also incorporates identification character matrices in tabular form for the ease of the user. In the character matrices the dental formula for each genus dealt with in this key is provided. The dental formula includes details of incisors, canines, premolars, molars of one side for both upper and lower jaw followed by total number of teeth present. Variations within genus are denoted by the numerical in parenthesis.

We encourage readers to contact the lead author with suggestions and recommendations so that the key may be kept current.

Taxonomic composition and endemism of the bats of South Asia:

A total of 128 species of bats are reported from South Asia (Table 2), with the microchiropterans being better represented than the megachiropterans (115 vs. 13 species). Among the Microchiroptera large number of species are from the family Vespertilionidae, followed by the families Rhinolophidae, Hipposideridae, Emballonuridae, Molossidae, Rhinopomatidae and Megadermatidae (Fig. 1). Of this diversity, 10 species, namely, *Pteropus faunulus* (Pteropodidae), *Pteropus melanotus* (Pteropodidae), *Latidens salimalii* (Pteropodidae), *Rhinolophus cognatus* (Rhinolophidae), *Rhinolophus mitratus* (Rhinolophidae), *Hipposideros durgadasi* (Hipposideridae), *Hipposideros hypophyllus* (Hipposideridae), *Myotis csorbai* (Vespertilionidae), *Eptesicus tatei* (Vespertilionidae) and *Harpiola grisea* (Vespertilionidae) are endemic to the region. Family-wise percent endemism is greatest in Pteropodidae (21.4) followed by Hipposideridae (13.3), Rhinolophidae (11.7 species) and Vespertilionidae (4.8).

Among the countries representing South Asia, India has more than 90% of the total bat diversity of this region, while others have less than 50% diversity (Fig. 2). Bhutan has 51%, Nepal has 40%, Pakistan has 33%, Bangladesh has 29%, Afghanistan has 28%, Sri Lanka has 23% and Maldives has 2% of the total bat diversity of South Asia.

Characters used for identification of bats

Identification of bats depends upon a series of external, cranial and dental measurements. Besides mensural characters many qualitative characters also help in easy identification, and wherever applicable we




Figure 1. Chiropteran diversity in South Asia




Figure 2. Number of bat species occurring in each country of South Asia

have included these. In this section, various external and cranial (including dental) measurements dealt in the present article as well as those useful in taxonomic studies have been detailed.

External (see Figs. 3a and 3b for details)

E (Ear length): from the lower border of the external auditory canal to the tip of the pinna, excluding hairs.

FA (Forearm length): taken with the wings folded, from the outer end of the elbow to the outer end of the wrist joint (or carpus).

HB (Head Body length): taken dorsally, from the tip of the snout to the base of the tail.

HF (Hindfoot length): from the outer end of the heel at the base of the calcar to the outer end of the longest digit, excluding hairs or claws.

TL (Tail length): from the base of the tail near the anal opening to tip of the tail.

Some other external measurements that are considered for taxonomic studies include:

3mt (Third metacarpal length): from the outer end of wrist joint (or carpus) to the distal outer end of the metacarpal.

1ph 3mt (Length of the first phalanx of the third metacarpal): from the proximal to the distal end of the first phalanx.

2ph 3mt (Length of the second phalanx of the third metacarpal): from the proximal to the distal end of the second phalanx.

(Note: Similarly, 4mt (length of the fourth metacarpal), 5mt (length of the fifth metacarpal), 1ph 4mt (Length of the first phalanx of the fourth metacarpal), and 2ph 4mt (Length of the second phalanx of the fourth metacarpal) is also taken into consideration.)

TIB (Tibia length): taken from the knee joint to the ankle.

Thumb (Thumb length): length of the first digit including metacarpal and phalanx excluding claw.

WSP (Wingspan): maximum spread of the wing from tip to tip taken with wings fully stretched.

Cranial (see Figs. 4 and 5 for details)

CBL (Condyllobasal length): from occipito-condyle to the anterior edge of alveolus of the anterior incisor.

CCL (Condylolanine length): from occipito-condyle to the anterior edge of alveolus of the canine.

CM^u (Maxillary toothrow): from the front of the upper canine to the back of the crown of the last upper molar.

CM_d (Mandibular toothrow): from the front of the lower canine to the back of the crown of the last lower molar.

GTL (Greatest length of the skull): from the extreme end of the anterior to the extreme end of the posterior parts of the skull.

M (Mandible length): from the extreme end of the condyle to the extreme end of the anterior of the mandible including the incisors.

M^u-M_d (Width across the last molars): taken from the outer borders of the crown of the last upper molars.

ZB (Zygomatic breadth): Greatest width of the skull across the zygomatic arches.

Some other dental and cranial measurements that are considered for taxonomic studies include:

BB (Breadth of the braincase): Greatest width of the brain case.

IC (Interorbital constriction): the narrowest width across the interorbital region.

RW (Rostral width): taken across the front of the orbits

at their most anterior point.

Some important terms defined

Antebrachial membrane: membrane in front of the arm extending between shoulder to forearm, wrist or thumb.

Antitragus: a lobe developed from the basal part of the outer margin of the ear.

Bicuspidate: a tooth possessing two cusps.

Bifid: a structure having two distal processes.

Calcar: a cartilaginous or bony spur like projection arising from the ankle that supports the interfemoral membrane.

Canine: a single tall and pointed tooth situated behind the incisors in each toothrow.

Cusp: a prominence or point on tooth.

Echolocation: navigation in flight by means of the echo of sound pulses.

Gular sac: a glandular pouch like structure in the skin of the throat.

Incisor: a front tooth situated in front of the canine tooth in each toothrow.

Interfemoral membrane: also uropatagium, a membrane extending between inner margins of the legs and the distal end of the body enclosing all or a part of the tail.

Jugal: also malar or the cheekbone, present in the middle of the zygomatic arch.

Lancet: present in the forms belonging to the genus *Rhinolophus*, it is the erect, subtriangular, posterior part of the nose leaf.

Mandible: the lower jaw composed of two bones, fused to different degrees.

Metacarpal: one of the long bones of the hand of the bat extending from the carpal bones to the proximal phalanx of the finger.

Molar: a posterior cheektooth.

Narial: pertaining to nasal region.




Figure 3a. Schematic diagram of a bat showing important external measurements




Figure 3b. Important external measurements of *Taphozous longimanus*

Noseleaf: a simple to complex structure derived from the skin around the nose in some bats.

Paranhinal glands: specialized sebaceous glands on the side of the muzzle.

Phalanx: (plural Phalanges) Digital bone of a finger or toe.

Premolar: a cheektooth in front of the first molar.

Radio-metacarpal pouch: a pocket on the ventral side




Figure 4. Dorsal view of the skull of *Pteropus giganteus*




Figure 5. Ventral view of the skull of *Pteropus giganteus*

of the wing extending between the radius and the fifth metacarpal in some bats.

Rostrum: the facial part of the skull in front of the orbits.


Sella: a median anterior projection of the noseleaf of the genus *Rhinolophus* (Fig. 6).

Tibia: the bone extending between the knee and the ankle.

Tragus: a cutaneous projection at the opening of the external ear.

Unicuspid: a tooth with single cusp.

Zygoma: (plural Zygomata) The arch of the cheek bone comprising part of squamosal at the base, jugal in centre and part of maxilla in front.

Figure 6. Lateral view of noseleaf of *Rhinolophus sinicus*

BRIEF NOTES ON BATS OF SOUTH ASIA

Order Chiroptera

Suborder Megachiroptera

Family Pteropodidae

Includes about 186 species of bats that feed chiefly on fruits, leaves, flowers and flower products. Distributed in Europe, Africa, Asia, Australia and Oceania (Mickleburgh et al. 2002; Simmons 2005). They have strong muzzle and jaws. Do not possess noseleaf or tragus. Eyes are large. They possess keen sense of smell. Ears simple (Fig. 7A). Tail small or absent, and proximal part of the caudal vertebrae included, the distal ones are free (Fig. 8A). Fourteen species belonging to eight genera have been reported from South Asia.

Brief descriptions of the genera of the family Pteropodidae present in South Asia:

Rousettus Gray, 1821 – Medium-sized fruit bats (FA – 75.0-90.0 mm) of rather heavier built and short tail (8.0-21.0 mm). Muzzle heavy and has deep emargination between the projecting nostrils. First digit has large claws while the second has smaller ones. Males have well-developed glandular hairs on the throat than the females. Echolocate for orientation within roost by clicking tongue against the roof of mouth. Rostrum moderately elongated. Two pairs of lower and upper incisors present. Two species – *R. aegyptiacus* (E. Geoffroy, 1810) and *R.*

leschenaultii (Desmarest, 1820) – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.1.


Pteropus Brisson, 1762 – Medium to large-sized fruit bats (FA – 110.0-209.0 mm) without tail. Patagium arises from sides of dorsum and the back of the second toe. Uropatagium less developed. Second digit has a small claw. Rostrum moderate. Two pairs of lower and upper incisors present. Four species – *P. giganteus* Brünnich, 1782, *P. hypomelanus* Temminck, 1853, *P. melanotus* Blyth, 1863 and *P. faunulus* Miller, 1902 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.2.

Cynopterus Cuvier, F., 1824 – Medium-sized fruit bats (FA – 57.0-79.0 mm) with short tail (2.0-19.0 mm) that is half enclosed within the interfemoral membrane. Muzzle short, and has deep emargination between the projecting nostrils. Both the first and second fingers have distinct claws. Rostrum short. Two pairs of lower and upper incisors present. Two species – *C. sphinx* (Vahl, 1797) and *C. brachyotis* (Müller, 1838) – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.3.

Megaerops Peters, 1865 – Medium-sized fruit bats (FA – 52.0-63.0 mm) without any external tail. Muzzle short, and has deep emargination between the projecting nostrils. Ears simple with broadly rounded tips. Interfemoral membrane narrow with dorsal medial parts hairy. Rostrum short. Two pairs of upper and one pair of lower incisors present. One species – *M. niphanae* Yenbutra & Felten, 1983 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.4.

Latidens Thonglongya, 1972 – Medium-sized fruit bats (FA – 66.0-69.0 mm) without tail. Muzzle long, and has deep emargination between the projecting nostrils. Ears simple and oval, with narrowly rounded tips. Interfemoral membrane with some hairs on upper and lower sides. Rostrum elongated and narrow. Only one pair of lower and upper incisors present. One species – *L. salimalii* Thonglongya, 1972 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.4.

Sphaerias Miller, 1906 – Medium-sized fruit bats (FA – 52.0-61.0 mm) without tail. Muzzle long, and has deep emargination between the projecting nostrils. Ears with well defined pale anterior margin, and with small triangular antitragal lobe. Interfemoral membrane very narrow and calcar absent. Rostrum long and narrow. Two pairs of lower and upper incisors present. One species

**Figure 7. Outline of ear and shape of tragus in select species of bats**

[A - *Pteropus giganteus*; B - *Rhinopoma microphyllum*; C - *Megaderma lyra*; D - *Rhinolophus luctus*; E - *Myotis* sp.; F - *Plecotus wardi*; G - *Pipistrellus pipistrellus*; H - *Miniopterus schreibersii*]

– *S. blanfordi* (Thomas, 1891) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.4.

Eonycteris Dobson, 1873 – Medium-sized fruit bats (FA – 66.0-78.0 mm) with well-developed tail (11.5-23.0 mm). Muzzle long and thin, and has deep emargination between the projecting nostrils. Only the thumb is clawed, second digit lacks claw. Ears are narrowly rounded. Interfemoral membrane very moderately broad, tail and calcar well-developed. A pair of large anal glands present. Tongue is sharply pointed and highly protrusible with well-developed unfringed filiform papillae at the tip. Rostrum long and narrow. Two pairs of lower and upper incisors present. There are usually eight palatal ridges. One species – *E. spelaea* (Dobson, 1871) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.4.

Macroglossus Cuvier, F., 1824 – Small to medium-

sized fruit bats (FA – 44.0-52.0 mm) without or with rudimentary tail (3.5-5.5 mm). Specialized for nectar feeding, muzzle long and narrow. The thumb and the second digit are clawed. Ears medium with narrowly rounded tips, and small antitragal lobes. Interfemoral membrane very narrow and is thickly haired. Tongue is sharply pointed and highly protrusible. Rostrum relatively long and narrow, braincase strongly deflected downwards. Two pairs of lower and upper incisors present. One species – *M. sobrinus* (K. Andersen, 1911) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 3 and those of the species belonging to it in Table 3.4.

Suborder Microchiroptera **Family Rhinopomatidae**

Includes five species of insectivorous bats. Distributed in dry regions of Africa and Asia (Mickleburgh et al. 2002; Simmons 2005). They have strong muzzle with thickened narial pads. Possess rudimentary noseleaf (a distinct




Figure 8. View of interfemoral membrane and tail in different families of bats
[A - Pteropodidae; B - Rhinolophidae, Hipposideridae, Vespertilionidae, Miniopteridae; C - Molossidae; D - Rhinopomatidae; E - Emballonuridae]

dermal ridge). Ears with simple tragus (Fig. 7B) and joined over the forehead by thin membrane. The second and the third digits of each wing have two distinct bony phalanges. Tail very long, and is only partly enclosed by a small interfemoral membrane (Fig. 8D). Three species belonging to a single genus is reported from South Asia.

Brief description of the genus of the family Rhinopomatidae present in South Asia:

Rhinopoma E. Geoffroy, 1818 – Small to medium-sized bats (FA – 46.0-74.0 mm) with long tail (49.0-78.0 mm). Tail mostly projecting free from the interfemoral membrane. One pair of upper and two pairs of lower incisors present. Other characters as outlined above. Three species – *R. microphyllum* (Brünnich, 1782), *R. hardwickii* Gray, 1837 and *R. muscatellum* Thomas, 1903 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 4 and those of the species belonging to it in Table 4.1.

Family Emballonuridae

Includes about 51 species of insectivorous bats with wide distribution in Americas, Europe, Africa, Asia, Australia and Oceania (Mickleburgh et al. 2002; Simmons 2005). They have strong muzzle, and lack noseleaf. Ears with simple tragus. The basal part of the tail is enclosed loosely in the interfemoral membrane and the tip pierces the upper surface of the membrane and lies free on the dorsal side (Fig. 8E). Wings long and narrow and the

second digit of each lack phalanges. Males usually have well-developed glands. Six species belonging to two genera are reported from South Asia.

Brief descriptions of the genera of the family Emballonuridae present in South Asia:

Taphozous E. Geoffroy, 1818 – Small to medium-sized bats (FA – 55.6-88.0 mm) with relatively medium sized and stout tail (20.0-46.0 mm). In some the chin is either naked or haired, and gular sac and/or glands on throat present or lacking. Some species have radio-metacarpal pouch on the wing. Muzzle simple, lacks noseleaf, and nostrils open forward. Rostrum moderately elongated. One pair of upper and two pairs of lower incisors present. Five species – *T. perforatus* E. Geoffroy, 1818, *T. longimanus* Hardwicke, 1825, *T. nudiventris* Cretzschmar, 1830-31, *T. melanopogon* Temminck, 1841 and *T. theobaldi* Dobson, 1872 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 4 and those of the species belonging to it in Table 4.2.

Saccopteryx Lesson, 1842 – Medium-sized bats (FA – 63.0-68.0 mm) with relatively medium sized and stout tail (21.0-35.0 mm). The chin is covered with short hairs. Gular sac on throat well-developed in males and less developed in females. Radio-metacarpal pouch absent. Muzzle simple, lacks noseleaf, and nostrils open forward. Rostrum moderately elongated. One pair of upper and two pairs of lower incisors present. One species – *S. saccolaimus* (Temminck, 1838) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 4 and those of the species belonging to it in Table 4.2.

Family Megadermatidae

Includes five species of insectivorous and carnivorous bats with distribution restricted to Africa, Asia and Australia (Mickleburgh et al. 2002; Simmons 2005). Ears are large and oval, joined over the forehead by a membrane. Tragus is distinctly bifid (Fig. 7C). They have strong muzzle and noseleaf is simple and erect. On each wing the second digit possess one phalanx and the third digit possess two phalanges. Tail absent. Two species belonging to one genus is reported from South Asia.

Brief description of the genus of the family Megadermatidae present in South Asia:

Megaderma E. Geoffroy, 1810 – Small to medium-sized bats (FA – 54.0-71.5 mm) without a tail. Upper incisors totally lacking and two pairs of lower incisors present. Other characters as outlined above. Two species – *M. spasma* Linnaeus, 1758 and *M. lyra* E. Geoffroy, 1810 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 4 and those of the species belonging to it in Table 4.3.

Family Rhinolophidae

Includes about 77 species of insectivorous bats with

wide distribution in Europe, Africa, Asia and Australia (Mickleburgh *et al.* 2002; Simmons 2005). They have short muzzle and complex noseleaf that essentially bears an erect posterior lancet, lower horizontal horseshoe surrounding the nostrils and a perpendicular median sella (Fig. 9A). Ears pointed with well-developed antitragus (Fig. 7D). Tragus absent. The tail (13.0-55.0 mm) is enclosed in the interfemoral membrane (Fig. 8B). Twenty species belonging to the single genus is reported from South Asia.

Brief description of the genus of the family Rhinolophidae present in South Asia:

Rhinolophus Lacépède, 1799 – Small to medium-sized bats (FA – 33.4-80.5 mm) with relatively medium sized tail (13.0-55.0 mm). One pair of upper and two pairs lower incisors present. Other characters as outlined above. Twenty species – *R. ferrumequinum* (Schreber, 1774), *R. hipposideros* (Bechstein, 1800), *R. affinis* Horsfield, 1823, *R. pusillus* Temminck, 1834, *R. trifoliatus* Temminck, 1834, *R. luctus* Temminck, 1835, *R. rouxii* Temminck, 1835, *R. lepidus* Blyth, 1844, *R. macrotis* Blyth, 1844, 1844 *R. mitratus* Blyth, 1844, *R. subbadius* Blyth, 1844, *R. pearsonii* Horsfield, 1851, *R. blasii* Peters, 1867, *R. yunanensis* Dobson, 1872, *R. mehelyi* Matschie, 1902, *R. beddomei* Andersen, 1905, *R. sinicus* Andersen, 1905, *R. cognatus* Andersen 1906, *R. bocharicus* Kastchenko and Akimov, 1917 and *R. shortridgei* K. Andersen, 1918 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 4 and those of the species belonging to it in Table 4.4.

Family Hipposideridae

Includes about 81 species of insectivorous bats with wide distribution in Africa, Asia, Australia and Oceania (Mickleburgh *et al.* 2002; Simmons 2005). They have short muzzle and complex noseleaf that bears a horizontal horseshoe surrounding the nostrils, often thrown into skin folds and associated leaflets (Fig. 9B). Intermediate leaf may or may not be clearly differentiated. Posterior leaf simple to complex. Sella and connecting process absent. The tail is well-developed and enclosed in the interfemoral membrane (Fig. 8B). Other characters vary at the generic level. Fifteen species belonging to four genera are reported from South Asia.

Brief descriptions of the genera of the family Hipposideridae present in South Asia:

Hipposideros Gray, 1831 – Small to medium-sized leaf-nosed bats (FA – 33.0-99.0 mm) with a medium-sized tail (20.0-64.0 mm). Noseleaf with anterior leaf with or without median emargination, an intermediate leaf and a posterior leaf (Fig. 8). Supplementary leaflets may be absent or present, if present may vary from 1 to 4 and in some species the last being much reduced. Ears pointed with large antitragus. Tragus absent. The interfemoral

membrane is broad and completely encloses the tail except the extreme tip. One pair of upper incisors and two pairs of lower incisors are present. Twelve species – *H. speoris* (Schneider, 1800), *H. diadema* (Geoffroy, E., 1813), *H. larvatus* (Horsfield, 1823), *H. armiger* (Hodgson, 1835), *H. fulvus* Gray, 1838, *H. galeritus* Cantor, 1846, *H. ater* Templeton, 1848, *H. lankadiva* Kelaart, 1850, *H. cineraceus* Blyth, 1853, *H. pomona* Andersen, 1908, *H. durgadasi*/Khajuria, 1970 and *H. hypophyllus* Kock & Bhat, 1994 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 5 and those of the species belonging to it in Table 5.1.

Triaenops Dobson, 1871 – Medium-sized leaf-nosed bats (FA – 48.3-54.0 mm) with a long tail (31.0-39.0 mm). Muzzle elongated and broad. Noseleaf with anterior leaf with a deep median emargination, and internarial septum greatly expanded. Intermediate and posterior leaves undifferentiated, and upper border bears three vertical pointed processes. Narial lappets present. A single pair of supplementary leaflets present. Ears small. Tragus absent. The interfemoral membrane is broad and completely encloses the tail except the extreme tip. A bony spicule at the base of the terminal phalanx of the third finger is characteristic. One pair of upper and two pairs of lower incisors are present. One species – *T. persicus* Dobson, 1871 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 5 and those of the species belonging to it in Table 5.2.

Asellia Gray, 1838 – Medium-sized leaf-nosed bats (FA – 50.1-52.3 mm) with a moderately small tail (25.0-28.0 mm). Muzzle elongated and narrow. Noseleaf with simple anterior leaf lacking any emargination, and internarial septum not expanded. Intermediate leaf is smooth and slightly protuberant. Posterior leaf divided into four shallow cells with three ill-defined septa. The upper margin has three vertical processes of which the central one is pointed while those on the sides are blunt. Narial lappets little developed. Two pairs of supplementary leaflets present. The interfemoral membrane is broad and completely encloses the tail except the extreme tip (ca. 3-5 mm). One pair of upper and two pairs of lower incisors are present. One species – *A. tridens* Geoffroy, E., 1813 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 5 and those of the species belonging to it in Table 5.2.

Coelops Blyth, 1848 – Small-sized leaf-nosed bats (FA – 37.8-42.0 mm) with a rudimentary tail (< 2.0 mm). Noseleaf distinct with anterior leaf – distinctly divided into two by deep emargination and also bearing two elongated narrow supplementary lappets projecting beyond the muzzle. Intermediate leaf has moderately developed median process. Posterior leaf has a single cell and a median process. Noseleaf is covered with hairs. Ears



Figure 9. Frontal view of noseleaf of a Rhinolophid [A] and a Hipposiderid [B] bats

broad and have large antitragal lobe. The interfemoral membrane is narrow and poorly developed. One pair of upper and two pairs of lower incisors are present. One species – *C. fritii* Blyth, 1848 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 5 and those of the species belonging to it in Table 5.2.

Family Molossidae

Includes about 100 species of insectivorous bats with wide distribution in Americas, Europe, Africa, Asia, Australia and Oceania (Mickleburgh et al. 2002; Simmons 2005). Lacks noseleaf. Tail stout and conspicuously projecting out of a narrow interfemoral membrane (Fig. 8C). Ears variable, usually fleshy could either be free or joined by membrane over the head. Tragus is rudimentary, while the antitragus is rudimentary to large. The upper lip often wrinkled. Four species belonging to three genera are reported from South Asia.

Brief descriptions of the genera of the family Molossidae present in South Asia:

Tadarida Rafinesque, 1814 – Medium-sized free-tailed bats (FA – 43.1-63.9 mm) with a medium-sized tail (30.0-54.8 mm). Ear, tail and other characters as outlined above. In some species ears are not joined over the forehead, while in some they are joined. Normally one pair of upper and two pairs of lower incisors are present, but one species has three pairs of lower incisors. Two species – *T. teniotis* (Rafinesque, 1814) and *T. aegyptiaca* (E. Geoffroy, 1818) – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 6 and those of the species belonging to it in Table

6.1.

Chaerephon Dobson, 1874 – Moderate-sized free-tailed bats (FA – 43.1-50.5 mm) with a relatively long tail (30.0-44.0 mm). Ears large and connected by a membrane over the forehead. Tragus quadrate and minute, and antitragus half oval, separated posteriorly by a deep notch. One pair of upper and two pairs of lower incisors are present. One species – *C. plicatus* (Buchanan, 1800) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 6 and those of the species belonging to it in Table 6.1.

Otomops Thomas, 1913 – Moderately large-sized free-tailed bats (FA – 63.0-67.0 mm) with a relatively long tail (41.0-49.0 mm). Ears large and connected by a membrane over the forehead. Tragus triangular and minute, and antitragus absent. One pair of upper and two to three pairs of lower incisors are present. One species – *O. wroughtoni* (Thomas, 1913) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 6 and those of the species belonging to it in Table 6.1.

Family Vespertilionidae

Includes more than 400 species of insectivorous bats with wide distribution in Americas, Europe, Africa, Asia, Australia and Oceania (Mickleburgh et al. 2002; Simmons 2005). Muzzle simple and lacks the noseleaf. Ears separate from each other, however in few species they are joined over the forehead. Tragus is well-developed (Fig. 7E, F, G, H). The shape of antitragus can also be diagnostic. There is a considerable variation in the number of teeth among the genera. The long tail

is completely enclosed in the interfemoral membrane or the extreme tip protrudes out (Fig. 8B). Sixty-three species belonging to twenty three genera are reported from South Asia.

Brief descriptions of the genera of the family Vespertilionidae present in South Asia:

Murina Gray, 1842 – Small to medium-sized bats (FA – 27.7-40.9 mm) with a moderately long tail (27.0-41.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears short and broad with long and narrow tragus. Projecting tubular nostrils characteristic. Wings attached either to the base of the claw of the first toe or to the base of the toe. Two pairs of upper incisors and three pairs of lower incisors are present. Two pairs each of upper and lower premolars are present. Five species – *M. leucogaster* Milne-Edwards, 1872, *M. aurata* Milne-Edwards, 1872, *M. cyclotis* Dobson, 1872, *M. huttoni* (Peters, 1872) and *M. tubinaria* (Scully, 1881) – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.1.

Harpiola Thomas, 1915 – Medium-sized bats (FA – 32.4-32.8 mm) with a moderately long tail (~27.5 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears short and broad with triangular terminal half. Tragus long and narrow, acutely pointed and curved outwards. Projecting tubular nostrils characteristic. Wings attached to the base of the first toe. Two pairs of upper incisors and three pairs of lower incisors are present. Two pairs each of upper and lower premolars are present. One species – *H. grisea* (Peters, 1872) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.1.

Harpiocephalus Gray, 1842 – Small to medium-sized bats (FA – 44.1-50.1 mm) with a moderately long tail (40.0-50.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Muzzle short. Ears moderate with long tragus. Projecting tubular nostrils characteristic. Two pairs of upper incisors and three pairs of lower incisors are present. Two pairs each of upper and lower premolars are present. One species – *H. harpia* (Temminck, 1840) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and that of the species belonging to it in Table 7.1.

Kerivoula Gray, 1842 – Small-sized bats (FA – 31.5-42.0 mm) with a long tail (35.0-55.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears moderately long and funnel shaped with long and slender tragus. Muzzle simple. Two pairs of upper incisors and three pairs of lower incisors are present. Three pairs each of upper and lower premolars are present. Three species – *K. picta* (Pallas, 1767),

K. hardwickii (Horsfield, 1825) and *K. lenis* Thomas, 1916 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.2.

Myotis Kaup, 1829 – Small to medium-sized bats (FA – 30.0-58.3 mm) with a long tail (25.0-68.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears usually tall and slender, and occasionally large. Tragus well-developed and spear shaped (Fig. 7E). Two pairs of upper and three pairs of lower incisors are present. Some species have three pairs each of upper and lower premolars, while in some species consistently only two pairs of both upper and lower premolars present. Fourteen species – *M. emarginatus* (E. Geoffroy, 1806), *M. laniger* (Peters, 1871), *M. formosus* (Hodgson, 1835), *M. hasseltii* (Temminck, 1840), *M. horsfieldii* (Temminck, 1840), *M. muricola* (Gray, 1846), *M. siligorensis* (Horsfield, 1855), *M. blythii* (Tomes, 1857), *M. annectans* (Dobson, 1871), *M. nipalensis* (Dobson, 1871), *M. longipes* (Dobson, 1873), *M. montivagus* (Dobson, 1874), *M. sicarius* Thomas, 1915 and *M. csorbai* Topal, 1997 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.3.

Plecotus Geoffroy, E., 1818 – Small-sized bats (FA – 36.5-45.1 mm) with a long tail (48.0-54.0 mm). The interfemoral membrane encloses entire but the extreme tip of the tail. Ears usually large and joined over the forehead. Tragus well-developed and antitragus absent (Fig. 7F). Nostrils open upwards with their orifices extended backwards by a fissure. Two pairs of upper and three pairs of lower incisors are present. Two pairs of upper and three pairs of lower premolars are present. Three species – *P. homochrous* Hodgson, 1847, *P. wardi* Thomas, 1911 and *P. strelkovi* Spitzenberger, 2008 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.4.

Barbastella Gray, 1821 – Small-sized bats (FA – 38.7-42.1 mm) with a moderately long tail (40.0-47.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears broad and not greatly elongated, forward facing and joined over the forehead. Tragus triangular and antitragus undefined. Nostrils open upwards and outwards. Two pairs of upper and three pairs of lower incisors are present. Two pairs each of upper and lower premolars are present. One species – *B. leucomelas* (Cretzschmar, 1826) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.4.

Otonycteris Peters, 1859 – Large-sized bats (FA – 64.7-65.6 mm) with a moderately long tail (40.0-58.0 mm). The interfemoral membrane encloses the entire

but the extreme tip of the tail. Ears elongated. Tragus large and antitragus small with a shallow notch. Nostrils are crescent shaped. One pair of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. One species – *O. hemprichii* Peters, 1859 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.4.

Scotoecus Thomas, 1901 – Small-sized bats (FA – 34.1-37.3 mm) with a moderately long tail (34.0-41.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears moderate with long and narrow tragus. Muzzle broadened and flattened. One pair of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. One species – *S. pallidus* Dobson, 1876 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.5.

Scotomanes Dobson, 1875 – Medium to large-sized bats (FA – 56.1-61.2 mm) with a moderately long tail (52.0-66.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears elongated with broad tragus. Antitragus undefined. Nostrils simple and face slightly outward. One pair of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. One species – *S. ornatus* (Blyth, 1851) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.5.

Scotophilus Leach, 1821 – Medium to large-sized bats (FA – 44.0-65.8 mm) with a moderately long tail (40.0-71.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears relatively small with crescent shaped tragus. Antitragus well-developed. Nostrils simple and face slightly outward. One pair of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. Two species – *S. kuhlii* Leach, 1821 and *S. heathi* Horsfield, 1831 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.5.

Arielulus Hill & Harrison, 1987 – Large-sized bats (FA – 41.8-43.6 mm) with a moderately long tail (~40.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears short and broad, tragus well-developed and broadest in the middle. Usually two pairs of upper and three pairs of lower incisors; and two pairs each of upper and lower premolars are present. One species – *A. circumdatus* (Temminck, 1840) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species

belonging to it in Table 7.6.

Hesperoptenus Peters, 1869 – Medium to large-sized bats (FA – 50.0-60.4 mm) with a moderately long tail (44.0-63.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears large, thick and fleshy with large crescent shaped tragus. Antitragus well-developed. Muzzle broadened. Two pairs of upper incisors and three pairs of lower incisors are present. One pair of upper premolars and two pairs of lower premolars are present. One species – *H. tickelli* (Blyth, 1851) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.6.

Eptesicus Rafinesque, 1820 – Small to medium-sized bats (FA – 35.4-55.1 mm) with a moderately long tail (38.0-58.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears large with short and blunt tragus. Antitragus not well defined. Two pairs of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. Seven species – *E. serotinus* (Schreber, 1774), *E. bottae* (Peters, 1869), *E. pachyotis* (Dobson, 1871), *E. nasutus* (Dobson, 1877), *E. dimissus* Thomas, 1916, *E. gobiensis* Bobrinskii, 1926 and *E. tatei* Ellerman & Morrison-Scott, 1951 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.6.

Nyctalus Bowdich, 1825 – Medium-sized bats (FA – 42.1-57.8 mm) with a moderately long tail (31.0-55.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears relatively short with a club shaped tragus. Two pairs of upper and three pairs of lower incisors are present. Two pairs each of upper and lower premolars are present. Three species – *N. noctula* (Schreber, 1774); *N. leisleri* (Kuhl, 1817); and *N. montanus* (Barrett-Hamilton, 1906) – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.7.

Pipistrellus Kaup, 1829 – Small to medium-sized bats (FA – 25.0-42.6 mm) with a moderately long tail (20.0-49.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears short and broad, tragus well-developed (Fig. 7G). Antitragus is not well defined. On the muzzle, pararhinal glands and internarial groove are distinct. Usually two pairs of upper and three pairs of lower incisors; and two pairs each of upper and lower premolars are present. Some species have reduction in dentition. Eight species – *P. pipistrellus* (Schreber, 1774), *P. kuhlii* (Kuhl, 1817), *P. coromandra* (Gray, 1838), *P. javanicus* (Gray, 1838), *P. abramus* (Temminck, 1840), *P. tenuis* (Temminck, 1840), *P. ceylonicus* (Kelaart, 1852) and *P. paterculus* Thomas, 1915 – occur in South Asia. Diagnostic morphological

characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.7.

Scotozous Dobson, 1875 – Medium-sized bats (FA – 32.7-36.0 mm) with a moderately long tail (27.0-41.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears short and broad, tragus well-developed with small triangular lobe near the base of the outer margin. Penis enlarged. One pair of upper (second upper usually absent, when present very minute) and three pairs of lower incisors; and two pairs each of upper and lower premolars are present. One species – *S. dormeri* Dobson, 1875 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.7.

Tylonycteris Peters, 1872 – Small-sized bats (FA – 26.1-29.0 mm) with a long tail (26.0-33.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears triangular, with short and broad tragus. Head characteristically broadened and flattened. Fleshy pads on the ball of thumb and sole of the foot characteristic. Two pairs of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. Two species – *T. pachypus* (Temminck, 1840) and *T. robustula* Thomas, 1915 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.8.

Ia Thomas, 1902 – Large-sized bats (FA – 70.9-77.3 mm) with a moderately long tail (~ 65.0mm). The interfemoral membrane encloses the entire but the extreme tip of the tail (~ 6.0mm). Ears broad with moderately long tragus. Nostrils simple and face slightly outward. Two pairs of upper and three pairs of lower incisors are present. Two pairs each of upper and lower premolars are present. One species – *I. io* Thomas, 1902 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.8.

Falsistrellus Throughton, 1943 – Medium-sized bats (FA – 38.4-41.4 mm) with a moderately long tail (30.0-41.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears squarish with broadly rounded tips, tragus well-developed with a small triangular lobe near the base of the outer margin. On the muzzle, pararhinal glands and internarial groove are distinct. Two pairs of upper and three pairs of lower incisors; and two pairs each of upper and lower premolars are present. One species – *F. affinis* (Dobson, 1871) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.8.

Philetor Thomas, 1902 – Small-sized bats (FA – 31.7-35.7 mm) with a moderately long tail (27.1-32.2 mm). The interfemoral membrane encloses the entire

but the extreme tip of the tail. Ears short with broad, short fleshy and thick tragus. Two pairs of upper incisors and three pairs of lower incisors are present. One pair of upper premolars and two pairs of lower premolars are present. One species – *P. brachypterus* (Temminck, 1840) – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.8.

Vespertilio Linnaeus, 1758 – Medium-sized bats (FA – 42.0-45.5 mm) with a moderately long tail (40.0-48.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears relatively small and broad, with small, short, blunt and rounded tragus. Antitragus well defined. Nostrils simple and face slightly outward. Two pairs of upper and three pairs of lower incisors are present. One pair of upper and two pairs of lower premolars are present. One species – *V. murinus* Linnaeus, 1758 – occurs in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.8.

Hypsugo Kolenati, 1856 – Medium-sized bats (FA – 32.1-38.0 mm) with a moderately long tail (30.0-49.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears short and broad, tragus well-developed. Antitragus is not well defined. Usually two pairs of upper and three pairs of lower incisors; and one pair of upper and two pairs lower premolars are present. Two species – *H. savii* (Bonaparte, 1873) and *H. cadornae* (Thomas, 1916) – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 7 and those of the species belonging to it in Table 7.8.

Family Miniopteridae

Includes about 19 species of insectivorous bats with wide distribution in Americas, Europe, Africa, Asia, Australia and Oceania (Mickleburgh et al. 2002; Simmons 2005). Muzzle simple and lacks the noseleaf. Ears separate from each other. Tragus is well-developed. The long tail is completely enclosed in the interfemoral membrane or the extreme tip protrudes out. Characteristically posses greatly elongated third digit whose second phalanx is markedly longer than that of the first, owing to which the distal end of the wing bends over the body while at rest. Three species belonging to one genus are reported from South Asia.

Miniopterus Bonaparte, 1837 – Medium-sized bats (FA – 39.6-49.6 mm) with a long tail (39.6-61.0 mm). The interfemoral membrane encloses the entire but the extreme tip of the tail. Ears small with tall, slender and slightly forwardly curved tragus (Fig. 7H). Muzzle short. Two pairs of upper incisors and three pairs of lower incisors are present. Two pairs of upper premolars and three pairs of lower premolars are present. Three

species – *M. fuliginosus* (Hodgson, 1835), *M. pusillus* Dobson, 1876 and *M. magnater* Sanborn, 1931 – occur in South Asia. Diagnostic morphological characters of the genus are provided in Table 8 and those of the species belonging to it in Table 8.1.

REFERENCES

- Bates, P.J.J. & D.L. Harrison (1997).** *Bats of the Indian Subcontinent*. Harrison Zoological Museum Publications, Sevenoaks, UK, 258 pp.
- Blanford, W.T. (1888-91).** *The Fauna of British India, Mammalia*. Taylor & Francis, London, 617pp. (in two parts).
- Corbet, G.B. & J.E. Hill (1992).** *The Mammals of the Indomalayan Region*. British Museum (Natural History)/ Oxford University Press, London, 488pp.
- Eick, G.N., D.S. Jacobs & C.A. Matthee (2005).** A nuclear DNA phylogenetic perspective on the evolution of echolocation and historical biogeography of extant bats (Chiroptera). *Molecular Biology and Evolution* 22(9): 1869-1886
- Gunnel, G.F. & N.B. Simmons (2005).** Fossil evidence and the origin of bats. *Journal of Mammalian Evolution* 12: 209-246.
- Hutson, A.M., S.P. Mickleburgh & P.A. Racey (compilers) (2001).** *Microchiropteran Bats: Global Status, Survey and Conservation Action Plan*. International Union for Conservation of Nature and Natural Resources, Chiroptera Specialist Group (IUCN/SSC). Gland, Switzerland, 258pp.
- Jones, G. & E.C. Teeling (2006).** The evolution of echolocation in bats. *Trends in Ecology and Evolution* 21(3): 149-156.
- Koopman, K.F. (1993).** Order Chiroptera, pp.137-241. In: Wilson, D.E. & D.M. Reeder (eds.). *Mammal Species of the World, 2nd Edition*. Smithsonian Institution Press, Washington, 1206pp.
- Mickleburgh, S.P., A.M. Hutson & P.A. Racey (compilers) (1992).** *Old World Fruit Bats. An Action Plan for their Conservation*. International Union for Conservation of Nature and Natural Resources, Chiroptera Specialist Group (IUCN/SSC), Gland, Switzerland, 252pp.
- Mickleburgh, S.P., A.M. Hutson & P.A. Racey (2002).** A review of the global conservation status of bats. *Oryx* 36(1): 18-34.
- Molur, S., G. Marimuthu, C. Srinivasulu, S. Mistry, A.M. Hutson, P.J.J. Bates, S. Walker, K.P. Priya & A.R.B. Priya (eds.) (2002).** *Status of South Asian Chiroptera: Conservation Assessment and Management Plan (C.A.M.P.) Workshop Report*. Zoo Outreach Organisation, Conservation Breeding Specialist Group South Asia, and Wildlife Information & Liaison Development Society, Coimbatore, India, viii+154pp.+CD.
- Simmons, N.B. (2005).** Order Chiroptera, pp.312-529. In: D.E. Wilson and D.M. Reeder (eds.), *Mammal Species of the World, 3rd Edition, Volume 1*. The John Hopkins University Press, Baltimore, MD, USA. 2142 pp.
- Simmons, N.B. & J.H. Geisler (1998).** Phylogenetic relationships of *Icaronycteris*, *Archaeonycteris*, *Hassianycteris* and *Palaeochiroptryx* to extant bat lineages with comments on the evolution of echolocation and foraging strategies in Microchiroptera. *Bulletin of the American Museum of Natural History* 235: 1-182.
- Springer, M.S., E.C. Teeling, O. Madsen, M.J. Stanhope & W.W. De Jong (2001).** Integrated fossil and molecular data reconstruct bat echolocation. *Proceedings of the National Academy of Sciences (USA)* 98: 6241-6246.
- Srinivasulu, C. & B. Srinivasulu (2001).** Bats of the Indian subcontinent. *Current Science* 80(11): 1378-1380.
- Srinivasulu, C., B. Srinivasulu & Y.P. Sinha (in review).** Bats (Mammalia: Chiroptera) of South Asia: Biogeography, Diversity, Taxonomy and Distribution. *Journal of Threatened Taxa*.
- Teeling, E.C., M.S. Springer, O. Madsen, P. Bates, S.J. O'Brien & W.J. Murphy (2005).** A molecular phylogeny for bats illuminates biogeography and the fossil record. *Science* 307: 580-584.

Appendix I. Some important literature consulted to prepare the key

- Advani, R. (1982).** Distribution and status of chiroptera species in Rajasthan, India. *Saugertierkundliche Mitteilungen* 30(1): 49-52.
- Aellen, V. (1959).** Contribution à l'étude de la faune d'Afghanistan. Chiroptères. *Revue suisse Zoologie* 66: 353-386.
- Agrawal, V.C. (1973).** On a collection of bats from Goa. *Records of the Zoological Survey of India* 67: 261-280.
- Agrawal, V.C. & T.P. Bhattacharyya (1976).** Report on a collection of mammals from Nagarjuna Sagar, Andhra Pradesh. *Newsletter of the Zoological Survey of India* 2(5): 212-216.
- Agrawal, V.C. & T.P. Bhattacharyya (1977).** Report on a collection of mammals from Tripura. *Records of the Zoological Survey of India* 73(1-4): 135-157.
- Agrawal, V.C. & S. Chakraborty (1971).** Notes on a collection of small mammals from Nepal, with the description of a new mouse-hare (Lagomorpha: Ochotonidae). *Proceedings of the Zoological Society, Calcutta* 24(1): 41-46.
- Agrawal, V.C., P.K. Das, S. Chakraborty, R.K. Ghose, A.K. Mandal, T.K. Chakraborty, A.K. Poddar, J.P. Lal, T.P. Bhattacharyya & M.K. Ghosh (1992).** Mammalia. pp.27-169. In: Director, ZSI (ed.). *State Fauna Series 3: Fauna of West Bengal, Part 1*. Zoological Survey of India, Calcutta.
- Agrawal, V.C. & Y.P. Sinha (1973).** Studies on the bacula of some oriental bats. *Anatomischer Anzeiger* 133: 180-192.
- Ahmed, S.K. & K.Z. Husain (1982).** Bats of Bangladesh. *Journal Asiatic Society of Bangladesh (Science)* 8: 89-99.
- Allen, G.M. (1936).** Two new races of Indian bats. *Records Indian Museum* 38: 343-346.
- Allen, G.M. (1938).** *The Mammals of China and Mongolia*. American Museum of Natural History, New York, 620pp.
- Andersen, K. (1908).** Twenty new forms of *Pteropus*. *Annals and Magazine of Natural History* 2: 361-370.
- Andersen, K. (1910).** Ten new fruit-bats of the genera *Nyctimene*, *Cynopterus* and *Eonycteris*. *Annals and Magazine of Natural History* 7: 641-643.
- Andersen, K. (1911).** Six new fruit-bats of the genera *Macroglossus* and *Syconycteris*. *Annals and Magazine of Natural History* 7: 641-643.

Table 1. Classical and modern classification of bats

| Classical | Modern | | |
|--------------------------|--|---|--|
| Koopman 1993 | Simmons & Geisler 1998 Gunnels & Simmons 2005 | Springer et al. 2001 Teeling et al. 2005 | Eick et al. 2005 Jones & Teeling 2006 |
| Order Chiroptera | Order Chiroptera | Order Chiroptera | Order Chiroptera |
| Suborder Megachiroptera | Suborder Megachiroptera | Suborder Yinpterochiroptera | Suborder Yinpterochiroptera |
| Family Pteropodidae | Family Pteropodidae | Superfamily Pteropodoidea | Superfamily Pteropodoidea |
| Suborder Microchiroptera | Suborder Microchiroptera | Family Pteropodidae | Family Pteropodidae |
| Family Rhinopomatidae | Superfamily Emballonuroidea | Superfamily Rhinolophoidea | Superfamily Rhinolophoidea |
| Family Craseonycteridae | Family Emballonuridae | Family Rhinopomatidae | Family Rhinolophidae |
| Family Nycteridae | Infraorder Yinochiroptera | Family Megadermatidae | Family Hipposideridae |
| Family Megadermatidae | Superfamily Rhinopomatoidea | Family Rhinolophidae | Family Megadermatidae |
| Family Rhinolophidae | Family Rhinopomatidae | Subfamily Rhinolophinae | Family Craseonycteridae |
| Family Hipposideridae | Superfamily Rhinolophoidea | Subfamily Hipposiderinae | Family Rhinopomatidae |
| Family Mormoopidae | Family Nycteridae | Suborder Yangochiroptera | Suborder Yangochiroptera |
| Family Noctilionidae | Family Megadermatidae | Family Nycteridae <i>incertae sedis</i> | Family Nycteridae |
| Family Phyllostomidae | Family Rhinolophidae | Superfamily Emballonuroidea | Superfamily Emballonuroidea |
| Family Mormoopidae | Subfamily Hipposiderinae | Family Emballonuridae | Family Emballonuridae |
| Family Noctilionidae | Subfamily Rhinolophinae | Superfamily Noctilionoidea | Superfamily Noctilionoidea |
| Family Furipteridae | Infraorder Yangochiroptera | Family Noctilionidae | Family Phyllostomidae |
| Family Thyropteridae | Superfamily Noctilionoidea | Family Phyllostomidae | Family Mormoopidae |
| Family Natalidae | Family Noctilionidae | Superfamily Vespertilioidea | Family Noctilionidae |
| Family Mystacinidae | Family Phyllostomidae | Family Natalidae | Family Furipteridae |
| Family Vespertilionidae | Superfamily Nataloidea | Family Vespertilionidae | Family Thyropteridae |
| Family Molossidae | Family Natalidae | Family Molossidae | Family Mystacinidae |
| | Superfamily Molossoidea | | Superfamily Vespertilioidea |
| | Family Antrozoidae | | Family Vespertilionidae |
| | Family Molossidae | | Family Molossidae |
| | Superfamily Vespertilioidea | | Family Miniopteridae |
| | Family Vespertilionidae | | Family Natalidae |

Table 2. Checklist of bats of South Asia

| | |
|---|--|
| Family Pteropodidae (Old World fruit bats) [8 Genera, 13 Species] | |
| 1. <i>Rousettus aegyptiacus</i> (E. Geoffroy, 1810) | 37. <i>Rhinolophus blasii</i> Peters, 1867 |
| 2. <i>Rousettus leschenaultii</i> (Desmarest, 1820) | 38. <i>Rhinolophus macrotis</i> Blyth, 1844 |
| 3. <i>Pteropus giganteus</i> Brunnich, 1782 | 39. <i>Rhinolophus luctus</i> Temminck, 1835 |
| 4. <i>Pteropus faunulus</i> Miller, 1902 | 40. <i>Rhinolophus beddomei</i> Andersen, 1905 |
| 5. <i>Pteropus hypomelanus</i> Temminck, 1853 | 41. <i>Rhinolophus trifoliatus</i> Temminck, 1834 |
| 6. <i>Pteropus melanotus</i> Blyth, 1863 | 42. <i>Rhinolophus pearsonii</i> Horsfield, 1851 |
| 7. <i>Cynopterus sphinx</i> (Vahl, 1797) | 43. <i>Rhinolophus yunanensis</i> Dobson, 1872 |
| 8. <i>Cynopterus brachyotis</i> (Muller, 1838) | 44. <i>Rhinolophus mitratus</i> Blyth, 1844 |
| 9. <i>Megaerops niphanae</i> Yenbutra & Felten, 1983 | Family Hipposideridae (Leaf-nosed bats) [4 Genera, 15 Species] |
| 10. <i>Latidens salimalii</i> Thonglongya, 1972 | 45. <i>Hipposideros ater</i> Templeton, 1848 |
| 11. <i>Sphaerias blanfordi</i> (Thomas, 1891) | 46. <i>Hipposideros cineraceus</i> Blyth, 1853 |
| 12. <i>Eonycteris spelaea</i> (Dobson, 1871) | 47. <i>Hipposideros durgadasi</i> Khajuria, 1970 |
| 13. <i>Macroglossus sobrinus</i> K. Andersen, 1911 | 48. <i>Hipposideros fulvus</i> Gray, 1838 |
| Family Rhinopomatidae (Mouse-tailed bats) [1 Genus, 3 Species] | 49. <i>Hipposideros pomona</i> K. Andersen, 1918 |
| 14. <i>Rhinopoma microphyllum</i> (Brunnich, 1782) | 50. <i>Hipposideros hypophyllus</i> Kock & Bhat, 1994 |
| 15. <i>Rhinopoma hardwickii</i> Gray, 1831 | 51. <i>Hipposideros galeritus</i> Cantor, 1846 |
| 16. <i>Rhinopoma muscatellum</i> Thomas, 1903 | 52. <i>Hipposideros speoris</i> (Schneider, 1800) |
| Family Emballonuridae (Sheath-tailed bats) [2 Genera, 6 Species] | 53. <i>Hipposideros larvatus</i> (Horsfield, 1823) |
| 17. <i>Taphozous longimanus</i> Hardwicke, 1825 | 54. <i>Hipposideros armiger</i> (Hodgson, 1835) |
| 18. <i>Taphozous melanopogon</i> Temminck 1841 | 55. <i>Hipposideros lankadiva</i> Kelaart, 1850 |
| 19. <i>Taphozous perforatus</i> E. Geoffroy, 1818 | 56. <i>Hipposideros diadema</i> (E. Geoffroy, 1813) |
| 20. <i>Taphozous theobaldi</i> Dobson, 1872 | 57. <i>Triaenops persicus</i> Dobson, 1871 |
| 21. <i>Taphozous nudiventris</i> Cretzschmar, 1830 | 58. <i>Asellia tridens</i> E. Geoffroy, 1813 |
| 22. <i>Saccoaimus saccoaimus</i> (Temminck, 1838) | 59. <i>Coelops frithi</i> Blyth, 1848 |
| Family Megadermatidae (False Vampire bats) [1 Genus, 2 Species] | Family Molossidae (Free-tailed bats) [3 Genera, 4 Species] |
| 23. <i>Megaderma lyra</i> E. Geoffroy, 1810 | 60. <i>Tadarida teniotis</i> (Rafinesque, 1814) |
| 24. <i>Megaderma spasma</i> (Linnaeus, 1758) | 61. <i>Tadarida aegyptiaca</i> (E. Geoffroy, 1818) |
| Family Rhinolophidae (Horseshoe bats) [1 Genus, 20 Species] | 62. <i>Chaerephon plicatus</i> (Buchanan, 1800) |
| 25. <i>Rhinolophus mehelyi</i> Matschie, 1901 | 63. <i>Otomops wroughtoni</i> (Thomas, 1913) |
| 26. <i>Rhinolophus ferrumequinum</i> (Schreber, 1774) | Family Vespertilionidae (Evening bats) [23 Genera, 63 Species] |
| 27. <i>Rhinolophus bocharicus</i> Kastschenko and Akimov, 1917 | 64. <i>Myotis blythii</i> (Tomes, 1857) |
| 28. <i>Rhinolophus affinis</i> Horsfield, 1823 | 65. <i>Myotis sicarius</i> Thomas, 1915 |
| 29. <i>Rhinolophus rouxii</i> Temminck, 1835 | 66. <i>Myotis formosus</i> (Hodgson, 1835) |
| 30. <i>Rhinolophus sinicus</i> Andersen, 1905 | 67. <i>Myotis nipalensis</i> (Dobson, 1871) |
| 31. <i>Rhinolophus hipposideros</i> (Bechstein, 1800) | 68. <i>Myotis muricola</i> (Gray, 1846) |
| 32. <i>Rhinolophus pusillus</i> Temminck, 1834 | 69. <i>Myotis siligorensis</i> (Horsfield, 1855) |
| 33. <i>Rhinolophus subbadius</i> Blyth, 1844 | 70. <i>Myotis montivagus</i> (Dobson, 1874) |
| 34. <i>Rhinolophus lepidus</i> Blyth, 1844 | 71. <i>Myotis annectans</i> (Dobson, 1871) |
| 35. <i>Rhinolophus shortridgei</i> Andersen, 1918 | 72. <i>Myotis longipes</i> (Dobson, 1873) |
| 36. <i>Rhinolophus cognatus</i> K. Andersen, 1906 | 73. <i>Myotis csorbai</i> Topal, 1997 |
| | 74. <i>Myotis laniger</i> (Peters, 1871) |
| | 75. <i>Myotis horsfieldii</i> (Temminck, 1840) |

| | |
|--|---|
| 76. <i>Myotis hasseltii</i> (Temminck, 1840) | 121. <i>Harpiola grisea</i> Peters, 1872 |
| 77. <i>Myotis emarginatus</i> (E. Geoffroy, 1806) | 122. <i>Harpiocephalus harpia</i> (Temminck, 1840) |
| 78. <i>Plecotus homochrous</i> Hodgson, 1847 | 123. <i>Kerivoula picta</i> (Pallas, 1767) |
| 79. <i>Plecotus wardi</i> Thomas, 1911 | 124. <i>Kerivoula hardwickii</i> (Horsfield, 1824) |
| 80. <i>Plecotus strelkovi</i> Spitzenberger, 2008 | 125. <i>Kerivoula lenis</i> Thomas, 1916 |
| 81. <i>Barbastella leucomelas</i> (Cretzschmar, 1826) | Family Miniopteridae (Long-fingered Bats) [1 Genus, 3 Species] |
| 82. <i>Otonycteris hemprichii</i> Peters, 1859 | 126. <i>Miniopterus schreibersii</i> (Kuhl, 1817) |
| 83. <i>Scotomanes ornatus</i> (Blyth, 1851) | 127. <i>Miniopterus pusillus</i> Dobson, 1876 |
| 84. <i>Scotophilus heathi</i> Horsfield, 1831 | 128. <i>Miniopterus magnater</i> Sanborn, 1931 |
| 85. <i>Scotophilus kuhlii</i> Leach, 1821 | |
| 86. <i>Eptesicus serotinus</i> (Schreber, 1774) | |
| 87. <i>Eptesicus bottae</i> (Peters, 1869) | |
| 88. <i>Eptesicus pachyotis</i> (Dobson, 1871) | |
| 89. <i>Eptesicus dimissus</i> Thomas, 1916 | |
| 90. <i>Eptesicus gobiensis</i> Bobrinskii, 1926 | |
| 91. <i>Eptesicus nasutus</i> (Dobson, 1877) | |
| 92. <i>Eptesicus tatei</i> Ellerman & Morrison-Scott, 1951 | |
| 93. <i>Vespertilio murinus</i> Linnaeus, 1758 | |
| 94. <i>Ia io</i> Thomas, 1902 | |
| 95. <i>Tylonycteris pachypus</i> (Temminck, 1840) | |
| 96. <i>Tylonycteris robustula</i> Thomas, 1915 | |
| 97. <i>Pipistrellus pipistrellus</i> (Schreber, 1774) | |
| 98. <i>Pipistrellus paterculus</i> Thomas, 1915 | |
| 99. <i>Pipistrellus javanicus</i> (Gray, 1838) | |
| 100. <i>Pipistrellus coromandra</i> (Gray, 1838) | |
| 101. <i>Pipistrellus tenuis</i> (Temminck, 1840) | |
| 102. <i>Pipistrellus ceylonicus</i> (Kelaart, 1852) | |
| 103. <i>Pipistrellus kuhlii</i> (Kuhl, 1817) | |
| 104. <i>Pipistrellus abramus</i> (Temminck, 1840) | |
| 105. <i>Hypsugo savii</i> (Bonaparte, 1837) | |
| 106. <i>Hypsugo cadornae</i> (Thomas, 1916) | |
| 107. <i>Falsistrellus affinis</i> (Dobson, 1871) | |
| 108. <i>Arielulus circumdatus</i> (Temminck, 1840) | |
| 109. <i>Scotozous dormeri</i> (Dobson, 1875) | |
| 110. <i>Scotoecus pallidus</i> (Dobson, 1876) | |
| 111. <i>Nyctalus noctula</i> (Schreber, 1774) | |
| 112. <i>Nyctalus leisleri</i> (Kuhl, 1817) | |
| 113. <i>Nyctalus montanus</i> (Barrett-Hamilton, 1906) | |
| 114. <i>Philetor brachypterus</i> (Temminck, 1840) | |
| 115. <i>Hesperoptenus tickelli</i> (Blyth, 1851) | |
| 116. <i>Murina leucogaster</i> Milne-Edwards, 1872 | |
| 117. <i>Murina aurata</i> Milne-Edwards, 1872 | |
| 118. <i>Murina cyclotis</i> Dobson, 1872 | |
| 119. <i>Murina tubinaria</i> (Scully, 1881) | |
| 120. <i>Murina huttoni</i> (Peters, 1872) | |

Key to the Suborders and the Families

- 1A. Eyes conspicuous (>4mm in diameter); ear pinna simple, tragus and anti tragus absent; noseleaf absent; second digit of the wing with well-developed phalanges, with claw (with one exception); tail absent, if present small (shorter than ¼ of tibia); interfemoral membrane reduced or virtually absent; head dog-like in profile

Family Pteropodidae Key 1

- 1B. Eyes medium to small (<3mm in diameter); ear pinna with distinctive tragus or/and antitragus; second digit of the wing with reduced phalanges and no claw; tail present (lacking in one family), long (longer than ½ of tibia); interfemoral membrane moderate to wide; head not dog-like in profile

Go to 2

- 2A. Muzzle with a simple or complex noseleaf, with or without supplementary leaflets and other processes; tragus present or absent

3

- 2B. Noseleaf usually absent; may or may not have dermal ridge between the nostrils; tragus present

4

- 3A. Noseleaf simple, with an erect leaf behind nostrils and a heart-shaped leaf surrounding nostrils; tragus bifid; antitragal lobe not developed; ear pinnae large (nearly 2/3rd of the forearm length), fused at bases; tail absent

Family Megadermatidae Key 2

- 3B. Noseleaf complex, consisting of a horseshoe; tragus absent; antitragal lobe conspicuous; tail enclosed within the interfemoral membrane

5

- 4A. Muzzle with small dermal ridge; second and third digits of each wing with two distinct bony phalanges; tail thin with greatest part projecting free from the narrow interfemoral membrane

Family Rhinopomatidae Key 3

- 4B. Muzzle without a dermal ridge; ears may or may not be joined over the forehead; interfemoral membrane well-developed; tail may or may not be free

6

- 5A. Noseleaf triangular with sella between the nostrils; lancet pointed and raising to near or above the forehead; anterior noseleaf horseshoe shaped; antitragus distinct; toes with three joints each

Family Rhinolophidae Key 4

- 5B. Noseleaf squarish without sella or lancet; anterior noseleaf lobate or horseshoe shaped; posterior leaf simple or complex; supplementary leaflets may or may not be present; antitragus notch-like; toes with two joints each

Family Hipposideridae Key 5

- 6A. Tail completely enclosed within the wide interfemoral membrane or the last one or two vertebrae are free

7

- 6B. At least a part of the tail free from the interfemoral membrane **8**
- 7A. Tail completely enclosed within the wide interfemoral membrane or the last one or two vertebrae are free; second phalanx of third finger only slightly longer than first
Family Vespertilionidae Key 6
- 7B. Tail completely enclosed within the wide interfemoral membrane; second phalanx of third finger much longer than first, the lengthened distal part folds upon the wing at rest
Family Miniopteridae Key 7
- 8A. Tail emerging on the upper surface from the mid-point of the interfemoral membrane
Family Emballonuridae Key 8
- 8B. Tail emerging out from the outer edge of the interfemoral membrane, wrinkle-lipped
Family Molossidae Key 9

Keys to the Genera and Species of Bats of South Asia

Key 1, Family Pteropodidae (13 species)

- 1A. Second digit of the wing lacks a claw (claw present only on thumb); tongue sharply pointed and protrusible; tail relatively long, about equal to hind foot length; a pair of large anal glands distinct; Four upper and four lower incisors; FA: 66.0-78.0 mm; CBL: 31.7-36.3 mm; CM²: 11.9-13.4 mm
Eonycteris spelaea
- 1B. Claws present on both thumb and second digit; tail usually shorter than hind foot or obscure **2**
- 2A. Large body size; in adults, forearm length more than 110mm **3**
- 2B. Small to medium body size; in adults, forearm length less than 100mm **4**
- 3A. Forearm less than 150mm; ears moderate in length, bluntly rounded **5**
- 3B. Forearm usually longer than 150mm; ears long (E: 28.0-57.0 mm), more or less pointed **6**
- 4A. Tail extremely short (5mm or less) or absent **7**
- 4B. Tail longer than 10mm **8**

5A. Pelage on the back pale russet brown with gray and black hairs; ear length 25.0-28.0 mm, short and broad, tips rounded; FA: 135.0-145.0 mm; CBL: 59.0-64.1 mm; CM²: 22.8-25.7 mm

Pteropus hypomelanus

5B. Pelage on the back blackish-brown with white hairs; ear length about 22.0mm, moderate and broad, tips rounded; FA: 110.0-116.0 mm; CBL: ~ 48.6mm; CM²: 19.0-19.2 mm

Pteropus faunulus

6A. Pelage on the back blackish-brown with a few paler hairs and on ventral side pale tan to deep orange red or chestnut brown; FA: 145-183 mm; CBL: 61.5-74.9 mm; CM²: 24.0-29.0 mm

Pteropus giganteus

6B. Pelage on the back blackish brown with a few paler hairs and on ventral side dark brown or blackish-brown; FA: 148.0-163.0 mm; CBL: 63.3-70.7 mm; CM²: 24.6-27.9 mm

Pteropus melanotos

7A. Muzzle narrow and elongate, slightly curved downwards; tongue very long and slender with papillae on distal part; tail absent or rudimentary (3.5-5.5 mm); wing membrane attached to dorsal side of foot; FA: 44.0-52.0 mm; CBL: 26.6-27.1 mm; CM²: 8.6-8.8 mm

Macroglossus sobrinus

7B. Muzzle not narrow and elongate, not curved downwards; tongue moderately long, lacks papillae on distal part; tail totally absent; wing membrane attached to the outer side of the foot or to outmost digit

9

8A. Forearm less than 75mm; muzzle short and relatively broad; ears with conspicuous white margins

10

8B. Forearm longer than 75mm; muzzle moderate in length and breadth; ears without any white margins

11

9A. Forearm greater than 65mm; one pair each of upper and lower incisors present; FA: 66.0-69.0 mm; CBL: 31.4-33.0 mm; CM¹: 11.3-11.8 mm

Latidens salimalii

9B. Forearm lesser than 65mm; two pairs of upper incisors, one or two pairs of lower incisors present

12

10A. Ears large (E: 17.5-24.0 mm) with well-developed pale anterior and posterior border; metacarpals and phalanges pale; FA: 64.0-79.0 mm; CBL: 28.4-33.3 mm; CM¹: 10.2-12.2 mm

Cynopterus sphinx

10B. Ears small (E: 14.5-18.0 mm) with poorly developed or no pale anterior and posterior border; metacarpals and phalanges dark; FA: 57.3-63.3 mm; CBL: 26.0-28.8 mm; CM¹: 8.9-10.7 mm

Cynopterus brachyotis

- 11A. FA: 83.0-90.0 mm; T: 14.0-20.0 mm; CBL: 36.9 -40.4 mm; CM²: 14.9-16.6 mm
Rousettus aegyptiacus
- 11B. FA: 75.0-86.0 mm; T: 8.0-21.0 mm; CBL: 33.5-37.7 mm; CM²: 13.5-15.2 mm
Rousettus leschenaultii
- 12A. Muzzle moderate; distance between eye and nostril equal to that between eye and ear; ears with thin white margins; rudimentary interfemoral membrane entirely concealed with dense and long fur on hind limbs; two pairs of lower incisors; FA: 52.0-61.0 mm; CBL averages 26.2mm; CM¹: 8.0-8.1 mm
Sphaerias blanfordi
- 12B. Muzzle very short and broad; distance between eye and nostril less than that between eye and ear (auditory meatus); ears without any white margins; interfemoral membrane visible, not entirely concealed by fur; one pair of lower incisors; FA: 52.0-63.0 mm; CBL: 25.6-26.7 mm; CM¹: 8.4-8.6 mm
Megaerops niphanae

Key 2 Family Megadermatidae (2 species)

- 1A. Vertical noseleaf tall (c'10mm), straight sided with longitudinal ridge and a simple rounded horizontal base; inner margins of the ears fused at about 30% of its height; FA: 56.0-71.5 mm; CCL: 24.5-27.8 mm; CM³: 10.6-12.1 mm
Megaderma lyra
- 1B. Vertical noseleaf short (c' 6.5mm); convex sided with longitudinal ridge and a distinctly heart-shaped base; inner margins of the ears fused at about 15% or less of its height; FA: 54.0-62.0 mm; CCL: 21.9-23.6 mm; CM³: 9.3-10.5 mm
Megaderma spasma

Key 3, Family Rhinopomatidae (3 species)

- 1A. Tail (50.0-77.0 mm) shorter than FA; FA: 59.5-74.6 mm; CCL: 17.2-22.7 mm; CM³: 7.0-8.0 mm
Rhinopoma microphyllum
- 1B. Tail (49.0-78.0 mm) longer than FA; FA: 46.0-64.0 mm
2
- 2A. Dermal ridge on the muzzle well-developed; FA: 52.9-64.0 mm; CCL: 15.5-17.5 mm; CM³: 6.0-6.8 mm
Rhinopoma hardwickii
- 2B. Dermal ridge on the muzzle poorly developed; FA: 46.0-53.2 mm; CCL: 14.7-15.1 mm; CM³: 5.3-5.9 mm
Rhinopoma muscatellum

Key 4, Family Rhinolophidae (20 species)

- 1A. Sella with lateral projecting lappets at its base 2
- 1B. Sella lacking lateral basal lappets 3
- 2A. Noesleaf and ears brownish; inferior surface of the sella broad (c' 4.3mm) 4
- 2B. Noseleaf and ears yellowish; inferior surface of the sella relatively narrow (c' 1.4mm at the apex); FA: 50.0-53.3 mm; CCL: 19.5-21.5 mm; CM³: 8.4-8.9 mm
Rhinolophus trifoliatus
- 3A. Connecting process pointed 5
- 3B. Connecting process rounded, not pointed 6
- 4A. Large in size (FA >70.0mm); inferior surface of the sella parallel-sided and broad (c' 4.3mm); FA: 70.0 -80.5 mm; CCL: 27.5-28.7 mm; CM³: 11.8-12.6 mm
Rhinolophus luctus
- 4B. Small in size (FA <65.0mm); inferior surface of the sella parallel-sided and moderately broad (c' 2.4mm); FA: 54.9-64.3 mm; CCL: 22.2-24.4 mm; CM³: 9.6-10.4 mm
Rhinolophus beddomei
- 5A. Anterior median swellings moderately low, protruding anteriorly 7
- 5B. Anterior median swellings higher, directing upwards 8
- 6A. Connecting process low, rounded, its tip pointed more or less downwards; superior connecting process broadly rounded and low; FA: 35.9-40.0 mm; CCL: 13.0-14.0 mm; CM³: 4.6-5.5 mm
Rhinolophus hipposideros
- 6B. Connecting process well-developed, rounded, its tip pointed more or less forwards 9
- 7A. Sella parallel-sided; dorsal pelage light brown, well marked dark brownish patches under each eye; ventral pelage light grayish-brown or even purely white; FA: 47.0-53.0 mm; GTL: 18.8-20.4 mm; CM³: 6.2-7.2 mm
Rhinolophus mehelyi
- 7B. Sella wedge-shaped; dorsal pelage light brown to yellowish-brown, ventral pelage grayish, white or creamy; FA: 43.0-50.0 mm; CCL: 16.0-17.0 mm; CM³: 6.0-7.6 mm
Rhinolophus blasii
- 8A. Superior process more horn like 10

| | |
|---|--------------------------------|
| 8B. Superior process not horn like | 11 |
| 9A. Connecting process forming a continuous arch or obsolete; lower lip with one mental groove, internarial not expanded | 12 |
| 9B. Connecting process not forming a continuous arch | 13 |
| 10A. Base of the sella broadly rounded; FA: 34.9- 37.8 mm; CCL: 13.0-14.0 mm; CM ³ : 5.4-6.0 mm | <i>Rhinolophus pusillus</i> |
| 10B. Base of the sella not broadly rounded | 14 |
| 11A. Small skull (GTL usually <17.0mm); small upper canine, feeble saggital crest; FA: 37.0-41.8 mm; GTL: 15.7-17.3 mm; CM ³ : 5.6-6.8 mm | <i>Rhinolophus lepidus</i> |
| 11B. Heavy built skull (GTL usually >17.0mm); very long upper canine, strong saggital crest; FA: 39.0-42.0 mm; GTL: 17.2-17.9 mm; CM ³ : 6.5-7.2 mm | <i>Rhinolophus shortridgei</i> |
| 12A. Inferior surface of the sella not rounded and the base expanded | 15 |
| 12B. Inferior surface of the sella rounded and the base narrow; FA: ~ 57.1mm; T: ~ 30.0mm | <i>Rhinolophus mitratus</i> |
| 13A. Sella long and wide; palatal bridge more than 1/3 rd length of maxillary toothrow CM ³ ; superior connecting process broadly rounded; inferior extremity short and emarginated; lancet short with pointed tip; FA: 39.0-46.0 mm; CCL: 15.2-16.6 mm; CM ³ : 6.2-6.7 mm | <i>Rhinolophus macrotis</i> |
| 13B. Sella shorter and narrower; palatal bridge less than 1/3 rd length of maxillary toothrow CM ³ | 16 |
| 14A. Large in size; horn like superior connecting process slightly curved downwards; FA: 37.5-40.7 mm; CCL: 14.2-15.8 mm; CM ³ : 6.0-7.0 mm | <i>Rhinolophus cognatus</i> |
| 14B. Small in size; horn like superior connecting process not curved downwards; FA: 33.4-35.9 mm; CCL: 11.9-12.9 mm; CM ³ : 5.1-5.5 mm | <i>Rhinolophus subbadius</i> |
| 15A. Small; FA: 50.8-54.8 mm; CCL: 20.2-21.6 mm; CM ³ : 9.2-9.9 mm | <i>Rhinolophus pearsonii</i> |
| 15B. Large; FA: 54.2-59.5 mm; CCL: 22.3-22.9 mm; CM ³ : 9.5-10.6 mm | <i>Rhinolophus yunanensis</i> |

- 16A. Connecting process high and bluntly rounded, anterior upper premolar when present always minute and fully extruded from the tooth row **17**
- 16B. Connecting process lower and rounded, anterior premolar larger and usually in row or only slightly extruded, very rarely totally external **18**
- 17A. Large (FA >53.0mm); FA: 54.8-62.1 mm; GTL: 20.8-24.9 mm; CM³: 8.5-9.6 mm
Rhinolophus ferrumequinum
- 17B. Small (FA <53.0mm); FA: 47.5-53.0 mm; GTL: 19.0-22.2 mm; CM³: 6.7-7.2 mm
Rhinolophus bocharicus
- 18A. Lancet triangular, lateral margins more or less straight; FA: 50.0-55.0 mm; CCL: 18.7-20.5 mm; CM³: 8.5-9.5 mm
Rhinolophus affinis
- 18B. Lancet abruptly narrowed at central, its lateral margins strongly concave **19**
- 19A. Second phalanx of third metacarpal 52.8-66.0% of its length; FA: 44.4-52.3 mm; CCL: 17.5-21.0 mm; CM³: 7.9-9.2 mm
Rhinolophus rouxii
- 19B. Second phalanx of third metacarpal 65.0-75.3% of its length; FA: 45.7-49.6 mm; CCL: 17.4-18.4 mm; CM³: 7.6-8.2 mm
Rhinolophus sinicus

Key 5, Family Hipposideridae (15 species)

- 1A. Tail long and enclosed within interfemoral membrane, except for the extreme tip; supplementary lappets absent; noseleaf with or without supplementary leaflets **2**
- 1B. Tail rudimentary (<2.0mm); interfemoral membrane much reduced and narrow; anterior noseleaf, deeply emarginated anteriorly, extends beyond muzzle; with forwardly projecting two supplementary lappets; FA: 37.8-42.0 mm; CCL: 14.3-14.8 mm; CM³: 5.8-6.0 mm
Ceolops frithi
- 2A. Posterior leaf of the noseleaf with three vertical processes **3**
- 2B. Posterior leaf of the noseleaf without three vertical processes **4**
- 3A. Narial lappets well-developed; ears small funnel shaped; noseleaf with a single pair of supplementary leaflets; a fourth vertical process projecting downwards also present; FA: 48.3-54.0 mm; CBL: 16.9-18.9 mm; CM³: 6.1-7.1 mm
Triaenops persicus

3B. Narial lappets less developed; ears broad with pointed tips; noseleaf with two pairs of supplementary leaflets; the central vertical process taller and pointed than others; FA: 50.1-52.3 mm; CCL: 15.7-17.0 mm; CM³: 6.5-7.2 mm

Asellia tridens

- | | |
|--|----|
| 4A. Noseleaf without supplementary leaflets | 5 |
| 4B. Noseleaf with supplementary leaflets | 6 |
| 5A. Ears moderate, averages less than 18.0mm (E: 13.0-20.0 mm) | 7 |
| 5B. Ears large, averages more than 20.0mm (E: 18.5-26.0 mm) | 8 |
| 6A. Two or less than two supplementary leaflets | 9 |
| 6B. Three or four supplementary leaflets | 10 |

7A. Jugal bone of each zygoma with dorsal process; fourth metacarpal exceeds fifth; FA: 34.9-42.3 mm; CCL: 13.2-15.0 mm; CM³: 5.1-5.9 mm

Hipposideros ater

- | | |
|--|----|
| 7B. Jugal bone of each zygoma lacking the dorsal process | 11 |
| 8A. Zygomatica exceeds braincase in width; third metacarpal shorter than fourth; FA: 38.4-44.0 mm; CCL: 15.0-16.4 mm; CM ³ : 6.0-6.9 mm | |

Hipposideros fulvus

8B. Zygomatica equal to braincase in width; third metacarpal shorter than fifth; FA: 38.1-43.2 mm; CCL: 14.6-2-16.0 mm; CM³: 5.5-6.2 mm

Hipposideros pomona

9A. One supplementary leaflet; fifth metacarpal shorter than the third; FA: 37.1-40.0 mm; CCL: 14.2-14.8 mm; CM³: 5.3-5.7 mm

Hipposideros hypophyllus

9B. Two supplementary leaflets; fifth metacarpal significantly shorter than the third; FA: 45.0-51.3 mm; CCL: 14.6-15.8 mm; CM³: 5.3-6.2 mm

Hipposideros galeritus

10A. Three supplementary leaflets, third sometimes reduced or absent

12

10B. Four supplementary leaflets, fourth sometimes reduced or absent

13

11A. Small; tail projects up to 1.0mm beyond the interfemoral membrane; FA: 33.0-36.3 mm; CCL: 12.7 -13.7 mm; CM³: 4.9-5.3 mm

Hipposideros cineraceus

11B. Large; tail projects up to 2.5-4.0 mm beyond the interfemoral membrane; FA: 36.0-37.5 mm; CBL: 13.0-13.9 mm; CM³: 5.0-6.0 mm

Hipposideros durgadasi

12A. Small in size; outer supplementary leaflet distinctly small; ears markedly small (E: 12.5-9.0 mm); FA: 45.6-54.0 mm; CCL: 15.9-17.5 mm; CM³: 6.6-7.5 mm

Hipposideros speoris

12B. Large in size; outer supplementary leaflet normal; ears relatively large (E: 23.0-26.0 mm); FA: 61.2-64.8 mm; CCL: 20.8-21.4 mm; CM³: 8.9-9.4 mm

Hipposideros larvatus

13A. Small in size, forearm less than 65.0; outer leaflet distinctly small or even absent; fleshy outgrowths behind posterior leaf or even protuberances before eyes lacking; FA: 58.8-64.2 mm; CCL: 22.6-22.9 mm; CM³: 9.7-10.0 mm

Hipposideros diadema

13B. Large in size, forearm in the range of 75.0-99.0 mm; fleshy outgrowths present behind posterior leaf, at least in the form of small protuberances before the eyes

14

14A. Second upper incisor greatly enlarged; dorsal pelage fawn, yellow, brown or red; ears short (E: 19.5-30.0 mm); outer leaflet slightly reduced; FA: 75.0-99.0 mm; CCL: 24.8-31.5 mm; CM³: 11.6-14.5 mm

Hipposideros lankadiva

14B. Second upper incisor not greatly enlarged; dorsal pelage dark brown; ears large (E: 26.0-34.0 mm); outer leaflet distinctly small; FA: 85.4-95.0 mm; CCL: 25.6-29.2 mm; CM³: 11.3-12.5 mm

Hipposideros armiger

Key 6, For subfamilies of the family Vespertilionidae (63 species)

1A. Nostrils elongated and tubular

Murininae, Key 6A

1B. Nostrils not elongated

2

2A. Ears funnel-shaped

Kerivoulinae, Key 6B

2B. Ears not funnel-shaped

3

3A. Tragus long, slender, bluntly pointed more or less straight, its length along the anterior margin at least three times its greatest width; usually three pairs of upper and lower premolars

Myotinae, Key 6C

3B. Tragus shorter, wider, generally rounded at tip, usually curved forwards, its length along the anterior margin usually less than three times its greatest width; less than three pairs of upper and lower premolars

Vespertilioninae, Key 6D

Key 6A, Subfamily Murininae (8 species)

- 1A. Forearm less than 44.0mm, last upper molar normal 2
 1B. Forearm more than 44.0mm, last upper molar reduced, often deciduous; FA: 44.1-50.1 mm; CCL: 19.0-19.5 mm; CM³: 6.5-6.9 mm
Harpiocephalus harpia
- 2A. Upper incisors (I² and I³) are approximately two-third that of the C¹ in height; wings attached to the base of the first toe; interfemoral membrane densely covered with rather long hairs above and naked below; FA: ~ 32.8mm; CCL: ~ 12.1mm; CM³: ~ 4.9mm
Harpiola grisea
- 2B. Upper incisors (I² and I³) are almost one-half that of the C¹ in height 3
 3A. Wings attached to the base of the first toe; ears with faint to distinct emargination on the posterior border 4
 3B. Wings attached to the base of the claw of the first toe; ears without emargination on the posterior border 5
 4A. Emargination on the posterior border of the ear indistinct; interfemoral membrane moderately covered with pale gray or ferruginous hairs above and naked below; FA: 30.1-34.1 mm; CCL: 13.4-14.2 mm; CM³: 5.0-5.4 mm
Murina tubinaris
- 4B. Emargination on the posterior border of the ear distinct and conspicuous; Interfemoral membrane moderately covered with long thick reddish hair above and naked below; FA: ~ 40.9mm; CCL: ~ 16.3mm; CM³: ~ 6.1mm
Murina leucogaster
- 5A. Interfemoral membrane naked below 6
 5B. Interfemoral membrane moderately covered with golden hairs above and gray hairs below; FA: 27.7-29.6 mm; CCL: ~ 12.3mm; CM³: ~ 4.5mm
Murina aurata
- 6A. FA: 32.8-35.4 mm; CCL: 14.9-15.5 mm; CM³: 5.8-6.1 mm; Skull relatively long with average CCL being 15.2mm
Murina huttoni
- 6B. FA: 29.7-34.5 mm; CCL: 123.9-15.0 mm; CM³: 5.2-5.7 mm; Skull relatively small with average CCL being 14.4mm
Murina cyclotis

Key 6B, Subfamily Kerivoulinae (3 species)

- 1A. Wing membranes parti-coloured orange and black; interfemoral membrane orange to scarlet, hairy above; FA: 31.5-37.9; CCL: 12.2-13.3 mm; CM³: 5.5-5.8 mm
Kerivoula picta
- 1B. Wing membranes uniformly brown 2
- 2A. Interfemoral membrane sparsely haired above near body parts and posterior border; FA: 31.7-36.0 mm; CCL: 12.4-13.0 mm; CM³: 5.5-5.8 mm
Kerivoula hardwickii
- 2B. Interfemoral membrane lacks hair, sometimes with whitish veins; FA: 37.2-41.0 mm; CCL: 14.5-15.1 mm; CM³: 6.6-6.8 mm
Kerivoula lenis

Key 6C, Subfamily Myotinae (14 species)

- 1A. Wings, forearm and interfemoral membrane conspicuously parti-coloured black, red, orange and yellow; FA: 44.5-49.1 mm; CCL: 16.3-16.6 mm; CM³: 6.8-7.2 mm
Myotis formosus
- 1B. Wings, forearm and interfemoral membrane not parti-coloured 2
- 2A. Outer margin of the ear with conspicuous emarginations; FA: 36.0-41.0 mm
Myotis emarginatus
- 2B. Outer margin of ear lacking any conspicuous emarginations 3
- 3A. Foot including the claws less than half of the tibia 4
- 3B. Foot including the claws half to three fourth of the tibia 5
- 4A. Small (FA <37.0mm); Forearm in the range of 30.0-37.0 mm 6
- 4B. Large (FA >44.0mm); Forearm in the range of 44.7-54.6 mm 7
- 5A. Wings attached to the side of the foot immediately below the ankle 8
- 5B. Wings attached to the foot at the ankle or little above 9

| | |
|--|----------------------------|
| 6A. Canines and molars very weak; FA: 30.0-31.5 mm; CCL: ~ 11.2mm; CM ³ : 4.7-4.8 mm | <i>Myotis siligorensis</i> |
| 6B. Canines and molars well-developed | 10 |
| 7A. Ear relatively large (about 32% of the HB); FA: 48.1-54.6 mm; CCL: 16.9-17.7 mm; CM ³ : 7.1-7.5 mm | <i>Myotis sicarius</i> |
| 7B. Ears not large (<32% of the HB) | 11 |
| 8A. Dorsal pelage sooty gray brown; FA: ~ 34.8mm; CBL: 10.6-11.9 mm; CM ³ : 4.8-5.0 mm | <i>Myotis laniger</i> |
| 8B. Dorsal pelage black; FA: 36.5-41.5 mm; CCL: 13.4-14.3 mm; CM ³ : 5.5-5.9 mm | <i>Myotis horsfieldii</i> |
| 9A. Tragus short and narrow; dorsal pelage dark brown; FA: 37.2-40.3 mm; CCL: 14.0-14.6 mm; CM ³ : 5.8-6.0 mm | <i>Myotis hasseltii</i> |
| 9B. Tragus long and narrow; dorsal pelage brown | 12 |
| 10A. Braincase domed or bulbous; ventral pelage pale gray to cream; FA: 34.5-36.8 mm; CCL: 11.6-12.4 mm; CM ³ : 4.7-5.3 mm | <i>Myotis nipalensis</i> |
| 10B. Braincase flatter; ventral pelage brownish; FA: 31.2-37.0 mm; CCL: 11.5-12.3 mm; CM ³ : 4.8-5.3 mm | <i>Myotis muricola</i> |
| 11A. Ears small, about 25% of HB; FA: 44.7-46.8 mm; CCL: 15.1-15.6 mm; CM ³ : 6.5-6.8 mm | <i>Myotis montivagus</i> |
| 11B. Ears large, about 30% of the HB; FA: 45.3-46.5 mm; CCL: 14.9-15.5 mm; CM ³ : 6.5-6.7 mm | <i>Myotis annectans</i> |
| 12A. Large in size (FA >55.0mm); interfemoral membrane pale brown and hairless throughout; FA: 55.5-58.3 mm; CCL: 18.1-18.9 mm; CM ³ : 8.2-8.5 mm | <i>Myotis blythii</i> |
| 12B. Small in size (FA <41.0mm) | 13 |
| 13A. FA: 36.5-39.0 mm; TIB: 14.4-17.80 mm; CBL: 12.77-13.76 mm; ZB: ~ 8.4 mm; CM ³ : 5.1-5.3 mm | <i>Myotis longipes</i> |
| 13B. FA: 34.80-37.50 mm; TIB: 14.84-15.80 mm; CBL: 12.11-12.71 mm; ZB: 7.59-8.19 mm; CM ³ : 4.72-5.19 mm | <i>Myotis csorbai</i> |

Key 6D, For tribes of the subfamily Vespertilioninae (38 species)

- 1A. Nostrils marginated by grooves on the upper surface of the muzzle; ears long, joined, or may not be joined, across the forehead

Plecotini, Key 6D i

- 1B. Nostrils simple, opening by crescentic or circular apertures at the extremity of the muzzle; ears generally moderate, forehead not grooved

2

- 2A. Total teeth 30; dental formula 1113/3123

Nycticeiini, Key 6D ii

- 2B. Total teeth 32-34; dental formula 211(2)3/3123

3

- 3A. First upper premolar usually absent, except in genus *Arielulus* where it is present; dental formula 2113/3123 in former, 2123/3123 in later

Eptesicini, Key 6D iii

- 3B. First upper premolar usually present; dental formula 2123/3123

4

- 4A. Medium to large bats (FA between 25.0-57.8 mm); pipistrelles, with pararhinal glands (highly variable genus *Pipistrellus*) or pipistrelle-like with relatively less depressed skull and absence or variable presence of I² (genus *Scotozous*) or with narrow wings, high and broad rostra (genus *Nyctalus*)

Pipistrellini, Key 6D iv

- 4B. Medium to very large bats (FA between 26.1- 77.3 mm); pipistrelle-like with variable characters, like enlarged pads on feet and base of thumbs (genus *Tylonycteris*), rounded braincase (genus *Philetor*), large size with forearm usually greater than 70.0mm (genus *Ia*), skull with full rostrum (genus *Vespertilio*), absence of keeled calcar (genus *Hypsugo*) or with pararhinal glands (genus *Falsistrellus*)

Vespertilionini, Key 6D v

Key 6D i, Tribe Plecotini (5 species)

- 1A. Ears long, greater than 30mm in length 2
- 1B. Ears, joined over forehead by a band of integument, are short, up to 18mm in length, barely extending beyond the tip of the muzzle when laid forward, and wide with 5-6 furrows; FA: 38.7-42.1 mm; CCL: 13.4-14.2 mm; CM³: 4.7-5.1 mm
Barbastella leucomelas
- 2A. Small, forearm less than 50mm; ears longer than 30 mm with numerous horizontal furrows, joined by an integument over forehead 3
- 2B. Large, forearm greater than 55mm; ears longer than 35 mm, but not joined by an integument over forehead; tragus half the height of the pinna; antitragus shallow notch-like; dorsal pelage buffy brown, ventral pelage uniformly pale grey; FA: 64.7-65.6 mm; CCL: ~ 21.7mm; CM³: ~ 8.5mm
Otonycteris hemprichii
- 3A. Large, forearm greater than 41mm 4
- 3B. Small, forearm lesser than 41mm; pelage dense and woolly, uniformly dark brown dorsally and ventrally; ears, wing and tail membranes dark brown; FA: 36.5-40.7 mm; CCL: 13.8-14.2 mm; CM³: 5.0-5.4 mm
Plecotus homochrous
- 4A. Pelage dense but not woolly, brownish dorsally and whitish ventrally; ears, wing and tail membranes pale to mid brown; FA: 41.4-44.7 mm; CCL: 14.8-15.5 mm; CM³: 5.4-5.8 mm
Plecotus wardi
- 4B. Pelage not very dense dorsally and shaggy ventrally, cold grayish-drab dorsally and yellowish white ventrally; ears, wing and tail membranes semitransparent cold brown; FA: 39.6-44.8 mm; CCL: 14.1-15.6 mm; CM³: 5.3-6.1 mm
Plecotus strelkovi

Key 6D ii, Tribe Nycticeiini (4 species)

1A. Small size (Forearm less than 38.0mm); head flattened with broad and flat muzzle; tragus long and narrow; FA: 34.1-37.3 mm; CCL: 13.8-14.8 mm; CM³: 5.5-5.9 mm

Scotoecus pallidus

1B. Large size (Forearm more than 44.0mm); head not flattened

2

2A. Dorsal pelage orange brown with white spots on crown, back and shoulders; ears large and lacking any ridges; FA: 56.1-61.2 mm; CCL: 20.1-20.9 mm; CM³: 7.8-8.5 mm

Scotomanes ornatus

2B. Dorsal pelage tawny or chestnut brown, lacks any marking; ears small with numerous transverse ridges

3

3A. Large (Forearm averaging 60.7mm); ventral pelage with distinctly yellow tinge; FA: 55.4-65.8 mm; CCL: 19.0-21.3 mm; CM³: 7.1-8.4 mm

Scotophilus heathi

3B. Smaller (Forearm averaging 49.0mm); ventral pelage buffy brown; FA: 44.0-56.4 mm; CCL: 16.3- 18.0 mm; CM³: 6.1-6.8 mm

Scotophilus kuhlii

Key 6D iii, Tribe Eptesicini (9 species)

- 1A. Ears with well-visible white margins; dorsal pelage is dark brown, the tip of longer guard hairs are bright cuprous-red or golden; interfemoral membrane uniformly dark brown; broad tragus has pale margin; FA: 41.8-43.6 mm; CCL: 14.6-15.6 mm; CM³: 6.0-6.5 mm
Arielulus circumdatus
- 1B. Ears lacking white or pale margins; dorsal pelage lacks cuprous-red or orange hue 2
- 2A. Ears thick and fleshy, yellowish-brown, without ridges; dorsal pelage with distinct yellowish tinge; FA: 50.0-60.4 mm; CCL: 17.2-19.6 mm; CM³: 6.5-6.9 mm
Hesperoptenus tickelli
- 2B. Ears with conspicuous ridges, dark; dorsal pelage without yellowish tinge 3
- 3A. Large in size (FA >50.0mm); ears moderately tall averaging 14.0mm, bears six transverse ridges; FA: 54.2-55.1 mm; CCL: 18.3-19.5 mm; CM³: 7.1-7.8 mm
Eptesicus serotinus
- 3B. Small in size (FA <50.0mm, in the range of 35.4 -45.3 mm) 4
- 4A. Membranes and naked areas pale yellow; FA: 41.0-41.5 mm; CBL: ~ 15.1mm; CM³: ~ 5.9mm
Eptesicus gobiensis
- 4B. Membranes and naked areas shades of brown 5
- 5A. Pelage black throughout; FA: ~ 43.4mm; HB: ~ 48.5mm; T: 45.9mm
Eptesicus tatei
- 5B. Pelage not uniformly black throughout 6
- 6A. Ears triangular in shape with rounded tips; tragus short; proximal part of ear thick and fleshy; FA: 38.0-40.3 mm; HB: 55.0-56.0 mm; T: 40.0-41.0 mm; GTL: ~ 21.2mm
Eptesicus pachyotis
- 6B. Ears not triangular in shape; proximal part of ear normal 7
- 7A. Tail nearly as long as the HB (>90%), rostrum short; FA: 35.4-36.9 mm; CCL: 11.7-12.2 mm; CM³: 4.4-4.8 mm
Eptesicus nasutus
- 7B. Tail not as long as the HB (<85%), rostrum long 8
- 8A. Basioccipital pits prominent; FA: ~ 42mm;
Eptesicus dimissus
- 8B. Basioccipital pits not very prominent; FA: ~ 42.1mm; CCL: ~ 15.0mm; CM³: ~ 5.8mm
Eptesicus bottae

Key 6D iv, Tribe Pipistrellini (12 species)

1A. Ears short (about 28% of the forearm length) and broad; tragus usually half the height of the pinna, with faintly concave anterior border 2

1B. Ears relatively short (about 24% of the forearm length); tragus club-shaped, more than twice wider in distal part than at base 3

2A. Inner upper incisor (i^2) large, unicuspide and in contact with canine or nearly so; outer incisor (i^3) absent, if present, no more than a minute spicule; interfemoral membrae uniformly mid brown, veins sometimes white; FA: 32.7-36.3 mm; CCL: 12.8-13.6 mm; CM³: 5.2-5.6 mm

Scotozous dormeri

2B. Inner upper incisor (i^2) usually bicuspidate; outer incisor large similar in crown area to inner and half or more its height 4

3A. Large (forearm more than 50mm); FA: 50.9-57.8 mm; CCL: 17.1-18.6 mm; CM³: 7.1-7.7 mm

Nyctalus noctula

3B. Small (forearm less than 50mm) 5

4A. Second upper premolar (pm^2) positioned usually in toothrow; interfemoral membrane haired only above but naked below; FA: 30.0-31.6 mm; CCL: 10.4-11.3 mm; CM³: 4.1-4.4 mm

Pipistrellus pipistrellus

4B. Second upper premolar (pm^2) intruded 6

5A. Fur bicoloured; FA: 42.1-45.2 mm; CCL: 14.7-15.2 mm; CM³: 5.5-5.9 mm

Nyctalus leisleri

5B. Fur uniform brown; FA: 42.9-43.0 mm; CCL: 15.4-16.0 mm; CM³: 6.5-6.6 mm

Nyctalus montanus

6A. Large (forearm averaging over 35.0mm); pelage on ventral side conspicuously paler than on dorsal side; interfemoral membrane sparsely haired above and below; FA: 33.0-42.0 mm; CCL: 13.1-14.3 mm; CM³: 5.2-5.9 mm

Pipistrellus ceylonicus

6B. Small (Forearm length averaging less than 35.0mm); pelage on ventral side slightly paler than on dorsal side 7

7A. Interfemoral membrane pallid and translucent; FA: 33.4-36.0 mm; CCL: 12.0-12.9 mm; CM³: 4.6-5.0 mm

Pipistrellus kuhlii

| | |
|--|--------------------------------|
| 7B. Interfemoral membrane not pallid and translucent | 8 |
| 8A. Canine lacks secondary posterior cusps; upper molars relatively massive; pelage dark brown; penis over 10mm in length; FA: 29.2-34.0 mm; CCL: 10.6-11.6 mm; CM ³ : 4.1-4.8 mm | <i>Pipistrellus paterculus</i> |
| 8B. Canine with posterior secondary cusp | 9 |
| 9A. Pelage colouration pale; tragus gradually narrowing along its terminal half, with somewhat pointed apex; penis long, over 8mm in length | 10 |
| 9B. Pelage colouration usually relatively dark; tragus more or less even together; penis short, less than 8mm in length | 11 |
| 10A. Large, forearm between 30.0 to 36 mm; interfemoral membrane not haired; penis short, about 8.0-10.0 mm in length; FA: 30.0-36.0 mm; CCL: 11.9 -13.1 mm; CM ³ : 4.6-5.2 mm | <i>Pipistrellus javanicus</i> |
| 10B. Small, forearm between 31.4 to 34.4 mm; penis long, about 10-13 mm in length; FA: 31.4-34.4 mm; CCL: 11.4-12.8 mm; CM ³ : 4.2-4.9 mm | <i>Pipistrellus abramus</i> |
| 11A. Large, forearm length averaging 30mm; interfemoral membrane haired near the body both above and below; FA: 25.5-34.3 mm; CCL: 10.6-11.9 mm; CM ³ : 3.9-4.6 mm | <i>Pipistrellus coromandra</i> |
| 11B. Small, forearm length averaging 28mm; interfemoral membrane not haired; FA: 25.0-30.2 mm; CCL: 9.3-10.7 mm; CM ³ : 3.5-4.1 mm | <i>Pipistrellus tenuis</i> |

Key 6D v, Tribe Vespertilionini (8 species)

- 1A. Soles of the feet expanded in to fleshy pads; head noticeably flattened **2**
- 1B. Soles of the feet normal; head not flattened **3**
- 2A. Pelage shades of golden brown; FA: 26.1-29.0 mm; CCL: 10.0-11.3 mm; CM³: 3.4-4.2 mm
Tylonycteris pachypus
- 2B. Pelage shades of brown; FA: 26.6-28.1 mm; CCL: 11.1-11.7 mm; CM³: 3.9-4.4 mm
Tylonycteris robustula
- 3A. Large bat, forearm greater than 70.0mm; FA: 70.9-77.3 mm; CCL: 25.2-26.2 mm; CM³: 10.5-11.0 mm
Ia io
- 3B. Small bat, forearm lesser than 70.0mm **4**
- 4A. Two upper premolars and two lower premolars **5**
- 4B. One upper premolar and two lower premolars **6**
- 5A. Large bat, forearm greater than 38mm; braincase small; basisphenoid pits absent; FA: 38.4-41.4 mm; CCL: 13.7-14.5 mm; CM³: 5.5-5.7 mm
Falsistrellus affinis
- 5B. Small bat, forearm lesser than 38mm **7**
- 6A. Tragus fleshy and thickened; FA: 31.7-38.0 mm; CCL: 12.4-13.7 mm; CM³: 4.5-4.8 mm
Philetor brachypterus
- 6B. Tragus small and short; FA: 42.0-45.5 mm; CCL: ~ 15.0mm; CM³: 4.9-5.5 mm
Vespertilio murinus
- 7A. Interfemoral membrane sparsely haired near the body; dorsal pelage long, silky and variably chestnut brown; FA: 32.1-38.0 mm; CCL: 12.4-13.3 mm; CM³: 4.6-5.1 mm
Hypsugo savii
- 7B. Interfemoral membrane mostly naked; dorsal pelage long, silky and dark brown; FA: 32.6-36.5 mm; CCL: 12.6-12.8 mm; CM³: 4.6-4.9 mm
Hypsugo cadornae

Key 7, Family Miniopteridae (3 species)

- 1A. Large with forearm greater than 47mm and condylobasal length greater than 15.5mm; FA: 47.5 -52.8 mm; CBL: 15.7-17.3 mm; CM³: 6.4-7.3 mm

Miniopterus magnater

- 1B. Small with forearm lesser than 50mm and condylobasal length lesser than 15.5mm

2

- 2A. Hairs on the interfemoral membrane near body parts; FA: 44.7-49.6 mm; CBL: 14.6-15.5 mm; CM³: 5.8-6.3 mm

Miniopterus schreibersii

- 2B. Hairs on the interfemoral membrane extend further away from body parts; FA: 39.6-40.2 mm; CBL: 12.7-13.2 mm; CM³: 5.1-5.3 mm

Miniopterus pusillus

Key 8, Family Emballonuridae (6 species)

- 1A. Radio-metacarpal pouch distinct; lower lip scarcely grooved

2

- 1B. Radio-metacarpal pouch absent; lower lip deeply grooved; gular sac present in both the sexes (smaller in females); wings attached to ankles; FA: 63.0-68.2 mm; CCL: 21.7-24.6 mm; CM³: 9.8-11.2 mm

Saccopteryx saccolaimus

- 2A. Radio-metacarpal pouch small; gular pouch less visible; wings attached to tibiae; posterior back and lower abdomen naked; FA: 71.0-80.0 mm; CCL: 21.6 -25.6 mm; CM³: 10.3-11.8 mm

Taphozous nudiventris

- 2B. Radio-metacarpal pouch well-developed

3

- 3A. Gular pouch present; wings attached either to tibiae or ankles

4

- 3B. Gular pouch absent; wings attached to tibiae

5

- 4A. Gular pouch present in males and absent in females; wings attached to tibiae; FA: 59.2-63.8 mm; CCL: 18.4-19.7 mm; CM³: 8.2-8.9 mm

Taphozous perforatus

- 4B. Gular pouch rudimentary in females; wings attached to ankles; posterior back and lower abdomen hairy; FA: 55.6-62.0 mm; CCL: 19.2-21.6 mm; CM³: 8.7-9.2 mm

Taphozous longimanus

5A. Large with forearm length greater than 70mm; wing and interfemoral membrane naked; FA: 71.0- 76.0 mm; CCL: 21.9-23.4 mm; CM³: 9.4-10.5 mm

Taphozous theobaldi

5B. Small with forearm length lesser than 70mm; dorsal fur extends over wing and interfemoral membranes; males with black beard; FA: 60.0-68.0 mm; CCL: 19.5-21.5 mm; CM³: 8.6-9.2 mm

Taphozous melanopogon

Key 9, Family Molossidae (4 species)

1A. Ears not joined by membrane over the forehead

2

1B. Ears joined by membrane over the forehead

3

2A. Forearm less than 52.3mm; two pairs of lower incisors (some aberrant individuals may have three pairs); FA: 46.0-52.3 mm; CCL: 17.1-18.8 mm; CM³: 7.1-7.8 mm

Tadarida aegyptiaca

2B. Forearm greater than 58.4mm; three pairs of lower incisors; FA: 58.4-63.9 mm; CCL: 21.6-23.1 mm; CM³: 8.2-9.3 mm

Tadarida teniotis

3A. Forearm less than 51mm; two pairs of lower incisors; FA: 43.1-50.2 mm; CCL: 15.9-17.1 mm; CM³: 6.4-7.3 mm

Chaerephon plicatus

3B. Forearm greater than 63.0mm; two (or sometimes three) pairs of small and bifid lower incisors; pelage diagnostic with glossy dark chocolate brown on the back, thin white lines on the flanks and antebrachial membrane; FA: 63.0-67.0 mm; CCL: 21.8-23.2 mm; CM³: 8.8-9.3 mm

Otomops wroughtoni

Table 3. Diagnostic morphological characters of eight genera of the family Pteropodidae present in South Asia

| Genus | <i>Rousettus</i> Gray, 1821 | <i>Pteropus</i> Brisson, 1762 | <i>Cynopterus</i> Cuvier, F., 1824 | <i>Megaptera</i> Peters, 1865 | <i>Latidens</i> Thonglongya, 1972 | <i>Sphaerias</i> Miller, 1906 | <i>Eonycteris</i> Dobson, 1873 | <i>Macroglossus</i> Cuvier, F., 1824 |
|---------------------------------------|--|---|--|---|---|--|---|---|
| External characters | | | | | | | | |
| Forearm length | 75.0-90.0 | 110.0-209.0 | 57.0-79.0 | 52.0-63.0 | 66.0-69.0 | 52.0-61.0 | 66.0-78.0 | 44.0-52.0 |
| Head Body length | 111.0-147.0 | 148.0-300.0 | 76.0-113.0 | 81.0-90.0 | 102.0-109.0 | NA | 92.0-130.0 | 78.0-89.0 |
| Tail length | 8.0-21.0 | Absent | 2.0-19.0 | Absent | Absent | Absent | 11.5-23.0 | Absent/ (3.5-5.5) |
| Hind Foot length | 15.0-25.0 | 38.0-65.0 | 11.0-18.0 | ~14.0 | 8.0-15.0 | ~12.0 | 17.0-21.0 | ~10.0 |
| Ear length | 17.5-24.0 | 22.0-57.0 | 14.5-24.0 | 18.5-19.0 | 15.5-18.5 | 16.1-19.5 | 16.9-21.0 | 15.0-17.0 |
| Cranial characters | | | | | | | | |
| Condylabasal length | 33.5-40.4 | 48.6-74.9 | 26.0-33.3 | 25.6-26.7 | 31.4-33.0 | ~26.2 | 31.7-36.3 | 26.6-27.1 |
| Maxillary toothrow (CM ^a) | 13.5-16.6 | 19.0-29.0 | 8.9-12.2 | 8.4-8.6 | 11.3-11.8 | 8.0-8.1 | 11.9-13.4 | 8.6-8.8 |
| Incisors – Upper – Lower | 2 pairs 2 pairs | 2 pairs 2 pairs | 2 pairs 2 pairs | 1 pair 1 pair | 1 pair 1 pair | 2 pairs 2 pairs | 2 pairs 2 pairs | 2 pairs 2 pairs |
| Cheek teeth | Robust | Robust | Robust | Short & broad | Broad | Narrow | Narrow | Extremely narrow |
| II upper molar | Present | Present | Absent | Absent | Absent | Absent | Present | Present |
| Dental formula | 2132 2133 = 34 | 2132 2133 = 34 | 2131 2132 = 30 | 2131 1132 = 28 | 1131 1132 = 26 | 2131 2132 = 30 | 2132 2133 = 34 | 2132 2133 = 34 |
| Species content | <i>R. aegyptiacus</i> (E. Geoffroy, 1810); <i>R. leschenaultii</i> (Desmarest, 1820) | <i>P. giganteus</i> Brünnich, 1782; <i>P. faunulus</i> Miller, 1902; <i>P. hypomelanus</i> Temminck, 1853; <i>P. melanotus</i> Blyth, 1863 | <i>C. sphinx</i> Vahl, 1797; <i>C. brachyotis</i> (Müller, 1838) | <i>M. niphanæ</i> Yenbutra & Felten, 1983 | <i>L. salvini</i> Thonglongya, 1972 | <i>S. blanfordi</i> (Thomas, 1891) | <i>E. spilacea</i> (Dobson, 1871) | <i>M. sobrinus</i> (K. Andersen, 1911) |
| Further table reference | 3.1 | 3.2 | 3.3 | 3.4 | 3.4 | 3.4 | 3.4 | 3.4 |

Note: All measurements are in mm

Table 3.1. Diagnostic morphological characters of the two species of the genus *Rousettus* Gray, 1821 present in South Asia

| Species | <i>Rousettus aegyptiacus</i> (E. Geoffroy, 1810) | <i>Rousettus leschenaultii</i> (Desmarest, 1810) |
|--|---|---|
| External characters | | |
| Forearm length | 83.0-90.0 | 75.0-86.0 |
| Head Body length | 113.0-126.0 | 111.0-147.0 |
| Tail length | 14.0-20.0 | 8.0-21.0 |
| Hind Foot length | 20.0-25.0 | 15.0-22.0 |
| Ear length | 18.0-22.0 | 17.5-24.0 |
| Thumb (including claw) | 30.0-33.4 | 24.0-31.3 |
| II phalanx of 3 rd metacarpal | 51.9-56.3 | 39.6-46.2 |
| Cranial characters | | |
| Condylabasal length | 36.9-40.4 | 33.5-37.7 |
| Maxillary toothrow (CM ²) | 14.9-16.6 | 13.5-15.2 |
| Mandibular toothrow (CM ₃) | 16.5-18.2 | 14.8-16.7 |
| Zygomatic breadth | 23.7-25.2 | 20.2-24.0 |
| Mandible length | 30.5-32.7 | 27.6-31.1 |

Note: All measurements are in mm

Table 3.2. Diagnostic morphological characters of four species of the genus *Pteropus* Brisson, 1762 present in South Asia

| Species | <i>Pteropus giganteus</i> (Brünnich, 1782) | <i>Pteropus faunulus</i> Miller, 1902 | <i>Pteropus hypomelanus</i> Temminck, 1853 | <i>Pteropus melanotus</i> Blyth, 1863 |
|--|--|---|---|--|
| External characters | | | | |
| Forearm length | 145.0-183.0 | 110.0-116.0 | 135.0-145.0 | 148.0-163.0 |
| Head Body length | 198.0-300.0 | ~ 170.0 | 199.0-220.0 | NA |
| Tail length | Absent | Absent | Absent | Absent |
| Hind Foot length | 38.0-58.0 | ~ 34.0 | ~ 59.2 | NA |
| Ear length | 33.0-45.0 Long & pointed, triangular shaped | ~ 22.0 Moderate length, tips rounded off | 25.0-28.0 Short & broad, tips rounded off | NA Large & broad, with narrow tips |
| Rostrum | Long & robust | Narrow | Narrow | Broad & robust |
| Pelage on the back | Blackish-brown with a few paler hairs | Blackish-brown with white hairs | Pale russet brown with gray & black hairs | Blackish-brown with a few gray hairs |
| Ventral surface | Pale tan to deep orange red or chestnut brown | Gray-chestnut brown | Fawn | Dark brown or blackish brown |
| Cranial characters | | | | |
| Condylabasal length | 61.5-74.9 | ~ 48.6 | 59.0-64.1 | 63.3-70.7 |
| Maxillary toothrow (CM ²) | 24.0-29.0 | 19.0-19.2 | 22.8-25.7 | 24.6-27.9 |
| Mandibular toothrow (CM ₃) | 25.6-33.0 | 21.5-22.0 | 25.7-28.6 | 27.8-30.6 |
| Zygomatic breadth | 32.2-44.0 | ~ 28.0 | 32.2-34.8 | 32.9-40.2 |
| Mandible length | 47.8-60.0 | 40.0-42.0 | 48.9-53.2 | 52.6-59.1 |

Note: All measurements are in mm; NA - Data not available

Table 3.3. Diagnostic morphological characters of the two species of the genus *Cynopterus* Cuvier, F., 1824 present in South Asia

| Species | <i>Cynopterus sphinx</i> (Vahl, 1797) | <i>Cynopterus brachyotis</i> (Müller, 1838) |
|---|---|--|
| External characters | | |
| Forearm length | 64.0-79.0 | 57.3-63.3 |
| Head Body length | 76.0-113.0 | 80.0-96.0 |
| Tail length | 4.5-19.0 | 2.0-13.0 |
| Hind Foot length | 12.6-18.0 | 11.0-15.0 |
| Ear length | 17.5-24.0 Large with pale anterior and posterior borders | 14.5-18.0 Small with pale anterior and posterior borders absent or poorly developed |
| Metacarpals & phalanges | Pale | Dark |
| Cranial characters | | |
| Condylar basal length | 28.4-33.3 | 26.0-28.8 |
| Maxillary toothrow (CM ¹) | 10.2-12.2 | 8.9-10.7 |
| Mandibular tooth row (CM ₂) | 12.3-13.5 | 10.0-11.9 |
| Zygomatic breadth | 18.8-23.1 | 17.6-19.8 |
| Mandible length | 22.7-27.5 | 20.5-23.8 |

Note: All measurements are in mm

Table 3.4. Diagnostic morphological characters of the species of the genera *Megaerops* Peters, 1865; *Latidens* Thonlongya, 1972; *Sphaerias* Miller, 1906; *Eonycteris* Dobson, 1873; and *Macroglossus* Cuvier, F., 1824 present in South Asia

| Species | <i>Megaerops niphanae</i> Yenbutra & Felten, 1983 | <i>Latidens salimalii</i> Thonlongya, 1972 | <i>Sphaerias blanfordi</i> (Thomas, 1891) | <i>Eonycteris spelaea</i> (Dobson, 1871) | <i>Macroglossus sobrinus</i> (K. Andersen, 1911) |
|--|--|--|--|---|--|
| External characters | | | | | |
| Forearm length | 52.0-63.0 | 66.0-69.0 | 51.7-60.5 | 66.0-78.0 | 44.0-52.0 |
| Head Body length | 81.0-90.0 | 102.0-109.0 | NA | 92.0-130.0 | 78.0-89.0 |
| Tail length | Absent | Absent | Absent | 11.5-23.0 | Absent/ if present 3.5-5.5 |
| Hind Foot length | ~ 14.0 | 8.0-15.0 | ~ 12.0 | 17.0-21.0 | ~ 10.0 |
| Ear length | 18.5-19.0 Small, broadly rounded | 15.5-18.5 Oval, narrowly rounded tip | 16.1-19.5 Narrowly rounded tip | 16.9-21.0 Narrowly rounded tip | 15.0-17.0 Medium with narrowly rounded tips |
| Interfemoral membrane | Narrow, medial part hairy | Narrow, sparsely haired above and below | Extremely narrow, sparsely haired | Moderately broad sparsely haired along the body | Narrow, thickly covered with short hairs above and below |
| Cranial characters | | | | | |
| Condylar basal length | 25.6-26.7 | 31.4-33.0 | ~ 26.2 | 31.7-36.3 | 26.6-27.1 |
| Maxillary toothrow (CM ¹) | 8.4-8.6 | 11.3-11.8 | 8.0-8.1 | 11.9-13.4* | 8.6-8.8* |
| Mandibular toothrow (CM ₃) | ~ 9.4 | 12.4-12.9 | ~ 9.6 | 13.0-13.7† | 9.7-10.1† |
| Zygomatic breadth | 17.9-18.6 | 21.0-22.1 | ~ 18.0 | 19.0-22.1 | 14.4-14.8 |
| Mandible length | 20.2-21.0 | 24.8-26.3 | ~ 19.4 | 25.1-28.8 | 21.6-22.5 |

Note: All measurements are in mm; * – CM²; † – CM₃

Table 4. Diagnostic morphological characters of the genera of the families Rhinopomatidae, Emballonuridae, Megadermatidae and Rhinolophidae present in South Asia

| Family | Rhinopomatidae | Emballonuridae | Megadermatidae | Rhinolophidae |
|---------------------------------------|---|---|---|--|
| Genus | <i>Rhinopoma</i> E. Geoffroy, 1818 | <i>Taphozous</i> E. Geoffroy, 1818 | <i>Saccopteryx</i> Temminck, 1838 | <i>Megaderma</i> E. Geoffroy, 1810 |
| External characters | | | | |
| Forearm length | 46.0-74.0 | 55.6-80.0 | 63.0-68.2 | 54.0-71.5 |
| Head Body length | 49.0-84.0 | 67.0-105.0 | 80.0-93.0 | 54.0-95.0 |
| Tail length | 49.0-78.0 | 20.0-46.0 | 21.0-35.0 | Absent |
| Hind Foot length | 11.0-18.0 | 8.0-18.0 | 12.0-17.0 | 13.0-20.0 |
| Ear length | 15.8-22.2 | 14.0-25.0 | 16.0-20.0 | 31.5-45.0 Ears joined above the forehead |
| Noseleaf | Absent | Absent | Absent | Present with erect posterior lancel & perpendicular median sella |
| Cranial characters | | | | |
| Condylar canine length | 14.7-22.7 | 18.4-25.6 | 21.7-24.6 | 21.9-27.8 |
| Maxillary toothrow (CM ³) | 5.3-5.9 | 8.2-11.8 | 9.8-11.2 | 9.3-12.1 |
| Incisors – Upper – Lower | 1 pair 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs | Absent 2 pairs |
| Dental formula | 1113 2123 = 28 | 1123 2123 = 30 | 1123 2123 = 28 | 1123 2133 = 32 |
| Species content | <i>R. microphyllum</i> (Brünnich, 1782); <i>R. hardwickii</i> Gray, 1837; <i>R. muscatellum</i> Thomas, 1903 | <i>T. longimanus</i> Hardwicke, 1825; <i>T. melanopogon</i> Temminck, 1841; <i>T. perforatus</i> E. Geoffroy, 1818; <i>T. thebaeus</i> Dobson, 1872; <i>T. nudiventris</i> Cretzschmar, 1830-31 | <i>S. saccolaimus</i> (Temminck, 1838) | <i>M. lyra</i> E. Geoffroy, 1810; <i>M. spasma</i> (Linnaeus, 1758) |
| Further table reference | 4.1 | 4.2 | 4.2 | 4.3 |
| | | | | 4.4 |

Note: All measurements are in mm

Table 4.1. Diagnostic morphological characters of the three species of the genus *Rhinopoma* E. Geoffroy, 1818 present in South Asia

| Species | <i>Rhinopoma microphyllum</i> (Brünnich, 1782) | <i>Rhinopoma hardwickii</i> Gray, 1831 | <i>Rhinopoma muscatellum</i> Thomas, 1903 |
|---|---|---|--|
| External characters | | | |
| Forearm length | 59.5-74.6 | 52.9-64.0 | 46.0-53.2 |
| Head Body length | 60.0-84.0 | 55.0-73.0 | 49.0-62.0 |
| Tail length | 50.0-77.0 | 56.0-78.0 | 49.0-64.3 |
| Hind Foot length | 14.0-18.0 | 11.0-15.0 | 11.0-13.0 |
| Ear length | 18.0-22.0 | 17.0-21.0 | 15.8-20.0 |
| Dermal ridge | Poorly developed | Well-developed | Poorly developed |
| Tail length in relation to forearm length | Shorter | Longer | Longer |
| Cranial characters | | | |
| Condylar canine length | 17.2-22.7 | 15.5-17.5 | 14.7-15.1 |
| Maxillary toothrow (CM ³) | 7.0-8.0 | 6.0-6.8 | 5.3-5.9 |
| Mandibular tooth row (CM ₃) | 7.6-8.6 | 6.5-7.5 | 5.1-6.7 |
| Zygomatic breadth | 11.4-13.4 | 10.1-11.7 | 8.8-9.8 |
| Mandible length | 13.7-15.8 | 11.8-13.6 | 10.5-11.4 |
| Nasal inflations | Small | Moderate | Large |

Note: All measurements are in mm

Table 4.3. Diagnostic morphological characters of the two species of the genus *Megaderma* E. Geoffroy, 1810 present in South Asia

| Species | <i>Megaderma lyra</i> E. Geoffroy, 1810 | <i>Megaderma spasma</i> (Linnaeus, 1758) |
|---|--|--|
| External characters | | |
| Forearm length | 56.0-71.5 | 54.0-62.0 |
| Head Body length | 70.0-95.0 | 54.0-81.0 |
| Tail length | Absent | Absent |
| Hind Foot length | 14.0-20.0 | 13.0-17.0 |
| Ear length | 31.5-45.0 Oval with white inner margin, joined by membrane over the forehead about 1/3 to 1/2 of its length | 33.0-40.0 Oval lacking white inner margin, joined over the forehead near the base |
| Tragus | Bifid with taller posterior process | Bifid with narrower and taller posterior process |
| Noseleaf – sides – longitudinal ridge – base | Tall (~ 10 mm) Straight Present Simple rounded horizontal | Short (~ 6.5 mm) Convex Present Heart-shaped |
| Cranial characters | | |
| Condylar canine length | 24.5-27.8 | 21.9-23.6 |
| Maxillary toothrow (CM ³) | 10.6-12.1 | 9.3-10.5 |
| Mandibular toothrow (CM ₃) | 11.6-12.2 | 10.2-11.4 |
| Zygomatic breadth | 15.4-17.1 | 13.6-14.8 |
| Mandible length | 18.8-21.2 | 16.8-18.5 |

Note: All measurements are in mm

Table 4.2. Diagnostic morphological characters of five species of the genera *Taphozous* E. Geoffroy, 1818 and *Saccopteryx* Lesson, 1842 present in South Asia

| Species | <i>Taphozous longimanus</i> Hardwicke, 1825 | <i>Taphozous melanopogon</i> Temminck, 1841 | <i>Taphozous perforatus</i> E. Geoffroy, 1818 | <i>Taphozous theobaldi</i> Dobson, 1872 | <i>Taphozous nudiventris</i> Cretzschmar, 1830-31 | <i>Saccopteryx saccostoma</i> (Temminck, 1838) |
|--|--|--|--|--|--|---|
| External characters | | | | | | |
| Forearm length | 55.6-62.0 | 60.0-68.0 | 59.2-63.8 | 71.0-76.0 | 71.0-80.0 | 63.0-68.2 |
| Head Body length | 73.0-86.0 | 67.0-86.0 | 71.0-80.0 | 88.0-89.0 | 90.0-105.0 | 80.0-93.0 |
| Tail length | 20.0-30.0 | 20.0-32.0 | 20.0-28.0 | 25.0-35.0 | 22.0-46.0 | 21.0-35.0 |
| Hind Foot length | 8.0-14.0 | 8.2-14.2 | 8.2-12.5 | 11.0-13.0 | 11.0-18.0 | 12.0-17.0 |
| Ear length | 16.0-19.0 | 16.5-22.0 | 14.0-20.0 | 21.0-24.0 | 18.0-25.0 | 16.0-20.0 |
| Chin | Naked | Hairy, males with black beard | Hairy | Hairy | Naked | Short hairs |
| Gular sac in males In females | Present Rudimentary | Absent Absent | Variably present Absent | Absent Absent | Present Present | Present Present |
| Wings attached to | Ankles | Tibiae | Tibiae | Tibiae | Tibiae | Ankles |
| Posterior back & lower abdomen | Hairy | Hairy | Hairy | Hairy | Naked | Hairy |
| III metacarpal length in relation to forearm length | Long (95.8-109.4%) | Short (84.9-92.7%) | Short (87.1-92.7%) | NA | NA | NA |
| Radio-metacarpal pouch | Absent | Absent | Absent | Present | Present | Absent |
| Cranial characters | | | | | | |
| Condylar canine length | 19.2-21.6 | 19.5-21.5 | 18.4-19.7 | 21.9-23.4 | 21.6-25.6 | 21.7-24.6 |
| Maxillary toothrow (CM ³) | 8.7-9.2 | 8.6-9.2 | 8.2-8.9 | 9.4-10.5 | 10.3-11.8 | 9.8-11.2 |
| Mandibular toothrow (CM ₃) | 9.4-10.2 | 9.3-10.2 | 9.0-9.6 | 10.7-12.0 | 11.4-13.3 | 10.8-12.3 |
| Zygomatic breadth | 12.0-12.9 | 12.0-13.1 | 11.5-12.1 | 13.2-14.0 | 14.4-17.8 | 14.2-15.6 |
| Mandible length | 15.4-16.4 | 15.5-16.5 | 14.6-15.6 | 18.0-18.9 | 18.2-21.5 | 17.5-19.8 |

Note: All measurements are in mm; NA - Data not available

Table 4.4. Diagnostic morphological characters of twenty species of the genus *Rhinolophus* Lacépède, 1799 present in South Asia

| Species | <i>Rhinolophus ferrumequinum</i> (Schreber, 1774) | <i>Rhinolophus bocharicus</i> Kastchenko & Akimov, 1917 | <i>Rhinolophus affinis</i> Horsfield, 1823 | <i>Rhinolophus rouxii</i> Temminck, 1835 | <i>Rhinolophus simicus</i> Andersen, 1905 | <i>Rhinolophus hippocideros</i> (Bechstein, 1800) |
|--|---|---|---|--|---|---|
| External characters | | | | | | |
| Forearm length | 54.8-62.1 | 47.0-53.0 | 50.0-55.0 | 44.4-52.3 | 45.7-49.6 | 35.9-40.0 |
| Head Body length | 56.0-79.0 | NA | 46.0-68.0 | 42.0-66.0 | 43.0-52.5 | 38.0-48.0 |
| Tail length | 30.0-42.0 | 22.0-32.0 | 20.0-30.0 | 22.0-33.0 | 21.5-30.0 | 22.0-32.0 |
| Hind Foot length | 9.9-14.0 | NA | 9.8-12.5 | 7.2-12.8 | 7.5-10.0 | 6.9-9.0 |
| Ear length | 20.0-28.5 | 19.3-24.0 | 14.0-23.0 | 14.5-22.0 | 15.8-20.0 | 13.9-19.0 |
| Horseshoe - Sella - Superior connecting Process | Narrow Relatively small Bluntly rounded | Narrow Small, constricted in mid Bluntly rounded | Relatively broad Relatively small Bluntly rounded | Broad Relatively small Bluntly rounded | Broad Relatively small Bluntly rounded | Broadly rounded off and low |
| Inferior extremity | Bluntly rounded | Bluntly rounded | Bluntly rounded | Bluntly rounded | Bluntly rounded, slightly projecting forward | Projects downwards and forwards |
| Lancelet | Narrowly pointed with concave sides | Narrowly pointed with concave sides | Narrowly pointed with concave sides | Triangular with straight sides | Triangular with straight sides | Tall and narrow with straight sides |
| Cranial characters | | | | | | |
| Condylar canine length | 19.7-22.3 | 19.0-20.2* | 18.7-20.5 | 17.5-21.0 | 17.4-18.4 | 13.0-14.0 |
| Maxillary toothrow (CM ³) | 8.5-9.6 | 6.7-7.2 | 8.5-9.5 | 7.9-9.2 | 7.6-8.2 | 5.4-6.0 |
| Mandibular toothrow (CM ₃) | 9.1-10.1 | 7.3-7.8 | 9.0-10.0 | 8.3-10.0 | 8.1-9.00 | 5.9-6.3 |
| Zygomatic breadth | 11.0-13.1 | 9.7-10.8 | 10.4-11.8 | 10.2-11.9 | 10.1-11.0 | 7.1-7.8 |
| Mandible length | 15.2-17.9 | 12.4-13.4 | 14.9-16.1 | 13.8-16.4 | 13.4-14.6 | 9.3-10.0 |

Note: All measurements are in mm; * - GTL

Contd...

Table 4.4. (Contd..) Diagnostic morphological characters of twenty species of the genus *Rhinolophus* Lacépède, 1799 present in South Asia

| Species | <i>Rhinolophus pusillus</i> Temminck, 1834 | <i>Rhinolophus subbadius</i> Blyth, 1844 | <i>Rhinolophus lepidus</i> Blyth, 1844 | <i>Rhinolophus shortridgei</i> Andersen, 1918 | <i>Rhinolophus cognatus</i> K. Andersen, 1906 | <i>Rhinolophus blasii</i> Peters, 1867 |
|--|---|---|--|--|---|--|
| External characters | | | | | | |
| Forearm length | 34.9-37.8 | 33.4-35.9 | 37.0-41.8 | 39.0-42.0 | 37.5-40.7 | 43.0-50.0 |
| Head Body length | 30.0-40.0 | 35.0-37.0 | 35.0-54.0 | NA | NA | 47.0-54.0 |
| Tail length | 13.5-26.0 | 16.0-19.0 | 14.0-28.0 | 16.0-19.0 | 13.0-21.0 | 21.0-35.0 |
| Hind Foot length | 6.6-8.0 | 6.7-8.0 | 5.5-10.0 | NA | 6.5-9.0 | 8.0-11.0 |
| Ear length | 15.0-17.5 | 14.1-18.0 | 14.5-20.6 | 14.0-18.0 | 13.5-18.3 | 15.8-22.0 |
| Horseshoe - Sella - Superior connecting process | Broad Small Broadly triangular (sometimes horn-like) | Broad Small Bluntly rounded (more horn-like) | Broad Small Bluntly rounded | Broad Small Bluntly rounded (more or less horn-like) | Relatively larger Bluntly rounded (horn-like, slightly curved downwards) | Broad Large Acute and high |
| Inferior extremity | Bluntly rounded, not projected downward | Bluntly rounded not projected downward | Bluntly rounded not projected downwards | Bluntly rounded not projected downwards | Bluntly rounded not projected downwards | Acute and projects slightly downwards |
| Lanceal | Equilateral triangle to elongate | Short & broad | Tip broadly rounded or pointed, sides deeply concave to straight | Tip broadly rounded or pointed, sides deeply concave to straight | Tall | Bluntly pointed, sides straight and wide at the base |
| Cranial characters | | | | | | |
| Condylar spine length | 13.0-14.0 | 11.9-12.9 | 13.3-15.5 | 17.2-17.9* | 14.2-15.8 | 16.0-17.0 |
| Maxillary toothrow (CM^3) | 5.4-6.0 | 5.1-5.5 | 5.6-6.8 | 6.5-7.2 | 6.0-7.0 | 6.0-7.6 |
| Mandibular toothrow (CM_3) | 5.9-6.3 | 5.2-5.8 | 6.0-7.4 | 7.0-7.3 | 6.2-7.7 | 6.9-7.2 |
| Zygomatic breadth | 7.1-7.8 | NA | 7.6-8.8 | 8.5-8.9 | 8.0-9.0 | 8.8-9.5 |
| Mandible length | 9.3-10.0 | 9.2-9.5 | 10.0-12.1 | 11.7-12.8 | 10.2-12.5 | 11.9-12.7 |

Note: All measurements are in mm; NA – Data not available; * - GTL

Contd...

Table 4.4. (Contd..) Diagnostic morphological characters of twenty species of the genus *Rhinolophus* Lacépède, 1799 present in South Asia

| Species | <i>Rhinolophus macrotis</i> Blyth, 1844 | <i>Rhinolophus mehelyi</i> Matschie, 1901 | <i>Rhinolophus luctus</i> Temminck, 1835 | <i>Rhinolophus beddomei</i> Andersen, 1905 | <i>Rhinolophus trifoliatus</i> Temminck, 1834 |
|--|---|--|---|---|--|
| External characters | | | | | |
| Forearm length | 39.0-46.0 | 47.0-53.0 | 70.0-80.5 | 54.9-64.3 | 50.0-53.3 |
| Head Body length | 39.0-48.7 | - | 85.0-90.0 | 65.0-75.0 | 62.0-65.0 |
| Tail length | 17.3-26.2 | 16.2-37.0 | 49.0-55.0 | 38.5-48.0 | 30.0-35.0 |
| Hind Foot length | 7.5-9.0 | - | 16.5-20.0 | 13.3-18.0 | ~13.0 |
| Ear length | 21.0-25.9 | 18.0-23.0 | ~40.0 | 21.0-34.0 | ~27.0 |
| Horseshoe - Sella - Superior connecting Process | Broad Small, projects strongly Broadly rounded | Narrow Moderate Broadly rounded | Very broad with emargination, noseleaf has basal lappets Moderate Broadly rounded, very reduced | Very broad, divided in two halves noseleaf has basal lappets Moderate Broadly rounded, very reduced | Broad divided in two halves noseleaf has basal lappets Moderate Broadly rounded very reduced |
| Inferior extremity | Short and emarginated | Broad | Broad | Moderately broad projecting forward and downwards | Narrow |
| Lancet | Short with pointed tip, with slightly convex sides | Abruptly narrow, linear tip | Well-developed, tip rounded | Well-developed, tip rounded | Well-developed, tip rounded |
| Cranial characters | | | | | |
| Condylar canine length | 15.2-16.6 | 18.8-20.3* | 27.5-28.7 | 22.2-24.4 | 19.5-21.5 |
| Maxillary toothrow (CM^3) | 6.2-6.7 | 6.2-7.2 | 11.8-12.6 | 9.6-10.4 | 8.4-8.9 |
| Mandibular toothrow (CM_3) | 6.3-6.9 | 6.9-7.7 | 12.7-13.3 | 10.3-11.1 | 8.7-9.7 |
| Zygomatic breadth | 7.9-8.6 | 10.0-10.9 | 14.8-16.2 | 13.1-14.6 | 11.2-12.5 |
| Mandible length | 10.9-12.7 | 11.8-13.2 | 22.6-23.2 | 17.9-19.3 | 15.0-16.8 |

Contd...

Note: All measurements are in mm; * - GTL

Table 4.4. (Contd..). Diagnostic morphological characters of twenty species of the genus *Rhinolophus* Lacépède, 1799 present in South Asia

| Species | <i>Rhinolophus pearsonii</i> Horsfield, 1851 | <i>Rhinolophus yunanensis</i> Dobson, 1872 | <i>Rhinolophus mitratus*</i> Blyth, 1844 |
|--|---|---|---|
| External characters | | | |
| Forearm length | 50.8-54.8 | 54.2-59.2 | ~ 57.1 |
| Head Body length | 51.0-64.0 | 60.5-68.0 | NA |
| Tail length | 20.0-29.0 | 21.5-26.0 | ~ 30.0 |
| Hind Foot length | 9.9-14.2 | 12.5-14.0 | NA |
| Ear length | 23.5-28.0 | 23.5-28.5 | NA |
| Horseshoe - Sella - Superior connecting process | Broad Moderate Bluntly rounded | Broad Moderate Bluntly rounded | Broad Moderate Bluntly rounded |
| Inferior extremity | Not deflected downwards or Forwards | Not deflected downwards or forwards | Narrow |
| Lancet | Well-developed tip rounded | Well-developed tip rounded | Well-developed tip rounded |
| Cranial characters | | | |
| Condylar canine length | 19.7-22.3 | 22.3-22.9 | NA |
| Maxillary toothrow (CM ³) | 8.5-9.6 | 9.5-10.6 | NA |
| Mandibular toothrow (CM ₃) | 9.1-10.1 | 10.8-11.5 | NA |
| Zygomatic breadth | 11.0-13.1 | 12.2-13.1 | NA |
| Mandible length | 15.2-17.9 | 17.7-18.3 | NA |

Note: All measurements are in mm; * - Known only from type specimen with damaged skull; NA – Data not available

Table 5. Diagnostic morphological characters of the genera of the family Hipposideridae present in South Asia

| Genus | <i>Hipposideros</i> Gray, 1831 | <i>Triaenops</i> Dobson, 1871 | <i>Asellia</i> Gray, 1838 | <i>Coelops</i> Blyth, 1848 |
|---------------------------------------|---|--|--|--|
| External characters | | | | |
| Forearm length | 33.0-99.0 | 48.3-54.0 | 50.1-52.3 | 37.8-42.0 |
| Head Body length | 38.0-106.0 | 51.0-64.0 | NA | NA |
| Tail length | 20.0-64.0 | 31.0-39.0 | 25.0-28.0 | Rudimentary (c' 2 mm) |
| Hind Foot length | 5.3-20.0 | 8.0-11.0 | NA | NA |
| Ear length | 13.0-34.0 | 13.0-16.0 | NA | NA |
| Noseleaf | Present with or without supplementary leaflets | Present with single pair of supplementary leaflets | Present with two pairs of supplementary leaflets | Present with two elongated supplementary lappets |
| Interfemoral membrane | Wide | Wide | Wide | Narrow |
| Cranial characters | | | | |
| Condylar canine length | 12.7-31.5 | 16.9-18.9* | 15.7-17.0 | 14.3-14.8 |
| Maxillary toothrow (CM ³) | 4.9-14.5 | 6.1-7.1 | 6.5-7.2 | 5.8-6.0 |
| Incisors – Upper – Lower | 1 pair 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs |
| Dental formula | 1123 2123 = 30 | 1123 2123 = 30 | 1113 2123 = 28 | 1123 2123 = 30 |
| Species content | <i>H. ater</i> , Templeton, 1848; <i>H. cineraceus</i> Blyth, 1853; <i>H. durgadasi</i> Khajuria, 1970; <i>H. fulvus</i> Gray, 1838; <i>H. pomona</i> K. Andersen, 1918; <i>H. hypophyllus</i> Kock & Bhat, 1994; <i>H. galeritus</i> Cantor, 1846; <i>H. speoris</i> (Schneider, 1800); <i>H. larvatus</i> (Horsfield, 1823); <i>H. armiger</i> Hodgson, 1835; <i>H. lankadiva</i> Kelaart, 1850; <i>H. diadema</i> Geoffroy, E., 1813 | <i>T. persicus</i> Dobson, 1871 | <i>A. tridens</i> Geoffroy, E., 1813 | <i>C. frithii</i> Blyth, 1848 |
| Further table reference | 5.1 | 5.2 | 5.2 | 5.2 |

Note: All measurements are in mm; * - Condylar basal length; NA – Data not available

Table 5.1. Diagnostic morphological characters of twelve species of the genus *Hipposideros* Gray, 1831 present in South Asia

| Species | <i>Hipposideros ater</i> Templeton, 1848 | <i>Hipposideros cinereus</i> Blyth, 1853 | <i>Hipposideros durgadasi</i> Khajuria, 1970 | <i>Hipposideros fulvus</i> Gray, 1838 | <i>Hipposideros pomona</i> K. Andersen, 1918 | <i>Hipposideros hypophyllus</i> Kock & Bhat, 1994 |
|--|---|---|---|--|---|--|
| External characters | | | | | | |
| Forearm length | 34.9-42.3 | 33.0-36.3 | 36.0-37.5 | 38.4-44.0 | 38.1-43.2 | 37.1-40.0 |
| Head Body length | ~38.0-48.0 | 33.0-42.0 | NA | 40.0-50.0 | ~36.0-52.0 | 42.0-47.3 |
| Tail length | 20.0-30.0 | 22.0-30.0 | 21.5-29.0 | 24.0-35.0 | ~28.0-35.0 | 21.5-23.5 |
| Hind Foot length | 5.3-7.2 | 6.0-7.0 | 5.5-8.0 | 6.0-9.8 | ~6.3-8.5 | 6.2-6.9 |
| Ear length | 14.8-20.0 | 13.0-17.0 | 13.0-19.0 | 19.0-26.0 | 18.5-25.0 | 18.0-20.3 |
| No. of supplementary leaflets | 0 | 0 | 0 | 0 | 0 | 1 |
| Median emargination on anterior leaf | Absent | Absent | Absent | Absent | Absent | Absent |
| Narial lappets | Absent | Absent | Absent | Absent | Absent | Absent |
| Cranial characters | | | | | | |
| Condylar canine length | 13.2-15.0 | 12.7-13.7 | 13.0-13.9* | 15.0-16.4 | 14.2-16.0 | 14.2-14.8 |
| Maxillary toothrow (CM ³) | 5.1-5.9 | 4.9-5.3 | 5.0-6.0 | 6.0-6.9 | 5.5-6.2 | 5.3-5.7 |
| Mandibular toothrow (CM ₃) | 5.2-6.4 | 5.2-5.8 | 5.0-6.0 | 6.4-7.5 | 6.0-6.8 | 5.6-6.1 |
| Zygomatic breadth | 7.7-8.7 | 6.9-7.6 | 6.8-9.0 | 8.6-9.6 | 7.9-9.0 | 7.8-8.3 |
| Mandible length | 9.4-10.6 | 8.8-9.4 | 9.0-9.5 | 11.1-12.0 | 9.8-11.3 | 9.5-10.6 |
| Dorsal process on jugal bone of zygoma | Present | Absent | Absent | NA | NA | Present |
| Rostral eminences | Present, 4, of which 2 less developed | Present, 4 all less developed | Present, 4 all less developed | NA | NA | NA |

Note: All measurements are in mm; * - Condylarbasal length; NA - Data not available

Contd...

Table 5.1 (Contd..) Diagnostic morphological characters of twelve species of the genus *Hipposideros* Gray, 1831 present in South Asia

| Species | <i>Hipposideros galeritus</i> Cantor, 1846 | <i>Hipposideros speoris</i> (Schneider, 1800) | <i>Hipposideros larvatus</i> (Horsfield, 1823) | <i>Hipposideros armiger</i> Hodgson, 1835 | <i>Hipposideros lankadiva</i> Kelaart, 1850 | <i>Hipposideros diadema</i> Geoffroy, E., 1813 |
|--|---|--|---|--|---|---|
| External characters | | | | | | |
| Forearm length | 45.0-51.3 | 45.6-54.0 | 61.2-64.8 | 85.4-95.0 | 75.0-99.0 | 58.8-64.2 |
| Head Body length | 45.0-59.5 | 46.0-62.0 | 74.0-78.0 | 82.0-105.0 | 87.0-106.0 | NA |
| Tail length | 29.5-37.0 | 20.0-29.0 | 37.0-44.0 | 50.0-64.0 | 35.0-58.0 | NA |
| Hind Foot length | 4.9-8.0 | 7.0-11.0 | NA | 13.0-17.0 | 12.0-20.0 | NA |
| Ear length | 14.5-17.0 | 12.5-19.0 | 23.0-26.0 | 26.0-34.0 | 19.5-30.0 | NA |
| No. of supplementary Leaflets | 2 | 3 rd much reduced | 3 | 4 th | 4 th much reduced, sometimes absent | 4 th much reduced, sometimes absent |
| Median emargination on anterior leaf | Absent | Present, faintly seen | Present, faintly seen | Absent | Absent | Absent |
| Narial apes | Small | Well-developed | Well-developed | Less developed | Well-developed | Well-developed |
| Cranial characters | | | | | | |
| Condylar canine length | 14.6-15.8 | 15.9-17.5 | 20.3-21.4 | 25.6-29.2 | 24.8-31.5 | 22.6-22.9 |
| Maxillary toothrow (CM ³) | 5.3-6.2 | 6.6-7.5 | 8.9-9.4 | 11.3-12.5 | 11.6-14.5 | 9.7-10.0 |
| Mandibular toothrow (CM ₃) | 5.9-6.3 | 7.2-8.3 | 9.9-10.5 | 12.6-14.0 | 12.6-16.5 | NA |
| Zygomatic breadth | 8.4-9.3 | 10.2-11.6 | 13.0-14.0 | 16.5-18.4 | 16.1-20.7 | 14.5-15.0 |
| Mandible length | 10.0-11.4 | 12.4-13.9 | 16.3-16.8 | 20.7-23.4 | 20.1-26.2 | NA |
| Dorsal process on jugal bone of zygoma | Present | Present | Present | Present | Present | NA |
| Rostral eminences | Present, 4 all well-developed | Present, 4 all less developed | Present, 4 all well-developed | NA all well-developed | Present, 4 all well-developed | NA |

Note: All measurements are in mm; NA – Data not available

Table 5.2. Diagnostic morphological characters of one species each of the genera *Triaenops* Dobson, 1871; *Asellia* Gray, 1838; and *Coelops* Blyth, 1848 present in South Asia

| Species | <i>Triaenops persicus</i> Dobson, 1871 | <i>Asellia tridens</i> Geoffroy, E., 1813 | <i>Coelops frithii</i> Blyth, 1848 |
|--|---|--|--|
| External characters | | | |
| Forearm length | 48.3-54.0 | 50.1-52.3 | 37.8-42.0 |
| Head Body length | 51.0-64.0 | 52.0-55.0 | 38.0-50.0 |
| Tail length | 31.0-39.0 | 25.0-28.0 | Rudimentary, <2.0 |
| Hind Foot length | 8.0-11.0 | 9.0-10.0 | 5.3-9.0 |
| Ear length | 11.0-16.0 Small, funnel-shaped | 19.0-20.0 Broad with pointed tips | 11.5-15.0 Broad with rounded tip |
| Noseleaf | Complex with three vertical pointed processes | Complex with three vertical processes, central one pointed, others blunt | Complex with anterior leaf divided in to two and with forwardly projecting supplementary lappets |
| No. of supplementary leaflets | 1 | 2 | 0 |
| Median emargination on anterior leaf | Deep | Absent | Deep |
| Narial lappets | Well-developed | Less developed | Well-developed |
| Cranial characters | | | |
| Condylarcanine length | 16.9-18.9* | 15.7-17.0 | 14.3-14.8 |
| Maxillary toothrow (CM ³) | 6.1-7.1 | 6.5-7.2 | 5.8-6.0 |
| Mandibular toothrow (CM ₃) | 7.0-7.7 | 7.3-8.3 | 6.0-6.5 |
| Zygomatic breadth | 8.7-9.5 | 9.7-10.7 | 7.2-7.5 |
| Mandible length | 11.7-13.1 | 12.2-13.2 | 9.8-10.2 |
| Dorsal process on jugal bone of zygoma | Present | Present, less developed | Absent |
| Rostral eminences | Present, 4, all well-developed | Present, 4, all well-developed | Present, 4, all less developed |

Note: All measurements are in mm; * - Condylarbasal length; NA – Data not available

Table 6. Diagnostic morphological characters of the genera of the family Molossidae present in South Asia

| Genus | <i>Tadarida</i> Rafinesque, 1814 | <i>Chaerephon</i> Dobson, 1874 | <i>Otomops</i> Thomas, 1913 |
|---------------------------------------|--|---|---|
| External characters | | | |
| Forearm length | 46.0-63.9 | 43.1-50.2 | 63.0-67.0 |
| Head Body length | 61.0-90.0 | 66.0-71.0 | 87.0-99.0 |
| Tail length | 33.0-60.0 | 30.0-44.0 | 41.0-49.0 |
| Hind Foot length | 7.0-12.1 | 9.0-11.3 | 10.0-14.0 |
| Ear length | 15.0-32.0 Not joined over forehead | 16.0-21.0 Joined by membrane over forehead | 31.2-34.0 Joined by membrane over forehead |
| Cranial characters | | | |
| Condylar canine length | 17.1-23.1 | 15.9-17.1 | 21.8-23.2 |
| Maxillary toothrow (CM ³) | 7.1-9.3 | 6.4-7.3 | 8.8-9.3 |
| Incisors – Upper – Lower | 1 pair 2/3 pairs | 1 pair 2 pairs | 1 pair 2/3 pairs |
| Dental formula | 1123 2(3)123 = 30/32 | 1123 2123 = 30 | 1123 2(3)123 = 30/32 |
| Species content | <i>T. teniotis</i> (Rafinesque, 1814); <i>T. aegyptiaca</i> (E. Geoffroy, 1818) | <i>C. plicatus</i> (Buchanan, 1800) | <i>O. wroughtoni</i> Thomas, 1913 |
| Further table reference | 6.1 | 6.1 | 6.1 |

Note: All measurements are in mm

Table 6.1. Diagnostic morphological characters of the species of the genera *Tadarida* Rafinesque, 1814, *Chaerephon* Dobson, 1874 and *Otomops* Thomas, 1913 present in South Asia

| Genus | <i>Tadarida teniotis</i> (Rafinesque, 1814) | <i>Tadarida aegyptiaca</i> (E. Geoffroy, 1818) | <i>Chaerephon plicatus</i> (Buchanan, 1800) | <i>Otomops wroughtoni</i> Thomas, 1913 |
|--|---|--|---|---|
| External characters | | | | |
| Forearm length | 58.4-63.9 | 46.0-52.3 | 43.1-50.2 | 63.0-67.0 |
| Head Body length | 74.0-90.0 | 61.0-77.0 | 66.0-71.0 | 87.0-99.0 |
| Tail length | 43.0-54.8 | 33.0-60.0 | 30.0-44.0 | 41.0-49.0 |
| Hind Foot length | 9.0-12.1 | 7.0-10.0 | 9.0-11.3 | 10.0-14.0 |
| Ear length | 25.0-32.0 Not joined over forehead | 15.0-23.0 Not joined over forehead | 16.0-21.0 Joined over forehead by a membrane | 31.2-34.0 Joined over forehead by a membrane |
| Antitragus | Large | Small | Small | Absent |
| Cranial characters | | | | |
| Condylar canine length | 21.6-23.1 | 17.1-18.8 | 15.9-17.1 | 21.8-23.2 |
| Maxillary toothrow (CM ³) | 8.2-9.3 | 7.1-7.8 | 6.4-7.3 | 8.8-9.3 |
| Mandibular toothrow (CM ₃) | 9.1-9.8 | 7.7-8.6 | 6.9-7.9 | 9.2-9.8 |
| Zygomatic breadth | 13.4-14.2 | 11.3-13.3 | 10.9-11.4 | 12.6-13.3 |
| Mandible length | 16.6-17.5 | 13.3-14.6 | 11.8-13.4 | 15.9-16.9 |
| Lower Incisors | 3 pairs | 2 pairs | 2 pairs | 2/3 pairs |

Note: All measurements are in mm

Table 7. Diagnostic morphological characters of the genera of the subfamilies Murininae, Kerivoulinae and Myotinae of family Vespertilionidae present in South Asia

| Genus | <i>Murina</i> Gray, 1842 | <i>Harpiola</i> Thomas, 1915 | <i>Harpiocephalus</i> Gray, 1842 | <i>Kerivoula</i> Gray, 1842 | <i>Myotis</i> Kaup, 1829 |
|---------------------------------------|--|--|--|---|---|
| External characters | | | | | |
| Forearm length | 27.7-40.9 | 32.4-34.2 | 44.1-50.1 | 31.5-42.0 | 30.0-58.3 |
| Head Body length | 38.0-50.0 | 35.6-42.7 | 60.0-75.0 | 39.0-55.0 | 38.0-80.0 |
| Tail length | 27.0-41.0 | 27.5 | 40.0-50.0 | 35.0-55.0 | 25.0-68.0 |
| Hind Foot length | 4.5-10.0 | 8.2 | 11.0-14.0 | 4.0-9.0 | 6.7-17.0 |
| Ear length | 13.0-17.0 Short, broad and tip rounded | 12.1 Short, broad, triangular and tip rounded | 17.0-18.0 Rounded | 11.0-17.0 Moderately long and funnel shaped | 6.0-19.0 Narrow and tip bluntly rounded |
| Cranial characters | | | | | |
| Condylar canine length | 12.1-16.3 | 12.1-14.8 | 19.0-19.5 | 12.2-15.9 | 11.2-18.9 |
| Maxillary toothrow (CM ³) | 4.5-6.1 | 4.9-5.3 | 6.5-6.9 | 5.5-7.4 | 4.5-8.5 |
| Incisors – Upper – Lower | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs |
| Premolars – Upper – Lower | 2 pairs 2 pairs | 2 pairs 2 pairs | 2 pairs 2 pairs | 3 pairs 3 pairs | 2-3 pairs 2-3 pairs |
| Dental formula | -23,1,-2-4,123 123,1,-2-4,123 = 34 | -23,1,-2-4,123 123,1,-2-4,123 = 34 | -23,1,-2-4,123 123,1,-2-4,123 = 34 | -23,1,-234,123 123,1,-234,123 = 38 | 213(2)3 313(2)3 = 38 / 34 |
| Species content | <i>M. leucogaster</i> Milne-Edwards, 1872; <i>M. aurata</i> Milne-Edwards, 1872; <i>M. cyclotis</i> Dobson, 1872; <i>M. tubinaris</i> (Scully, 1881); <i>M. huttoni</i> (Peters, 1872); | <i>H. grisea</i> (Peters, 1872) | <i>H. harpia</i> (Temminck, 1840) | <i>K. picta</i> (Pallas, 1767); <i>K. hardwickii</i> (Horsfield, 1825); <i>K. lenis</i> Thomas, 1916 | <i>M. blythii</i> (Tomes, 1857); <i>M. sicarius</i> Thomas, 1915; <i>M. formosus</i> (Hodgson, 1835); <i>M. nipalensis</i> (Dobson, 1871); <i>M. muricola</i> (Gray, 1846); <i>M. siligorensis</i> (Horsfield, 1855); <i>M. montivagus</i> (Dobson, 1874); <i>M. annectans</i> (Dobson, 1871); <i>M. longipes</i> (Dobson, 1873); <i>M. laniger</i> (Peters, 1871); <i>M. csorbai</i> Topal, 1997; <i>M. horsfieldii</i> (Temminck, 1814); <i>M. hasseltii</i> (Temminck, 1840) |
| Further table reference | 7.1 | 7.1 | 7.1 | 7.2 | 7.3 |

Note: All measurements are in mm

Contd...

Table 7 (Contd.). Diagnostic morphological characters of the genera of the tribes Plecotini, Nycticeiini and Eptesicini of Subfamily Vespertilioninae of family Vespertilionidae present in South Asia

| Genus | <i>Plecotus</i> Geoffroy, E., 1818 | <i>Barbastella</i> Gray, 1821 | <i>Otonycteris</i> Peters, 1859 | <i>Scotomantoides</i> Dobson, 1875 | <i>Scotophilus</i> Leach, 1821 | <i>Scotoecus</i> Thomas, 1901 | <i>Arielulus</i> Hill & Harrison, 1987 | <i>Hesperoptenus</i> Peters, 1869 | <i>Eptesicus</i> Rafinesque, 1820 |
|---------------------------------------|---|--|--------------------------------------|---------------------------------------|---|-------------------------------------|--|--------------------------------------|---|
| External characters | | | | | | | | | |
| Forearm length | 36.5-45.1 | 38.7-42.1 | 64.7-65.6 | 56.1-61.2 | 44.0-65.8 | 34.1-37.3 | 41.8-43.6 | 50.0-60.4 | 35.4-55.1 |
| Head Body length | 40.0-53.0 | 47.0-51.0 | 61.0-76.0 | 64.0-85.0 | 60.0-93.0 | 50.0-58.0 | 92.0-95.0 | 61.0-79.0 | 40.0-80.0 |
| Tail length | 48.0-54.0 | 40.0-47.0 | 40.0-58.0 | 52.0-66.0 | 40.0-71.0 | 34.0-41.0 | 39.0-40.0 | 44.0-63.0 | 38.0-58.0 |
| Hind Foot length | 7.0-9.5 | 7.0-8.0 | 11.3-12.9 | 12.0-15.0 | 8.0-15.0 | 6.0-10.0 | 9.0-10.0 | 9.0-14.0 | 7.0-10.0 |
| Ear length | 37.0-42.0 | 15.0-17.0 | 30.0-42.0 | 19.0-23.0 | 9.0-20.2 | Moderate and tip broadly rounded | 12.0-15.0 | 14.0-18.0 | 12.5-15.0 Large and thick, with ridges |
| Cranial characters | | | | | | | | | |
| Condylar canine length | 13.8-15.4 | 13.4-14.2 | ~21.7 | 20.1-20.9 | 16.3-21.3 | 13.8-14.8 | 14.6-15.6 | 17.2-19.6 | 11.7-19.5 |
| Maxillary toothrow (CM ³) | 5.0-5.8 | 4.7-5.1 | ~8.5 | 7.8-8.5 | 6.1-8.4 | 5.5-5.9 | 6.0-6.5 | 7.1-8.2 | 4.4-7.8 |
| Incisors – Upper – Lower | 2 pairs 3 pairs | 2 pairs 3 pairs | 1 pair 3 pairs | 1 pair 3 pairs | 1 pair 3 pairs | 1 pair 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs |
| Premolars – Upper – Lower | 2 pairs 3 pairs | 2 pairs 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs | 2 pairs 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs |
| Dental formula | 2123 3133 = 36 | 2123 3123 = 34 | 1113 3123 = 30 | 1113 3123 = 30 | 1113 3123 = 30 | 1113 3123 = 30 | 2123 3123 = 34 | 2113 3123 = 32 | 2112 3123 = 32 |
| Species content | | | | | | | | | |
| | <i>P. homochrous</i> Hodgson, 1847 <i>P. wardi</i> Thomas, 1911 <i>P. strelkovi</i> Spitzenberger, 2008 | <i>B. leucomelas</i> (Cretzschmar, 1826) | <i>O. hemprichii</i> Peters, 1859 | <i>S. ornatus</i> (Blyth, 1851) | <i>S. heathi</i> Horsfield, 1831; <i>S. kuhlii</i> Leach, 1821 | <i>S. pallidus</i> Dobson, 1876 | <i>A. circumdatus</i> (Temminck, 1840) | <i>H. tickelli</i> (Blyth, 1851) | <i>E. serotinus</i> (Schreber, 1774); <i>E. bottae</i> (Peters, 1869); <i>E. pachyotis</i> (Dobson, 1871); <i>E. dimissus</i> Thomas, 1916 <i>E. gobiensis</i> Bobrinskii, 1926; <i>E. nasutus</i> (Dobson, 1877); <i>E. tatei</i> Ellerman & Morrison- Scott, 1951 |
| Further table reference | 7.4 | 7.4 | 7.4 | 7.5 | 7.5 | 7.5 | 7.6 | 7.6 | 7.6 |

Note: All measurements are in mm

Contd...

Table 7 (Contd.). Diagnostic morphological characters of the genera of the tribes Pipistrellini and Vespertilionini of Subfamily Vespertilioninae of family Vespertilionidae present in South Asia

| Genus | <i>Scotozous</i> Dobson, 1875 | <i>Nyctalus</i> Bowdich, 1825 | <i>Pipistrellus</i> Kaup, 1829 | <i>Tylonycteris</i> Peters, 1872 | <i>Ia</i> Thomas, 1902 | <i>Falsistrellus</i> Troughton, 1943 | <i>Philetor</i> Thomas, 1902 | <i>Vespertilio</i> Linnaeus, 1758 | <i>Hypsugo</i> Kolenati, 1856 |
|---------------------------------------|----------------------------------|----------------------------------|--|---|---------------------------------------|--|---|--|--|
| External characters | | | | | | | | | |
| Forearm length | 32.7-36.3 | 42.1-57.8 | 25.0-42.0 | 26.1-29.0 | 70.9-77.3 | 38.4-41.4 | 31.7-35.7 | 42.0-45.5 | 32.1-38.0 |
| Head Body length | 39.0-55.0 | 68.0-80.0 | 33.0-64.0 | 34.0-46.0 | NA | 43.0-51.0 | NA | 55.0-66.0 | 47.0-60.0 |
| Tail length | 27.0-41.0 | 31.0-55.0 | 20.0-45.0 | 26.0-33.0 | ~65.0 | 30.0-41.0 | 27.1-32.2 | 40.0-48.0 | 30.0-49.0 |
| Hind Foot length | 5.0-8.0 | 6.0-11.4 | 3.0-11.0 | 5.0-7.0 | ~17.0 | 7.0-8.0 | 6.4-8.1 | 8.0-10.0 | 6.4-8.0 |
| Ear length | 10.0-18.0 | 7.0-17.0 Short | 5.0-14.0 | 8.5-10.5 Triangular | 23.7-24.0 Broad and tip rounded | 12.0-15.0 | 8.0-10.0 Short, broad & tip rounded | 14.7-16.0 Short and tip broadly rounded | 10.0-15.0 |
| Cranial characters | | | | | | | | | |
| Condylar canine length | 12.8-13.6 | 14.7-18.6 | 9.3-14.3 | 10.0-11.7 | 25.2-26.2 | 13.7-14.7 | 12.4-13.7 | ~15.0 | 11.4-13.3 |
| Maxillary toothrow (CM ³) | 5.2-5.6 | 5.5-7.7 | 3.5-5.9 | 3.4-4.2 | 10.5-11.0 | 5.5-5.7 | 4.5-4.8 | 4.9-5.5 | 4.6-5.1 |
| Incisors – Upper – Lower | 1 pair* 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs | 2 pairs 3 pairs |
| Premolars – Upper – Lower | 2 pairs 2 pairs | 2 pairs 2 pairs | 1 pair 2 pairs | 2 pairs 2 pairs | 2 pairs 2 pairs | 2 pairs 2 pairs | 1 pair 2 pairs | 1 pair 2 pairs | 2 pairs 2 pairs |
| Dental formula | 1123 3123 =32 | 2123 3123 =32 | 2123 3123 =32 | 2113 3123 =32 | 2123 3123 =32 | 2123 3123 =32 | 2112 3123 =32 | 212(1)3 3123 =32 | 212(1)3 3123 =34/32 |
| Species content | | | | | | | | | |
| S. dormieri Dobson, 1875 | | | <i>P. pipistrellus</i> (Schreber, 1774); <i>P. paterculus</i> Thomas, 1915; <i>P. javanicus</i> (Gray, 1838); <i>N. noctula</i> (Schreber, 1774); <i>N. leisleri</i> (Kuhl, 1817); <i>N. montanus</i> (Barrett-Hamilton, 1906) | <i>T. pachypus</i> (Temminck, 1840); <i>T. robustula</i> Thomas, 1915 | <i>I. io</i> Thomas, 1902 | <i>F. affinis</i> (Dobson, 1871) | <i>P. brachypterus</i> (Temminck, 1840) | <i>V. murinus</i> Linnaeus, 1758 | <i>H. savii</i> (Bonaparte, 1837) <i>H. cadornae</i> (Thomas, 1916) |
| Further table reference | 7.7 | 7.7 | 7.7 | 7.8 | 7.8 | 7.8 | 7.8 | 7.8 | 7.8 |

Note: All measurements are in mm; * - outer upper incisor, when present, no more than a minute pointed spicule; † Anterior upper premolar (pm^2) much reduced, rarely absent

Table 7.1. Diagnostic morphological characters of the species belonging to the genera *Harpiocephalus*, *Harpiola* and *Murina* present in South Asia

| Species | <i>Harpiocephalus harpia</i> (Temminck, 1840) | <i>Harpiola grisea</i> Peters, 1872 | <i>Murina leucogaster</i> Milne-Edwards, 1872 |
|--|--|--|--|
| External characters | | | |
| Forearm length | 44.1-50.1 | 32.8-34.2 | ~ 40.9 |
| Head Body length | 60.0-75.0 | ~ 35.6 | ~ 47.0 |
| Tail length | 40.0-50.0 | NA | ~ 35.0 |
| Hind Foot length | 11.0-14.0 | NA | ~ 9.0 |
| Ear length – Posterior border – Emargination on posterior border | 17.0-18.0 Evenly rounded | NA NA Conspicuous | ~ 15.0 Slightly convex Distinct |
| Interfemoral membrane | Covered with long reddish hairs | Densely covered with rather long hairs | Moderately covered with long fine reddish hair above and naked below |
| Cranial characters | | | |
| Condylar canine length | 19.0-19.5 | ~ 12.1 | ~ 16.3 |
| Maxillary toothrow (CM ³) | 6.5-6.9 | ~ 4.9 | ~ 6.1 |
| Mandibular toothrow (CM ₃) | 7.5-8.0 | ~ 5.8 | ~ 6.6 |
| Zygomatic breadth | 13.6-13.7 | NA | ~ 10.0 |
| Mandible length | 15.1-16.0 | ~ 4.1 | ~ 13.1 |

Note: All measurements are in mm; NA – Data not available

Contd...

Table 7.1 (Contd..). Diagnostic morphological characters of the species belonging to the genus *Murina* present in South Asia

| Species | <i>Murina aurata</i> Milne-Edwards, 1872 | <i>Murina cyclotis</i> Dobson, 1872 | <i>Murina tubinaris</i> (Scully, 1881) | <i>Murina huttoni</i> (Peters, 1872) |
|--|---|---|--|--|
| External characters | | | | |
| Forearm length | 27.7-29.6 | 29.7-34.5 | 30.1-34.1 | 32.8-35.4 |
| Head Body length | ~ 45.0 | 38.0-50.0 | 39.0-48.0 | ~ 48.0 |
| Tail length | ~ 27.0 | 32.0-41.0 | 22.0-35.0 | ~ 37.0 |
| Hind Foot length | ~ 4.5 | 7.0-10.0 | 6.0-9.0 | ~ 6.0 |
| Ear length – Posterior border – Emargination on posterior border | ~ 13.0 Slightly convex Lacking | 13.0-16.0 Slightly convex Lacking | 13.0-15.0 Smoothly convex Small | ~ 17.0 Smoothly convex Lacking |
| Interfemoral membrane | Moderately covered with golden hair above & gray hair below | Moderately covered with orange hairs above & nearly naked below | Moderately covered with pale gray or ferruginous hairs above & naked below | Moderately covered with pale brown hairs above & naked below |
| Cranial characters | | | | |
| Condylar canine length | ~ 12.3 | 13.9-15.0 | 13.4-14.2 | 14.9-15.5 |
| Maxillary toothrow (CM ³) | ~ 4.5 | 5.2-5.7 | 5.0-5.4 | 5.8-6.1 |
| Mandibular toothrow (CM ₃) | 4.5-4.9 | 5.6-6.2 | 5.2-5.8 | 6.4-6.9 |
| Zygomatic breadth | ~ 7.5 | 9.2-9.7 | 8.4-9.0 | 9.4-9.8 |
| Mandible length | 9.2-9.4 | 10.8-11.8 | 10.3-10.9 | 11.5-12.2 |

Note: All measurements are in mm; NA – Data not available

Table 7.2. Diagnostic morphological characters of the species of the genus *Kerivoula* present in South Asia

| Species | <i>Kerivoula picta</i> (Pallas, 1767) | <i>Kerivoula hardwickii</i> (Horsfield, 1824) | <i>Kerivoula lenis</i> Thomas, 1916 |
|--|--|--|--|
| External characters | | | |
| Forearm length | 31.5-37.9 | 31.7-36.0 | 37.2-41.0 |
| Head Body length | 45.0-48.0 | 39.0-55.0 | NA |
| Tail length | 43.0-48.0 | 35.0-43.0 | ~48.0 |
| Hind Foot length | 4.0-8.0 | 5.0-8.0 | NA |
| Ear length | 14.0-16.0 Relatively large, tip rounded | 11.0-15.0 Moderate-sized, tip rounded | ~13.8 Moderate-sized, tip rounded |
| Tragus | Tall and narrow | Narrow, sharply pointed | NA |
| Interfemoral membrane | Orange to scarlet hairy above along the body parts | Brown & nearly transparent, sparsely haired above near body parts & posterior border | Without hairs, veins sometimes whitish |
| Cranial characters | | | |
| Condylar canine length | 12.2-13.3 | 12.4-13.0 | 14.5-15.1 |
| Maxillary toothrow (CM ³) | 5.5-5.8 | 5.5-5.8 | 6.6-6.8 |
| Mandibular toothrow (CM ₃) | 5.9-6.1 | 5.9-6.1 | 7.2-7.6 |
| Zygomatic breadth | 8.4-8.8 | 8.6-8.7 | 9.6-10.6 |
| Mandible length | 10.0-10.7 | 9.9-10.6 | 11.8-12.4 |

Note: All measurements are in mm; NA – Data not available

Table 7.3. Diagnostic morphological characters of the species belonging to the genus *Myotis* Kaup, 1829 present in South Asia

| Species | <i>Myotis blythii</i> (Tomes, 1857) | <i>Myotis sicarius</i> Thomas, 1915 | <i>Myotis formosus</i> (Hodgson, 1835) | <i>Myotis nipalensis</i> (Dobson, 1871) | <i>Myotis muricola</i> (Gray, 1846) | <i>Myotis silvorensis</i> (Horsfield, 1855) |
|--|--|--|---|--|--|--|
| External characters | | | | | | |
| Forearm length | 55.5-58.3 | 48.1-54.6 | 44.5-49.1 | 34.5-36.8 | 31.2-37.0 | 30.0-31.5 |
| Head Body length | 65.0-80.0 | 50.0-56.0 | NA | 38.0-47.0 | 41.0-47.0 | NA |
| Tail length | 53.0-68.0 | 44.0-47.0 | 36.0-48.0 | 32.0-40.0 | 25.0-39.0 | NA |
| Hind Foot length | 11.0-17.0 | 10.0-11.0 | 10.2-11.6 | 7.0-8.0 | Small | 4.0-6.7 |
| Ear length | 19.0-26.0 | 17.0-19.0 Medium, tips bluntly rounded | 12.8-14.5 Ovate, orange coloured | 12.0-14.0 Relatively small, dark brown/black | 6.0-13.3 Relatively small, dark brow/black | NA |
| Tragus | Tall and narrow | Tall and narrow | Tall, narrow and obtusely pointed | Tall and narrow | Tall and narrow | Tall and narrow |
| Interfemoral membrane | Pale brown, hairless throughout | Semi-translucent, dark brown/black hairless throughout | Orange | Dark brownish with some hairs on dorsal side near the body | Dark brownish with some hairs on dorsal side near the body | Brown |
| Cranial characters | | | | | | |
| Condylar canine length | 18.1-18.9 | 16.9-17.7 | 16.3-16.6 | 11.6-12.4 | 11.5-12.3 | ~11.2 |
| Maxillary toothrow (CM ³) | 8.2-8.5 | 7.1-7.5 | 6.8-7.2 | 4.7-5.7 | 4.8-5.3 | 4.5-4.8 |
| Mandibular toothrow (CM ₃) | ~9.2 | 4.9-5.3 | 7.4-7.8 | 5.3-5.7 | 5.1-5.6 | 4.9-5.0 |
| Zygomatic breadth | NA | ~11.8 | 11.7-11.9 | NA | 7.3-8.5 | NA |
| Mandible length | 15.7-16.0 | 14.1-14.9 | 13.7-14.1 | 9.7-10.3 | 9.2-10.1 | 9.0-9.2 |

Contd...

Note: All measurements are in mm; NA – Data not available

Table 7.3 (Contd..) Diagnostic morphological characters of the species belonging to the genus *Myotis* Kaup, 1829 present in South Asia

| Species | <i>Myotis montivagus</i> (Dobson, 1874) | <i>Myotis annectans</i> (Dobson, 1871) | <i>Myotis longipes</i> (Dobson, 1873) | <i>Myotis corsbai</i> Topal, 1997 | <i>Myotis laniger</i> (Peters, 1871) | <i>Myotis horsfieldii</i> (Temminck, 1840) | <i>Myotis hasseltii</i> (Temminck, 1840) |
|--|--|---|--|--------------------------------------|---|---|--|
| External characters | | | | | | | |
| Forearm length | 44.7-46.8 | 45.3-46.5 | 36.5-39.0 | 34.8-37.5 | ~34.8 | 36.5-41.5 | 37.2-40.3 |
| Head Body length | 56.0-62.0 | 45.0-48.0 | 43.0-46.0 | NA | ~41.3 | 49.0-59.0 | 52.0-58.0 |
| Tail length | 42.0-48.0 | 39.0-45.0 | 37.0-42.0 | NA | ~38.6 | 34.0-42.0 | 35.0-42.0 |
| Hind Foot length | 9.0-10.0 Medium | ~10.0 Medium | 9.2-9.7 Large | 14.8-15.8* | ~7.9 Large | 7.0-11.2 Large | 9.0-11.0 Very large |
| Ear length | 14.2-15.5 Relatively short | 14.0-15.0 Small, tips bluntly rounded | 10.6-15.0 Narrow | NA | Relatively small, rounded | 13.0-15.2 Small with rounded tip | 15.5-16.5 Rather narrow, tips narrowly rounded |
| Tragus | Short | Relatively broad | Tall and narrow | NA | Short | Short | Short and narrow |
| Interfemoral membrane | Dark brown, hairyless throughout | Dark brown/black hairyless throughout | Brown with some hairs near the hindlimbs | NA | Brownish | Deep chocolate brown, some hairs on dorsal side near the body | Dark brown |
| Cranial characters | | | | | | | |
| Condylar canine length | 15.1-15.6 | 14.9-15.5 | 12.2-12.8 | 12.11-12.71† | 10.6-11.9† | 13.4-14.3 | 14.0-14.6 |
| Maxillary toothrow (CM ³) | 6.5-6.8 | 6.5-6.7 | 5.1-5.3 | 4.7-5.19 | 4.8-5.0 | 5.5-5.9 | 5.8-6.0 |
| Mandibular toothrow (CM ₃) | 6.9-7.2 | 7.1-7.2 | 5.5-5.9 | NA | 4.9-5.5 | 5.9-6.4 | 6.2-6.6 |
| Zygomatic breadth | NA | ~11.5 | ~8.4 | NA | ~8.9 | 9.3-10.2 | ~10.5 |
| Mandible length | 12.9-13.3 | 12.8-13.2 | 10.1-10.5 | NA | NA | 11.2-11.6 | 11.7-12.0 |

Note: All measurements are in mm; NA - Data not available; * - Tibia length; † - Condylarbasal length (CBL)

Table 7.4. Diagnostic morphological characters of the species belonging to the genera *Plecotus*, *Barbastella* and *Otonycteris* present in South Asia

| Species | <i>Plecotus homochrous</i> Hodgson, 1847 | <i>Plecotus wardi</i> Thomas, 1911 | <i>Plecotus strelkovi</i> Spitzenberger, 2008 | <i>Barbastella leucomelas</i> (Cretzschmar, 1826) | <i>Otonycteris hemprichii</i> Peters, 1859 |
|--|--|---|--|---|---|
| External characters | | | | | |
| Forearm length | 36.5-40.3 | 41.9-45.1 | 39.6-44.9 | 38.7-42.1 | 64.7-65.6 |
| Head Body length | 40.0-45.0 | 47.0-53.0 | - | 47.0-51.0 | 61.0-76.0 |
| Tail length | 48.0-50.0 | 49.0-54.0 | ~ 51.0 | 40.0-47.0 | 40.0-58.0 |
| Hind Foot length | 7.0-8.0 | 7.0-9.5 | - | 7.0-8.0 | 11.3-12.9 |
| Ear length | 39.0-41.0 Long and oval; joined over forehead | 37.0-42.0 Large and oval; joined over forehead | ~ 40.0 Large and oval; joined over forehead | 15.0-17.0 Small squarish; joined over forehead | 30.0-42.0 Large and tip broadly rounded |
| Tragus | Tall, ½ the pinna, tip bluntly pointed | Tall, ½ the pinna, tip bluntly pointed | Tall, ½ the pinna, tip bluntly pointed | Triangular | Tall, ½ the pinna |
| Antitragus | Indefinable | Indefinable | Indefinable | Absent | Shallow notch-like |
| Cranial characters | | | | | |
| Condylar canine length | 13.8-14.2 | 14.8-15.4 | 14.1-15.6 | 13.4-14.2 | ~ 21.7 |
| Maxillary toothrow (CM ³) | 5.0-5.3 | 5.4-5.8 | 5.4-6.1 | 4.7-5.1 | ~ 8.5 |
| Mandibular toothrow (CM ₃) | 5.4-5.7 | 5.9-6.3 | 5.8-6.4 | 5.1-5.7 | 9.7-9.8 |
| Zygomatic breadth | 8.1-8.2 | 8.5-8.9 | 8.4-9.2 | 7.3-7.6 | 13.6-15.8 |
| Mandible length | 9.8-10.2 | 10.6-11.2 | 10.4-11.4 | 9.6-10.1 | 17.1-17.4 |

Note: All measurements are in mm

Table 7.5. Diagnostic morphological characters of the species belonging to the genera *Scotoecus*, *Scotomanes* and *Scotophilus* present in South Asia

| Species | <i>Scotoecus pallidus</i> Dobson, 1876 | <i>Scotomanes ornatus</i> (Blyth, 1851) | <i>Scotophilus heathi</i> Horsfield, 1831 | <i>Scotophilus kuhlii</i> Leach, 1821 |
|--|---|--|--|--|
| External characters | | | | |
| Forearm length | 34.1-37.3 | 56.1-61.2 | 55.4-65.8 | 44.0-56.0 |
| Head Body length | 50.0-58.0 | 64.0-85.0 | 67.0-93.0 | 60.0-78.0 |
| Tail length | 34.0-41.0 | 52.0-66.0 | 43.0-71.0 | 40.0-65.0 |
| Hind Foot length | 6.0-10.0 | 12.0-15.0 | 9.0-15.0 | 8.0-13.0 |
| Ear length | 12.0-15.0 Moderate with rounded tip | 19.0-23.0 Large and tip broadly rounded | 13.0-20.2 Small with ridges | 9.0-17.0 Small with ridges |
| Tragus | Long and narrow | Tall, up to ½ the pinna, broad & crescent-shaped | Tall, ½ the pinna, crescent-shaped | Tall, ½ the pinna, crescent-shaped |
| Antitragus | Indefinable | Indefinable | Well formed | Well formed |
| Cranial characters | | | | |
| Condylar canine length | 13.8-14.8 | 20.1-20.9 | 19.0-21.3 | 16.3-18.0 |
| Maxillary toothrow (CM ³) | 5.5-5.9 | 7.8-8.5 | 7.1-8.4 | 6.1-6.8 |
| Mandibular toothrow (CM ₃) | 5.9-6.4 | 8.7-9.4 | 8.1-9.6 | 6.9-7.8 |
| Zygomatic breadth | ~ 10.5 | 15.9-17.3 | 14.5-16.9 | 12.4-13.7 |
| Mandible length | 10.9-12.0 | 16.9-17.9 | 14.8-18.0 | 12.9-14.4 |

Note: All measurements are in mm

Table 7.6. Diagnostic morphological characters of the species belonging to the genus *Arielulus*, *Hesperoptenus* and *Eptesicus* present in South Asia

| Species | <i>Arielulus circumdatus</i> (Temminck, 1840) | <i>Hesperoptenus tickelli</i> (Blyth, 1851) | <i>Eptesicus serotinus</i> (Schreber, 1774) | <i>Eptesicus bottae</i> (Peters, 1869) | <i>Eptesicus pachyotis</i> (Dobson, 1871) |
|--|--|--|---|---|--|
| External characters | | | | | |
| Forearm length | 41.8-43.6 | 50.0-60.4 | 54.2-55.1 | ~ 42.1 | 38.0-45.3 |
| Head Body length | ~ 95.0 | 61.0-79.0 | ~ 80.0 | ~ 57.0 | 55.0-56.0 |
| Tail length | ~ 40.0 | 44.0-63.0 | ~ 58.0 | ~ 45.0 | 40.0-41.0 |
| Hind Foot length | ~ 10.0 | 9.0-14.0 | ~ 10.0 | ~ 7.0 | 8.0-9.0 |
| Ear length | ~ 15.0 Dark brown to black, with pale margins in some specimens | 14.0-18.0 Moderately large, thick & fleshy | ~ 14.0 Moderately tall, dark | ~ 15.0 Moderately tall, dark | 13.0-14.0 Triangular with rounded tip |
| Tragus | Broad with pale margin | Tall, ½ the pinna, crescent-shaped | Small, less than ½ the pinna, tip bluntly pointed | NA | Short, broadly rounded & curved inwards |
| Cranial characters | | | | | |
| Condylar length | 14.6-15.6 | 17.2-19.6 | 18.3-19.5 | ~ 15.0 | ~ 21.2* |
| Maxillary toothrow (CM ³) | 6.0-6.5 | 7.1-8.2 | 7.1-7.8 | ~ 5.8 | NA |
| Mandibular toothrow (CM ₃) | 6.4-6.8 | 7.9-9.2 | 8.4-8.7 | ~ 6.8 | NA |
| Zygomatic breadth | ~ 11.7 | 13.1-15.4 | 13.3-14.6 | NA | NA |
| Mandible length | 11.8-12.7 | 14.0-16.5 | 15.0-16.0 | ~ 6.2 | NA |

Note: All measurements are in mm; NA – Data not available; * - Greatest skull length

Table 7.6 (Contd..). Diagnostic morphological characters of the species belonging to the genus *Eptesicus* present in South Asia

| Species | <i>Eptesicus gobiensis</i> Bobrinskii, 1926 | <i>Eptesicus nasutus</i> Dobson, 1877 | <i>Eptesicus dimissus</i> Thomas, 1916 | <i>Eptesicus tatei</i> Ellerman & Morrison-Scott, 1951 |
|--|--|--|---|---|
| External characters | | | | |
| Forearm length | 41.0-41.5 | 35.4-36.9 | 38.0-42.0 | ~ 43.4 |
| Head Body length | NA | 40.0-46.0 | ~ 56.0 | ~ 48.5 |
| Tail length | NA | 38.0-46.0 | 36.0-41.0 | ~ 45.9 |
| Hind Foot length | NA | 7.0-8.0 | ~ 8.0 | NA |
| Ear length | 14.0-15.0 Moderately tall | 12.5-14.0 Small, narrowly rounded tip | 14.0-15.0 Small and rounded | ~ 15.3 Moderately tall, oval, with rounded tip |
| Tragus | NA | Tall, ½ the pinna | NA | Tall, ½ the pinna |
| Cranial characters | | | | |
| Condylar length | ~ 15.1† | 11.7-12.2 | ~ 15.4† | NA |
| Maxillary toothrow (CM ³) | ~ 5.9 | 4.4-4.8 | ~ 5.9 | NA |
| Mandibular toothrow (CM ₃) | NA | 4.8-5.1 | NA | NA |
| Zygomatic breadth | NA | 8.4-8.8 | NA | NA |
| Mandible length | NA | 8.6-9.6 | NA | NA |

Note: All measurements are in mm; NA – Data not available; * - Greatest skull length; † - Condylar length

Table 7.7. Diagnostic morphological characters of the species belonging to the genera *Nyctalus* and *Scotozous* present in South Asia

| Species | <i>Nyctalus noctula</i> (Schreber, 1774) | <i>Nyctalus leisleri</i> (Kuhl, 1817) | <i>Nyctalus montanus</i> (Barrett-Hamilton, 1906) | <i>Scotozous dormeri</i> (Dobson, 1875) |
|--|---|--|--|--|
| External characters | | | | |
| Forearm length | 50.9-57.8 | 42.1-45.2 | 42.9-43.0 | 32.7-36.3 |
| Head Body length | 68.0-80.0 | 62.0-72.0 | ~ 70.0 | 39.0-55.0 |
| Tail length | 33.0-55.0 | 31.0-45.0 | ~ 43.0 | 27.0-41.0 |
| Hind Foot length | 10.2-11.4 | 6.0-10.0 | NA | 5.0-8.0 |
| Ear length | 13.0-17.0 Moderate | 7.0-16.0 Moderate | ~ 14.0 Short | 10.0-18.0 |
| Tragus | Club-shaped expanded distally | Club-shaped expanded distally | Club-shaped expanded distally | Without hairs, veins sometimes whitish |
| Cranial characters | | | | |
| Condylar canine length | 17.1-18.6 | 14.7-15.2 | 15.4-16.0 | 12.8-13.6 |
| Maxillary toothrow (CM ³) | 7.1-7.7 | 5.5-5.9 | 6.5-6.6 | 5.2-5.6 |
| Mandibular toothrow (CM ₃) | 7.5-8.2 | 5.9-6.3 | 6.9-7.0 | 5.5-6.1 |
| Zygomatic breadth | 12.2-13.3 | 9.8-10.7 | NA | 9.6-10.5 |
| Mandible length | 13.8-14.7 | 11.4-11.7 | 12.7-12.8 | 10.4-11.2 |

Note: All measurements are in mm; NA – Data not available

Table 7.7 (Contd..). Diagnostic morphological characters of the species belonging to the genus *Pipistrellus* present in South Asia

| Species | <i>Pipistrellus pipistrellus</i> (Schreber, 1774) | <i>Pipistrellus paterculus</i> Thomas, 1915 | <i>Pipistrellus javanicus</i> (Gray, 1838) | <i>Pipistrellus coromandra</i> (Gray, 1838) |
|---|--|--|---|---|
| External characters | | | | |
| Forearm length | 30.0-30.6 | 29.2-34.0 | 30.0-36.0 | 25.5-34.3 |
| Head Body length | 40.0-48.0 | 42.0-48.0 | 40.0-55.0 | 34.0-49.0 |
| Tail length | 29.0-35.0 | 31.0-38.0 | 26.0-40.0 | 22.0-39.0 |
| Hind Foot length | 6.0-7.0 | 6.0-7.0 | 3.0-8.0 | 3.4-8.0 |
| Ear length | 10.5-12.0 | 10.0-13.0 | 5.0-15.0 | 7.1-14.0 |
| Interfemoral membrane | Sparsely haired above near the body parts | Sparsely haired above near the body | Without hairs | Sparsely haired above and below near the body parts |
| Cranial characters | | | | |
| Condylar canine length | 10.4-11.3 | 10.6-11.6 | 11.9-13.1 | 10.6-11.9 |
| Maxillary toothrow (CM ³) | 4.1-4.4 | 4.1-4.8 | 4.6-5.2 | 3.9-4.6 |
| Mandibular toothrow (CM ₃) | 4.3-4.7 | 4.4-5.0 | 4.8-5.5 | 4.1-5.1 |
| Zygomatic breadth | 7.2-7.9 | NA | 8.2-9.0 | 7.6-8.2 |
| Mandible length | 7.9-8.7 | 8.4-9.1 | 9.3-10.7 | 8.2-9.5 |
| Posterior palatal width | 4.8-5.2 | 5.3-5.9 | 5.6-6.7 | 5.0-6.0 |
| I ³ ½ or > than I ² | Yes | Yes | Yes | Yes |
| Upper canine | Bicuspidate | Unicuspidate | Bicuspidate | Bicuspidate |

Note: All measurements are in mm; NA – Data not available

Contd...

Table 7.7 (Contd.). Diagnostic morphological characters of the species belonging to the genus *Pipistrellus* present in South Asia

| Species | <i>Pipistrellus tenuis</i> (Temminck, 1840) | <i>Pipistrellus ceylonicus</i> (Kelaart, 1852) | <i>Pipistrellus kuhlii</i> (Kuhl, 1817) | <i>Pipistrellus abramus</i> (Temminck, 1840) |
|---|--|---|--|---|
| External characters | | | | |
| Forearm length | 25.0-30.2 | 33.0-42.0 | 33.4-36.0 | 31.4-34.4 |
| Head Body length | 33.0-45.0 | 45.0-64.0 | 35.0-49.0 | NA |
| Tail length | 20.0-35.0 | 30.0-45.0 | 33.0-45.0 | NA |
| Hind Foot length | 3.0-7.0 | 6.0-11.0 | 6.0-8.0 | 6.4-7.6 |
| Ear length | 5.0-11.0 | 9.5-14.0 | 10.0-13.0 | NA |
| Interfemoral membrane | Without hairs | Sparsely haired above and below near the body parts | Pallid & Translucent | NA |
| Cranial characters | | | | |
| Condylar canine length | 9.3-10.7 | 13.1-14.3 | 12.0-12.9 | 11.4-12.8* |
| Maxillary toothrow (CM ³) | 3.5-4.1 | 5.2-5.9 | 4.6-5.0 | 4.2-4.9 |
| Mandibular toothrow (CM ₃) | 3.8-4.4 | 5.7-6.5 | 5.0-5.5 | 4.6-5.3 |
| Zygomatic breadth | 7.3-7.6 | 9.2-11.0 | 8.4-8.7 | ~ 8.2 |
| Mandible length | 7.2-8.3 | 10.6-12.0 | 9.3-10.4 | 8.9-10.0 |
| Posterior palatal width | 4.5-5.2 | 6.2-7.2 | 5.5-5.9 | 5.0-6.0 |
| I ³ ½ or > than I ² | Yes | Yes | No | Yes |
| Upper canine | Bicuspidate | Bicuspidate | Faintly bicuspidate | Unicuspidate |

Note: All measurements are in mm; NA – Data not available; * - Condylabasal length

Table 7.8. Diagnostic morphological characters of species of the genera *Tylonycteris*, *Ia*, *Falsistrellus* and *Vespertilio* present in South Asia

| Species | <i>Tylonycteris pachypus</i> (Temminck, 1840) | <i>Tylonycteris robustula</i> Thomas, 1915 | <i>Ia io</i> Thomas, 1902 | <i>Falsistrellus affinis</i> (Dobson, 1871) | <i>Vespertilio murinus</i> Linnaeus, 1758 |
|--|--|--|--|--|--|
| External characters | | | | | |
| Forearm length | 26.1-29.0 | 26.6-28.1 | 70.9-77.3 | 38.4-41.4 | 42.0-45.5 |
| Head Body length | 34.0-46.0 | 40.0-44.0 | NA | 43.0-51.0 | 55.0-66.0 |
| Tail length | 26.0-33.0 | 26.0-31.0 | ~ 65.0 | 30.0-41.0 | 40.0-48.0 |
| Hind Foot length | 5.0-7.0 | 5.0-5.5 | ~ 17.0 | 7.0-8.0 | 8.0-10.0 |
| Ear length | 9.0-10.0 Small, triangular with broadly rounded tip | 8.5-10.5 Small, triangular with broadly rounded tip | 23.7-24.0 Large, broad with rounded tip | 12.0-15.0 | 14.7-16.0 Short and tip broadly rounded |
| Tragus | Short and broad | Short and broad | Small, 1/3 rd the pinna | | Small, short bluntly rounded |
| Cranial characters | | | | | |
| Condylar canine length | 10.0-11.3 | 11.1-11.7 | 25.2-26.2 | 13.7-14.7 | ~ 15.0 |
| Maxillary toothrow (CM ³) | 3.4-4.2 | 3.9-4.1 | 10.5-11.0 | 5.5-5.7 | 4.9-5.5 |
| Mandibular toothrow (CM ₃) | 3.6-4.6 | 5.4-5.8 | 11.6-12.1 | 5.6-5.8 | 5.3-6.1 |
| Zygomatic breadth | 8.4-8.5 | 8.9-9.1 | 16.7-18.0 | ~ 9.1 | 9.1-9.7 |
| Mandible length | 7.6-8.9 | 8.5-9.0 | 20.8-21.8 | 10.5-11.4 | 10.1-11.4 |

Note: All measurements are in mm; NA – Data not available

Contd...

Table 7.8 (Contd.). Diagnostic morphological characters of species of the genera *Philetor* and *Hypsugo* present in South Asia

| Species | <i>Philetor brachypterus</i> (Temminck, 1840) | <i>Hypsugo savii</i> (Bonaparte, 1837) | <i>Hypsugo cadornae</i> (Thomas, 1916) |
|--|---|--|---|
| External characters | | | |
| Forearm length | 31.7-35.7 | 32.1-38.0 | 32.6-36.5 |
| Head Body length | NA | 47.0-60.0 | 47.0-52.5 |
| Tail length | 27.1-32.2 | 30.0-35.0 | 34.0-49.0 |
| Hind Foot length | 6.4-8.1 | 6.4-8.0 | 6.5-7.0 |
| Ear length | 8.0-10.0 Short and broad, broadly rounded tip | 10.0-14.0 | 14.0-15.0 |
| Interfemoral membrane | Broad & short, fleshy & thick | Sparingly haired near the body and tail | Without hairs |
| Cranial characters | | | |
| Condylar canine length | 12.4-13.7 | 11.4-13.3 | 12.6-12.8 |
| Maxillary toothrow (CM ³) | 4.5-4.8 | 4.6-5.1 | 4.6-4.9 |
| Mandibular toothrow (CM ₃) | 4.7-5.0 | 4.9-5.2 | 4.8-5.1 |
| Zygomatic breadth | 10.0-10.7 | 8.5-9.1 | NA |
| Mandible length | 9.9-10.7 | 9.6-10.3 | 9.5-10.3 |

Note: All measurements are in mm; NA – Data not available

Table 8. Diagnostic morphological characters of one genus of the family Miniopteridae present in South Asia

| Genus | <i>Miniopterus</i> Bonaparte, 1837 |
|---------------------------------------|---|
| External characters | |
| Forearm length | 39.6-49.6 |
| Head Body length | 40.1-65.0 |
| Tail length | 39.6-61.0 |
| Hind Foot length | ~ 7.0-12.0 |
| Ear length | ~ 8.7-12.0 Small and tip broadly rounded |
| Cranial characters | |
| Condylar canine length | 12.0-14.8 |
| Maxillary toothrow (CM ³) | 5.1-6.3 |
| Incisors – Upper – Lower | 2 pairs 3 pairs |
| Premolars – Upper – Lower | 2 pairs 3 pairs |
| Dental formula | -23,1,-2-4,123 123,1,-234,123 = 36 |
| Species content | <i>M. schreibersii</i> (Kuhl, 1817); <i>M. pusillus</i> Dobson, 1876 <i>M. magnater</i> Sanborn, 1931 |
| Further table reference | 8.1 |

Note: All measurements are in mm

Table 8.1. Diagnostic morphological characters of species of the genera *Miniopterus* Bonaparte, 1837 present in South Asia

| Species | <i>Miniopterus schreibersii</i> (Kuhl, 1817) | <i>Miniopterus pusillus</i> (Dobson, 1876) | <i>Miniopterus magnater</i> Sanborn, 1931 |
|--|---|---|--|
| External characters | | | |
| Forearm length | 44.7-49.6 | 39.6-40.2 | 47.0-53.0 |
| Head Body length | 47.0-65.0 | NA | NA |
| Tail length | 44.0-61.0 | 40.1-44.1 | NA |
| Hind Foot length | 7.0-12.0 | NA | NA |
| Ear length | 8.7-12.0 Small, tip broadly rounded | NA | NA |
| Tragus | ½ the pinna, slightly curved forward | NA | NA |
| Interfemoral membrane | Dark brownish-black, sparingly haired near the body parts | Dark brownish-black, fur on membrane extends further away from body parts | NA |
| Cranial characters | | | |
| Condylarcanine length | 13.6-14.8 | 12.0-12.7 | 16.8-17.3 [†] |
| Maxillary toothrow (CM ³) | 5.8-6.3 | 5.1-5.3 | 7.0-7.2 |
| Mandibular toothrow (CM ₃) | 6.3-6.8 | 5.4-5.6 | 8.3-8.9 |
| Zygomatic breadth | 8.5-9.1 | 7.5-7.6 | 10.0-10.2 |
| Mandible length | 10.7-11.8 | 9.3-10.0 | 13.3-13.9 |

Note: All measurements are in mm; NA – Data not available; † - Condyllobasal length

Andersen, K. (1912). Catalogue of the Chiroptera in the collection of the British Museum, vol. I: Megachiroptera. British Museum (Natural History) London, 854pp.

Andersen, K. (1918). Diagnoses of new bats of the families Rhinolophidae and Megadermatidae. Annals and Magazine of Natural History 2: 374-384.

Andersen, K. & R.C. Wroughton (1907). On bats of the family Megadermatidae. Annals and Magazine of Natural History 19: 129-145.

Anderson, J. (1881). Catalogue of the Mammalia in the Indian Museum, Calcutta. Part I. Primates, Prosimiae, Chiroptera and Insectivora. Calcutta.

Barrett-Hamilton, G.E.H. (1906). Description of two new species of *Pterygistes*. Annals and Magazine of Natural History 17: 98-100.

Barrett-Hamilton, G.E.H. (1907). Description of two new species of *Plecotus*. Annals and Magazine of Natural History 20: 520-521.

Bates, P.J.J. & D.L. Harrison (1997). Bats of the Indian Subcontinent. Harrison Zoological Museum Publications, Sevenoaks, UK, 258pp.

Bates, P.J.J., D.L. Harrison & M. Muni (1994a). The Bats of Western India - Part 1. Journal of the Bombay Natural History Society 91(1): 1-15.

Bates, P.J.J., D.L. Harrison & M. Muni (1994b). The Bats of Western India - Part 2. Journal of the Bombay Natural History Society 91(2): 224-240.

Bates, P.J.J., D.L. Harrison & M. Muni (1994c). The Bats of Western India - Part 3: Journal of the Bombay Natural History Society 91(3): 360-380.

Bates, P.J.J., D.L. Harrison, N.M. Thomas & M. Muni (1994d).

The Indian fruit bat *Latidens salimalii* Thonglongya, 1972 (Chiroptera: Pteropodidae) rediscovered in southern India. Bonner Zoologische Beitrage 45(2): 89-98.

Bhat, H.R. (1968a). Dobson's long nosed fruit bat, *Eonycteris spelaea* (Dobson), from Kumaon Hills, U.P.: an addition to the chiropteran fauna of India. Journal of the Bombay Natural History Society 64: 550-551.

Bhat, H.R. (1968b). *Sphaerias blanfordi* (Thomas, 1891) from Himalayan region of Uttar Pradesh: an addition to the chiropteran fauna of India. Journal of the Bombay Natural History Society 65: 471-473.

Bhat, H.R. (1974). Records and observations on bats of Himalayan region of Uttar Pradesh and West Bengal, India. Journal of the Bombay Natural History Society 71(1): 51-57.

Bhattacharyya, T.P. (1975). Occurrence of Dobson's long tongued fruit bat, *Eonycteris spelaea* (Dobson) (Mammalia: Chiroptera: Pteropidae) in the Andaman Islands, India. Science Culture 41(7): 317-318.

Blanford, W.T. (1888-91). The Fauna of British India, Mammalia. Taylor & Francis, London, 617pp. (in two parts).

Blyth, E. (1846). Notes on the fauna of the Nicobar Islands. Journal of the Asiatic Society of Bengal 15: 367-369.

Blyth, E. (1863). Catalogue of the Mammalia in the Museum Asiatic Society. Calcutta, 187+xiii pp.

Bogdanowicz, W. (1994). *Myotis daubentonii*. Mammalian Species 475: 1-9.

Brosset, A. (1962a). The bats of central and western India - Part I. Journal of the Bombay Natural History Society 59: 1-57.

Brosset, A. (1962b). The bats of central and western India - Part II. Journal of the Bombay Natural History Society 59:

- 583-624.
- Brosset, A. (1962c).** The bats of central and western India - Part III. *Journal of the Bombay Natural History Society* 59: 707-746.
- Brosset, A. (1963).** The bats of central and western India. Part. IV. *Journal of the Bombay Natural History Society* 60: 337-355.
- Carter, T.D. (1943).** The Mammals of the Vernay-Chindwin Expedition, Northern Burma. *Bulletin of the American Museum Natural History* 82(4): 95-114.
- Chakraborty, S. (1975).** On a collection of Mammals from Bhutan. *Records of the Zoological Survey of India* 68: 1-20.
- Chakraborty, S. (1983).** Contribution to the knowledge of the mammalian fauna of Jammu and Kashmir, India. *Records of the Zoological Survey of India Miscellaneous Publications, Occasional Paper No. 38:* 1-129.
- Chasen, F.N. (1940).** A handlist of Malaysian mammals. *Bulletin Raffles Museum* 15: 209
- Corbet, G.B. (1978).** *The Mammals of the Palaearctic Region: a taxonomic review.* British Museum (Natural History), London, 314 pp.
- Corbet, G.B. & J.E. Hill (1986).** *A World List of Mammalian Species.* British Museum (Natural History), London, 226pp.
- Corbet, G.B. & J.E. Hill (1992).** *The Mammals of the Indomalayan Region.* British Museum (Natural History)/ Oxford University Press, London, 488pp.
- Csorba, G. & P.J.J. Bates (1995).** A new subspecies of the Horseshoe bat *Rhinolophus macrotis* from Pakistan (Chiroptera, Rhinolophidae). *Acta Zoologica Academia Scientorum Hungaricae* 41(3): 285-293.
- Das, P.K. (1986).** Studies on the taxonomy and geographical distribution of the species of bat obtained by the Silent Valley (Kerala, India) expedition, 1980. *Records of the Zoological Survey of India* 84(1): 259-276.
- Das, P.K. & Y.P. Sinha (1995).** Occurrence of the Japanese Pipistrelle, *Pipistrellus abramus* (Temminck, 1840) (Chiroptera: Vespertilionidae) in Myanmar (Burma) and India. *Journal of the Bombay Natural History Society* 92: 252-254.
- Das, P.K., R.K. Ghose, T.K. Chakraborty, T.P. Bhattacharyya & M.K. Ghosh (1995).** Mammalia, pp.23-128. In: Director, ZSI (ed.). *State Fauna Series 4: Fauna of Meghalaya, Part 1.* Zoological Survey of India, Calcutta.
- Das, P.K., J.P. Lal & V.C. Agrawal (1993).** Mammalia. pp.143-180. In: Director, ZSI (ed.), *State Fauna Series I: Fauna of Orissa, Part 4.* Zoological Survey of India, Calcutta.
- Dobson, G.E. (1872).** Brief description of five new species of rhinolophine bats. *Journal of the Asiatic Society of Bengal* 41: 336-338.
- Dobson, G.E. (1873a).** Description of a new species of *Vesptilio* from the northwestern Himalaya. *Journal of the Asiatic Society of Bengal* 42(2): 205-206.
- Dobson, G.E. (1873b).** On the genera *Murina* and *Harpyiocephalus* of Gray. *Proceedings of the Asiatic Society of Bengal*, pp.107-110.
- Dobson, G.E. (1874a).** On the Asiatic species of Molossi. *Journal of the Asiatic Society of Bengal* 43(2): 142-144.
- Dobson, G.E. (1874b).** List of Chiroptera inhabiting the Khasia Hills, with description of new species. *Journal of the Asiatic Society of Bengal* 43(2): 234-236.
- Dobson, G.E. (1874c).** Descriptions of new species of Chiroptera from India and Yunan. *Journal of the Asiatic Society of Bengal* 43(2): 237-238.
- Dobson, G.E. (1875).** Description of new species of Vespertilionidae. *Annals and Magazine of Natural History* 16: 260-262.
- Dobson, G.E. (1876).** *Monograph of the Asiatic Chiroptera and catalogue of the species of bats in the collection of the Indian Museum, Calcutta.* London.
- Dobson, G. E. (1878).** *Catalogue of the Chiroptera in the collection of the British Museum.* London, 567pp.
- Ellerman, J.R. & T.C.S. Morrison-Scott (1951).** *Checklist of Palaearctic and Indian Mammals 1758 to 1946.* British Museum (Nat Hist), London, 810pp.
- Finn, F. (1929).** *Sterndale's mammalia of India.* Thacker, Spink & Co., Madras, 347pp.
- Fry, T.B. (1925).** Report No 37a: Nepal. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 30: 525-530.
- Fry, T.B. (1928).** Report No 46: On a collection from Toungoo, Burma. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 30: 545-547.
- Gharaibeh, B.M. & M.Z. Qumsiyeh (1995).** *Otonycteris hemprichii.* *Mammalian Species* 514: 1-4.
- Gray, J.E. (1870).** *Catalogue of monkeys, lemurs, and fruit-eating bats in the collection of the British Museum.* Trustees of the British Museum, London (dated 1870 but published 1871). i-viii, 1-137.
- Hill, J.E. (1958).** Some observations on the fauna of the Maldives Islands. Part II. Mammals. *Journal of the Bombay Natural History Society* 55: 3-10.
- Hill, J.E. (1961).** Indo-Australian bats of the genus *Tadarida.* *Mammalia* 25: 29-56.
- Hill, J.E. (1962).** Notes on some insectivores and bats from Upper Burma. *Proceedings of the Zoological Society of London* 139(1): 119-137.
- Hill, J.E. (1963).** Occurrence of the European free-tailed bat [*Tadarida teniotis* (Rafinesque)] (Chiroptera: Molossidae) in India. *Journal of the Bombay Natural History Society* 60: 723-725.
- Hill, J.E. (1965).** Asiatic bats of the genera *Kerivoula* and *Phoniscus* (Vespertilionidae), with a note on *Kerivoula aerosa* Temes. *Mammalia* 29: 524-556.
- Hill, J.E. (1966).** A review of the genus *Philetor* (Chiroptera: Vespertilionidae). *Bulletin of the British Museum (Natural History), Zoology Series* 14: 371-387.
- Hill, J.E. (1967).** The bats of the Andaman and Nicobar Island. *Journal of the Bombay Natural History Society* 64(1): 1-9.
- Hill, J.E. (1971a).** A note on *Pteropus* (Chiroptera: Pteropidae) from the Andaman Islands. *Journal of the Bombay Natural History Society* 68(1): 1-8.
- Hill, J.E. (1971b).** The bats of Aldabra Atoll, western Indian Ocean. *Philosophical Transactions of the Royal Society (B)* 260: 573-577.
- Hill, J.E. (1972).** The status of *Vesptilio brachypterus* Temminck, 1840 (Chiroptera: Vespertilionidae). *Zoologische Mededelingen, Leiden* 45(12): 139-146.
- Hill, J.E. (1974).** A review of *Scotoecus* Thomas, 1901 (Chiroptera: Vespertilionidae). *Bulletin of the British Museum of Natural History (Zoology)* 27(4): 169-188.

- Hill, J. E. (1976a).** Bats referred to *Hesperoptenus* Peters, 1869 (Chiroptera: Vespertilionidae) with the description of a new subgenus. *Bulletin of the British Museum of Natural History (Zoology)* 32: 1-28.
- Hill, J.E. (1976b).** Further records of *Myotis peshwa* (Thomas 1915) (Chiroptera: Vespertilionidae) from the Indian peninsula. *Journal of the Bombay Natural History Society* 73(3): 433-437.
- Hill, J.E. (1977).** A review of the Rhinopomatidae (Mammalia: Chiroptera). *Bulletin of British Museum Natural History (Zoology)* 32(2): 29-43.
- Hill, J.E. (1983).** Bats (Mammalia: Chiroptera) from Indo-Australia. *Bulletin of British Museum Natural History (Zoology)* 45: 103-208.
- Hill, J.E. (1986).** A note on *Rhinolophus pearsoni* Horsfield, 1851 and *Rhinolophus yunanensis* Dobson, 1872 (Chiroptera: Rhinolophidae). *Journal of the Bombay Natural History Society* 83(Supl.) (Centenary issue): 12-18.
- Hill, J.E. & D.L. Harrison (1987).** The baculum in the Vespertilioninae (Chiroptera: Vespertilionidae) with a systematic review, a synopsis of *Pipistrellus* and *Eptesicus*, and the descriptions of a new genus and subgenus. *Bulletin of British Museum Natural History (Zoology)* 52: 225-305.
- Hill, J.E. & J.D. Smith (1984).** Bats: A natural history. British Museum (Natural History), London, 243pp.
- Hinton, M.A.C. & T.B. Fry (1923).** Report No. 37: Nepal. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 29: 399-428.
- Hinton, M.A.C. & H.M. Lindsay (1926).** Report No 41: Assam and Mishmi Hills. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 31: 383-403.
- Hinton, M.A.C. & O. Thomas (1926).** Report No 42: Kashmir and Punjab. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 31: 606-614.
- Hutson, A.M., S.P. Mickleburgh & P.A. Racey (compilers) (2001).** *Microchiropteran Bats: Global Status, Survey and Conservation Action Plan*. International Union for Conservation of Nature and Natural Resources, Chiroptera Specialist Group (IUCN/SSC). Gland, Switzerland.
- Hutton, A.F. (1949a).** Notes on snakes and mammals of the High Wavy Mountains, Madurai District, S. India. *Journal of the Bombay Natural History Society* 48: 454-460.
- Hutton, A.F. (1949b).** Notes on snakes and mammals of the High Wavy Mountains, Madurai District, S. India. *Journal of the Bombay Natural History Society* 48: 681-694.
- Hutton, T. (1872).** On the bats of the north-western Himalayas. *Proceeding zool. Society Lond.*, pp.690-714.
- Jerdon, T.C. (1874).** *The Mammals of India: a natural history of all the animals known to inhabit continental India*. John Wheldon, London, 335 pp.
- Johnson, D.H., S.D. Ripley & K. Thonglongya (1980).** Mammals from Nepal. *Journal of the Bombay Natural History Society* 77: 56-63.
- Khajuria, H. (1953).** Taxonomic studies on some Indian Chiroptera. *Records of the Indian Museum* 50: 113-128.
- Khajuria, H. (1970).** A new Leaf nosed bat from central India. *Mammalia* 34: 622-627.
- Khajuria, H. (1979a.)** Taxonomical and ecological studies on bats of Jabalpur District Madhya Pradesh, India. Part. I. (Families Pteropidae, Rhinopomatidae and Emballonuridae). *Records of the Zoological Survey of India Miscellaneous Publications, Occasional Paper No. 13*: 1-59.
- Khajuria, H. (1980).** Taxonomical and ecological studies on bats of Jabalpur District Madhya Pradesh, India. Part. 11. Families Megadermatidae, Rhinolophidae and Vespertilionidae. *Records of the Zoological Survey of India Miscellaneous Publications, Occasional Paper No. 19*: 1-69.
- Khajuria, H. (1984).** Notes on some central Indian bats. *Journal of the Zoological Society of India* 36 (1-2): 125-126.
- Khajuria, H., Y. Chaturvedi & D.K. Ghosal (1977).** Annotated catalogue of type specimens of mammals in zoological survey of India. *Records of the Zoological Survey of India Miscellaneous Publications, Occasional Paper No. 7*: 1-45.
- Khajuria, H. & D.K. Ghosal (1981).** Studies on wildlife of Narbada Valley. Part 4. Mammals. *Records of the Zoological Survey of India* 79(1-2): 235-257.
- Khan, R. (1985).** *Mammals of Bangladesh*. Nanna Reza, Dhaka.
- Kock, D. & H.R. Bhat (1994).** *Hipposideros hypophyllus* n. sp. of the *H. bicolor*-group from peninsular India (Mammalia: Chiroptera: Hipposideridae). *Senckenbergiana biologica* 73(1-2): 25-31.
- Koopman, K.F. (1993).** Order Chiroptera In: D. E. Wilson & D. M. Reeder (eds.), *Mammal Species of the World*, 2nd ed. Smithsonian Institution Press, Washington. 1206 pp. pp. 137-241.
- Kurup, G.U. (1968).** Mammals of Assam and adjoining areas. *Proceedings of the zoological Society of Calcutta* 21: 79-99.
- Lal, J.P. (1982).** Andersen's Rufous bat, *Rhinolophus rouxi sinicus* Andersen (Chiroptera: Rhinolophidae) from Arunachal Pradesh, India. *Journal of the Bombay Natural History Society* 79(2): 402.
- Lindsay, H.M. (1926a).** [i] Report No 38: Sind [40-42]; [ii] Report No 39: Mergui Archipelago [42-48]. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 30: 40-48.
- Lindsay, H.M. (1926b).** Report No 40: Gwalior; Bombay Natural History Society's Mammal Survey of India. *Journal of the Bombay Natural History Society* 31: 379-382.
- Lindsay, H.M. (1927).** [i] Report No 43: Nelliampathy plateau and Palni Hills [591-597]; [ii] Report No 44: Kangra and Chamba [597-606]; [iii] Report No 45: The Punjab Salt Range and Murree [606-614]; Bombay Natural History Society's Mammal Survey of India. *Journal of the Bombay Natural History Society* 31: 591-614.
- Meuhauer, H.H. (1970).** First positive record of *Pipistrellus savii* (Chiroptera: Vespertilionidae) from India. *Journal of the Bombay Natural History Society* 67(2): 319-320.
- Mickleburgh, S.P., A.M. Hutson & P.A. Racey (compilers) (1992).** *Old World fruit bats. An action plan for their conservation*. International Union for Conservation of Nature and Natural Resources, Chiroptera Specialist Group (IUCN/SSC). Gland, Switzerland, 252pp.
- Mickleburgh, S.P., A.M. Hutson & P.A. Racey (2002).** A review of the global conservation status of bats. *Oryx* 36(1): 18-34.
- Miller, G.S. (1902).** The mammals of the Andaman and Nicobar Islands. *Proceedings of the US National Museum* 24(1296):

- 751-795.
- Miller, G.S. (1903).** Seventy new Malaysian mammals. *Smithsonian Miscellaneous Collections* 45: 1-73.
- Miller, G.S. (1906a).** Twelve new genera of bats. *Proceedings of the Biological Society of Washington* 19: 83-86.
- Miller, G.S. (1906b).** Seven new Malayan bats. *Proceedings of the Biological Society of Washington* 19: 61-65.
- Miller, G.S. (1907).** The families and genera of bats. *Bulletin of the US National Museum* 57: xxvii, 282pp.
- Mills, I.C.S. (1921).** Report No 36: Naga Hills. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 29: 221-229.
- Mirza, Z.B. (1965).** Four new mammal records for West Pakistan. *Mammalia* 29: 205-210.
- Mitchell, R.M. (1980).** New records of bats (Chiroptera) from Nepal. *Mammalia* 44(3): 339-342.
- Molur, S., G. Marimuthu, C. Srinivasulu, S. Mistry, A.M. Hutson, P.J.J. Bates, S. Walker, K.P. Priya & A.R.B. Priya (eds.) (2002).** *Status of South Asian Chiroptera: Conservation Assessment and Management Plan (C.A.M.P.) Workshop Report*. Zoo Outreach Organisation, Conservation Breeding Specialist Group South Asia, and Wildlife Information & Liaison Development Society, Coimbatore, India, viii+154pp.+CD.
- Nath, B. (1952).** On a collection of mammals from Assam (India) with special reference to the rodents. *Records of the Indian Museum* 50: 271-285.
- Nath, S. (1979).** On a collection of microchiropterid bats from Kashmir Valley and adjacent areas, with some new records. *Proceedings of the 66th Indian Science Congress* III(D): 3.
- Phillips, W.W.A. (1923).** Further notes on some Ceylon bats. *Journal of the Bombay Natural History Society* 29: 154-156.
- Phillips, W.W.A. (1924).** A guide to the Mammals of Ceylon: Chiroptera. *Spolia Zeylanica* 13: 1-63.
- Phillips, W.W.A. (1980).** *Manual of the Mammals of Sri Lanka. Part I*. Wildlife & Nature Protection Society of Sri Lanka, 1-116pp.
- Prakash, I. (1961).** Taxonomic and biological observations on the bats of the Rajasthan desert. *Records of the Indian Museum* 59: 149170.
- Prakash, I. (1963).** Zoogeography and evolution of the mammalian fauna of Rajasthan desert, India. *Mammalia* 27: 342-351.
- Prater, S.H. (1914).** Notes on Wroughton's Free-tailed bat (*Otomops wroughtoni*). *Journal of the Bombay Natural History Society* 22: 788.
- Roberts, T.J. (1977).** *The Mammals of Pakistan*. Ernest Benn Ltd, 361pp.
- Ryley, K.V. (1913a).** Report No 9: Mysore. Bombay Natural History Society's Mammal Survey of India. *Journal of the Bombay Natural History Society* 22(2): 283-295.
- Ryley, K.V. (1913b).** [i] Report No 10: Kathiawar [464-486]; [ii] Report No 11: Coorg [486-513]. Bombay Natural History Society's Mammal Survey of India. *Journal of the Bombay Natural History Society* 22: 464-513.
- Ryley, K.V. (1914a).** Report No 12: Palanpur & Mount Abu. Bombay Natural History Society's Mammal Survey of India. *Journal of the Bombay Natural History Society* 22(4): 684-699.
- Ryley, K.V. (1914b).** [i] Report No 13: South Ceylon [700-710]; [ii] Report No 14: North Shan States, Burma [710-725]. Bombay Natural History Society's Mammal Survey of India. *Journal of the Bombay Natural History Society* 22(4): 700-725.
- Saha, S.S. (1979).** A supplementary note on some mammals recently collected in Bhutan. *Journal of the Bombay Natural History Society* 76(1): 155-156.
- Saha, S.S. (1980).** Notes on some mammals recently collected from the Andaman and Nicobar Islands. *Records of the Zoological Survey of India* 77: 119-126.
- Saha, S.S. (1984).** Occurrence of the tail-less fruit bat, *Megaerops ecaudatus* (Temminck, 1837), in Namdapha, Tirap District, Arunachal Pradesh: an addition to the Indian Fauna (Mammalia: Chiroptera: Pteropodidae). *Bulletin zool. surv. India* 6(I-3): 343-344.
- Saha, S.S. (1985).** Mammalia. In: Fauna of Namdapha: Arunachal Pradesh 'A proposed biosphere reserve'. *Records of the Zoological Survey of India* 82(I-4): 321-330.
- Scully, J. (1887).** On the Chiroptera of Nepal. *Journal Asiatic Society Bengal* 56: 233-259.
- Siddiqi, M.S. (1961).** Checklist of Mammals of Pakistan with particular reference to the Mammalian Collection in the British Museum (Natural History), London. *Biologia* 7(1-2): 93-225.
- Sinha, Y.P. (1970).** Taxonomic notes on some Indian bats. *Mammalia* 34: 81-92.
- Sinha, Y.P. (1973).** Taxonomic studies on the Indian horseshoe bats of the genus *Rhinolophus* Lacépède. *Mammalia* 37(4): 603-630.
- Sinha, Y.P. (1980).** The bats of Rajasthan: taxonomy and zoogeography. *Records of the Zoological Survey of India* 76(I-4): 7-63.
- Sinha, Y.P. (1986).** The Bats of Bihar: taxonomy and field ecology. *Records of the Zoological Survey of India Miscellaneous Publications, Occasional Paper* 77: 1-60.
- Sinha, Y.P. (1990).** Occurrence of Dobson's Long-tongued Fruitbat *Eonycteris spelaea* (Dobson, 1871) (Chiroptera: Pteropodidae) in Meghalaya. *Journal of the Bombay Natural History Society* 87: 134.
- Sinha, Y.P. (1994a).** Occurrence of the long-tongued fruit bat *Eonycteris spelaea* (Dobson, 1871) in Manipur and Nagaland, India. *Geobios New Reports* 13(2): 186-187.
- Sinha, Y.P. (1994b).** Occurrence of the Kashmir cave bat, *Myotis longipes* (Dobson, 1873) in Meghalaya, India. *Geobios New Reports* 13(1): 68.
- Srinivasulu, C. & B. Srinivasulu (2001).** Bats of the Indian subcontinent. *Current Science* 80(11): 1378-1380.
- Tate, G.H.H. (1941a).** A review of the genus *Hipposideros* with special reference to Indo-Australian species. *Bulletin of the American Museum of Natural History* 78: 353-393.
- Tate, G.H.H. (1941b).** Notes on vespertilionid bats. *Bulletin of the American Museum of Natural History* 78: 567-597.
- Thomas, N.M. (2000).** Morphological and mitochondrial-DNA variation in *Rhinolophus rouxii* (Chiroptera). *Bonner Zoologische Beiträge* 49(1-4): 1-18.
- Thomas, O. (1897).** On some bats obtained from the Surat and Thana districts. *Journal of the Bombay Natural History Society* 11: 274-276.
- Thomas, O. (1913).** On a remarkable new free-tailed bat from southern Bombay. *Journal of the Bombay Natural History Society* 22: 87-91.

- Thomas, O. (1915a).** Scientific Results from the Mammal Survey No 10: The Indian Bats assigned to the genus *Myotis*. *Journal of the Bombay Natural History Society* 23: 607-612.
- Thomas, O. (1915b).** On pipistrelles of the genera *Pipistrellus* and *Scotozous*. *Journal of the Bombay Natural History Society* 24: 29-36.
- Thomas, O. (1915c).** On bats of the genera *Nyctalus*, *Tylonycteris* and *Pipistrellus*. *Annals and Magazine of Natural History* 16: 225-232.
- Thomas, O. (1915d).** A special genus for the Himalayan bat known as *Murina grisea*. *Annals and Magazine of Natural History* 16: 309-3 10.
- Thomas, O. (1915e).** Notes on *Taphozous* and *Saccopteryx*. *Journal of the Bombay Natural History Society* 24: 57-63.
- Thomas, O. (1916a).** Scientific results from the Mammal Survey, XIII. *Journal of the Bombay Natural History Society* 24: 404-430.
- Thomas, O. (1916b).** Scientific results from the Mammal Survey, XIV. *Journal of the Bombay Natural History Society* 24: 639-644.
- Thomas, O. (1920).** A new bat of the genus *Myotis* from Sikkim. *Journal of the Bombay Natural History Society* 27: 248-250.
- Thomas, O. (1921).** The geographical races of *Scotomantulus ornatus*. *Journal of the Bombay Natural History Society* 27: 772-773.
- Thomas, O. (1923).** On the forms contained in the genus *Harpiocephalus*. *Journal of the Bombay Natural History Society* 29: 88-89.
- Thomas, O. (1926).** A new *Myotis* from Ladak. *Annals and Magazine of Natural History* 17: 609-610.
- Thonglongya, K. (1972).** A new genus and species of fruit bat from south India (Chiroptera: Pteropodidae). *Journal of the Bombay Natural History Society* 69(1): 151-158.
- Tiwari, K.K., R.K. Ghose & S. Chakraborty (1971).** Notes on a collection of small mammals from Western Ghats, with remarks on the status of *Rattus rufescens* (Gray) and *Bandicota indica malabarica* (Shaw). *Journal of the Bombay Natural History Society* 68: 378-384.
- Topal, G. (1970).** The first record of *Ia io* Thomas, 1902 in Vietnam and India, and some remarks on the taxonomic position of *Parascotomantulus beauforti* Bourret, 1942, *Ia longimana* Pen, 1962, and the genus *Ia* Thomas, 1902 (Chiroptera: Vespertilionidae). *Opuscula Zoologica, Budapest* 10: 341-347.
- Topal, G. (1971).** The taxonomic position of *Myotis dobsoni* (Trouessart, 1879) and some statistical data to the subspecific examination of *Myotis blythii* (Tomes, 1857). *Annales Historico-Naturales Musei Nationalis Hungarici (Budapest)* 63: 383-399.
- Topal, G. (1975).** Bacula of some Old World Leaf nosed bats (Rhinolophidae and Hipposideridae, Chiroptera: Mammalia). *Vertebrata Hungarica* 15: 21-45.
- Topal, G. (1997).** A new mouse-eared bat species, from Nepal, with statistical analysis of some other species of subgenus *Leuconycteris* (Chiroptera, Vespertilionidae). *Acta Zoologica Academiae Scientiarum Hungaricae* 43(4): 375-402.
- Walton, D.W. (1974).** New records of bats (Chiroptera) from Pakistan. *Journal of the Mammalian Society of Japan* 6(2): 43-50.
- Wroughton, R.C. (1899).** Some Konkan bats. *Journal of the Bombay Natural History Society* 12: 716-725.
- Wroughton, R.C. (1912a).** Report No 1: East Khandesh. *Bombay Natural History Society's Mammal Survey of India. Journal of the Bombay Natural History Society* 21(2): 391-410.
- Wroughton, R.C. (1912b).** [i] Report No 2: Berars [820-825]; [ii] Report No 3: Cutch [826-844]; [iii] Report No 4: Nimir [844-851]. *Bombay Natural History Society's Mammal Survey of India. Journal of the Bombay Natural History Society* 21(3): 820-851.
- Wroughton, R.C. (1912c).** Report No 5: Dharwar. *Bombay Natural History Society's Mammal Survey of India. Journal of the Bombay Natural History Society* 21(4): 1170-1195.
- Wroughton, R.C. (1912d).** Some new Indian mammals. *Journal of the Bombay Natural History Society* 21: 767-773.
- Wroughton, R.C. (1913).** [i] Report No 6: Kanara [29-44]; [ii] Report No 7 (with K.V. Ryley): Central Provinces [45-58]; [iii] Report No 8 (with K.V. Ryley): Vijayanagar [58-66]. *Bombay Natural History Society's Mammal Survey of India. Journal of the Bombay Natural History Society* 22(1): 29-66.
- Wroughton, R.C. (1914).** Report No 15: Kumaon. *Bombay Natural History Society's Mammal Survey of India. Journal of the Bombay Natural History Society* 22(4): 282-301.
- Wroughton, R.C. (1915a).** Report No 16: Dry Zone, Central Burma & Mount Popa. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 23: 460-480.
- Wroughton, R.C. (1915b).** Report No 17: South Tennasserim, Burma. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 23: 695-720.
- Wroughton, R.C. (1915c).** [i] Report No 18: Ceylon [79-96]; [ii] Report No 19: Bengal, Bihar & Orissa [96-110]. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 24(1): 79-1 10.
- Wroughton, R.C. (1916a).** [i] Report No 20: Chindwin River [291-309]; [ii] Report No 21: Gwalior [309-310]; [iii] Report No 22: Koyna Valley [311-316]. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 24: 291-316.
- Wroughton, R.C. (1916b).** Report No 23: Sikkim & Bengal Terai. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 24: 468-493.
- Wroughton, R.C. (1916c).** [i] Report No 24: Sind [749-758]; [ii] Report No 25: Chin Hills [758-773]; [iii] Report No 26: Darjiling District [773-781]. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 24(1): 749-781.
- Wroughton, R.C. (1917a).** Report No 27: Bhutan Duars. *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 25(1): 63-71.
- Wroughton, R.C. (1917b).** Report No 28: Kalimpong (Darjiling). *Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. Journal of the Bombay Natural History Society* 25: 274-279.
- Wroughton, R.C. (1918a).** Summary of the results from the Indian mammal survey. Part 1. *Journal of the Bombay*

- Natural History Society* 25: 547-598.
- Wroughton, R.C. (1918b).** Summary of the results from the Indian mammal survey. Part 2. *Journal of the Bombay Natural History Society* 26: 19-58.
- Wroughton, R.C. (1920).** Report No 32: Baluchistan. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 27: 314-322.
- Wroughton, R.C. (1921).** [i] Report No 33: High Wavy Mountains, Madura District [545-549]; [ii] Report No 34: Travancore [549-553]; [iii] Report No 35: Prome [553-554]; Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 27: 545554.
- Wroughton, R.C. & W.M. Davidson (1918).** Report No 29: Pegu. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 25: 472-481.
- Wroughton, R.C. & W.M. Davidson (1920).** [i] Report No 30: Dekhan, Poona District [1025-1030]; [ii] Report No 31: Nilgiris [1031-1035]. Bombay Natural History Society's Mammal Survey of India, Burma and Ceylon. *Journal of the Bombay Natural History Society* 25: 1025-1035.
- Wroughton, R.C. & K.V. Ryley (1913).** Scientific results from the Mammals Survey, III. *Journal of the Bombay Natural History Society* 22: 13-21.

