

OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

NOTE

GENTIANA APERTA (GENTIANACEAE) - A NEW RECORD TO INDIA FROM LADAKH HIMALAYA

Mohd Shabir, Priyanka Agnihotri, Jay Krishan Tiwari & Tariq Husain

26 August 2018 | Vol. 10 | No. 9 | Pages: 12286–12289

10.11609/jott.4233.10.9.12286-12289



For Focus, Scope, Aims, Policies and Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints contact info@threatenedtaxa.org

Partners



المحمد بن زايد
للمحافظة على
الكائنات الحية
The Mohamed bin Zayed
SPECIES CONSERVATION FUND



Member



Publisher & Host





ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

OPEN ACCESS



Gentiana L. is the largest and most diverse genus of family Gentianaceae, consisting of c. 362 species (Ho & Liu 2001; Mabberley 2008; Shabir et al. 2017b) and is largely distributed in the meadows of temperate, sub-alpine and alpine regions in Asia, Europe and North America and a few species occur in the Andes of South America, Central America, eastern Australia, and northwestern Africa (Ho & Liu 2001; Struwe & Albert 2002). In India, the genus is represented by 68 species (Gupta et al. 2012; Maity 2014; Shabir et al. 2017a,b) mainly distributed in alpine and sub-alpine meadows of both the eastern and western Himalaya.

During the ongoing revisionary study on the genus *Gentiana* in the Indian Himalaya, some interesting specimens of the genus were collected from alpine slopes of Ldokchan and Spang-rings of Tumail in Kargil District of Jammu & Kashmir, India, at an elevation of 3900–4200 m. After critical examination of the specimens, we identified the species as *Gentiana aperta* Maxim., which has not been recorded in the Indian flora so far. Therefore, a detailed description along with photo plates and other relevant information of the species has been provided to facilitate its easy identification.

Gentiana aperta

Maxim., Bull. Acad. Imp. Sci. Saint-Pétersbourg 3, 27: 500. 1881; T.N. Ho & S. Liu, Worldwide Monogr. *Gentiana*: 386. 2001; T.N. Ho & J.S. Pringle in Z.Y. Wu & P.H. Raven, Fl. China 16: 86. 1995 (Image 1).

GENTIANA APERTA (GENTIANACEAE) - A NEW RECORD TO INDIA FROM LADAKH HIMALAYA

Mohd Shabir¹ , Priyanka Agnihotri² ,
Jay Krishan Tiwari³ & Tariq Husain⁴

^{1,2,4} Plant Diversity, Systematics & Herbarium Division, CSIR-National Botanical Research Institute, Lucknow, Uttar Pradesh 226001, India

^{1,3} Plant Systematics Laboratory, Department of Botany and Microbiology, HNB Garhwal University, Srinagar Garhwal, Uttarakhand 246174, India

¹shabir1610@gmail.com (corresponding author), ²priyagni_2006@yahoo.co.in, ³jktiwari31@rediffmail.com, ⁴hustar_2000@yahoo.co.uk

Type: China, Qinghai (as W Kansu, Tangut region), to Huangsui river (fl. Rako-gol), 10,000–11,000 ped., in meadows, in 1880, Przewalski *s.n.* (holotype: LE not seen; isotypes: K, P Images)

Annual herbs, 3.0–6.0 cm high. Stem prostrate to ascending, striate, branched from the base. Basal leaves wither on anthesis, leaf blades ovate, 3.0–4.0 × 1.6–2.0 mm, apex obtuse, margin indistinctly membranous, veins distinct; cauline leaves, widely spaced, 3–6 paired, elliptic, spatulate to oblong, 2.5–3.5 × 1.0–1.5 mm, apex acute, mid-vein distinct, margin membranous. Inflorescence terminal, solitary; pedicels 3.0–4.0 mm long. Calyx 4.0–5.0 mm long; tube 2.8–3.4 mm long; lobes more or less equal, ovate, 1.2–1.5 × 0.6–0.8 mm, apex acute. Corolla white, pale blue to blue, 5.0–7.0 mm long, dark spot in the throat, tube 4.5–4.8 mm long; lobes ovate-oblong, 1.0–1.5 × 0.7–0.8, apex obtuse to sub-rounded, margin entire; plicae 0.6–0.9 mm long, 2-cleft, apex acute, margin entire. Stamens 5; filaments

DOI: <https://doi.org/10.11609/jott.4233.10.9.12286-12289>

Editor: P. Lakshminarasimhan, Botanical Survey of India, Pune, India.

Date of publication: 26 August 2018 (online & print)

Manuscript details: Ms # 4233 | Received 03 May 2018 | Final received 09 July 2018 | Finally accepted 25 July 2018

Citation: Shabir, M., P. Agnihotri, J.K. Tiwari & T. Husain (2018). *Gentiana aperta* (Gentianaceae) - a new record to India from Ladakh Himalaya. *Journal of Threatened Taxa* 10(9): 12286–12289; <https://doi.org/10.11609/jott.4233.10.9.12286-12289>

Copyright: © Shabir et al. 2018. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: University Grants commission, New Delhi has provided financial assistance in terms of fellowship to the first author (MS) to carry out the research work.

Competing interests: The authors declare no competing interests.

Acknowledgements: We are thankful to the Director, CSIR-National Botanical Research Institute, Lucknow for the necessary facilities and encouragement. The first author (MS) is also thankful to the UGC, New Delhi for providing fund in terms of fellowship (UGC-SRF) to carry out the research work.





Image 1. *Gentiana aperta* Maxim.: A– C - habitats; D - habit; E–F - flowers.

inserted near the middle of the corolla, 2.0–4.0 mm long; anther ellipsoid to orbicular, $0.35\text{--}0.45 \times 0.18\text{--}0.30$ mm. Style short; stigma bifid, recurved, lobes semi-orbicular; ovary $2.5\text{--}2.7 \times c. 1.8$ mm. Capsules 3.6–4.3 mm long; stipe 0.8–1.2 mm long. Seeds ellipsoid, $0.7\text{--}0.8 \times 0.28\text{--}0.33$ mm; seed coat reticulate.

Flowering & Fruiting: June–September.

Habitat: The species was found growing on the west facing alpine slopes of Ldokchan and Sprang-rings in Tumail, Kargil of the Ladakh Himalaya, India, associated with *Gentiana leucomelaena* Maxim, *Gentiana aquatica* L., *Gentianopsis detonsa* (Rottb.) Ma, and *Gentianella tumailica* M. Shabir, Agnihotri, Tiwari & Husain.

Distribution: China (Ho & Pringle 1995; Ho & Liu 2001), new to India.

Specimens examined: 309906 (LWG), India, Jammu & Kashmir, Ladakh, Kargil, Tumail, 3,900–4,000 m, 12.viii.2016, coll. Mohd Shabir; 309908 (LWG), India, Jammu and Kashmir, Ladakh, Kargil, Tumail, 4,200m, 12.viii.2016, coll. Mohd Shabir; China: Gansu, Regio Tangut, 30.viii.1980, N.M. Przewalski s.n. (P); Gansu, Regio Tangut, 30.viii.1980, N.M. Przewalski s.n. (K).

Conservation status: Data deficient (DD).

Taxonomic notes: *Gentiana aperta* Maxim., belonging to Section *Chondrophyllae* Bunge, under the genus *Gentiana* was described by Maximowicz (1881) from Gansu, China. In the present state of our knowledge, this species is so far known only from China, and is endemic to the mountains of northeastern Qinghai and northwestern Gansu. The species growing in the Indian Himalaya is characterized by white to bluish-white flowers, apex of corolla lobes acute to sub-acute, dark-blue spot on the corolla throat, spots less dense on the throat and plicae deeply bifurcate, 2-cleft with both segments acute and entire, whereas, the plants growing in China have flowers bluish-white, apex of corolla lobes obtuse, yellowish-white spots on the corolla throat and spots scattered densely up to the base. *Gentiana aperta* is allied to *Gentiana leucomelaena* Maxim. In India, *G. leucomelaena* is distributed in Jammu & Kashmir, Himachal Pradesh, Uttarakhand, and the Sikkim Himalaya, and differs from *G. aperta* in having calyx lobes lanceolate to linear-lanceolate, apex acuminate, mid-vein prominent, plicae oblong, apex obtuse and



Image 2. Map showing existing and new localities of *Gentiana aperta* Maxim.




Image 3. Herbarium image of *Gentiana aperta* Maxim., (Mohd Shabir 309908 (LWG)).




Image 4. Herbarium image of *Gentiana aperta* Maxim., (Mohd Shabir 309906 (LWG)).

margin irregularly laciniate.

Further, *G. aperta* also shows a taxonomic affinity with *Kuepferia pringlei* D. Maity & S.K. Dey, in the nature of habit and floral characters, but plicae well developed, as long as or near to the corolla lobe, apex

deeply segmented with both segments acute and entire, not forming auricle, corolla lobes much shorter than the tube differentiate, the former from the latter. The report of *G. aperta* from the Ladakh Himalaya extends its range of distribution further southwestward.

References

- Gupta, S.A., S.K. Mukherjee & M. Mondal (2012).** A census of *Gentiana* L. in India, systematics of flowering plants, pp. 53–58. In: Maiti, G. & S.K. Mukherjee (eds.). *Multidisciplinary approaches in Angiosperm Systematics*. University of Kalyani, Kalyani.
- Ho, T.N. & J.S. Pringle (1995).** Gentianaceae, pp. 86–87. In: Wu, Z.Y. & P.H. Raven (eds.). *Flora of China (Gentianaceae through Boraginaceae)*. Beijing: Science Press, and St Louis: Missouri Botanical Garden Press.
- Ho, T.N. & S. Liu (2001).** *A Worldwide Monograph of Gentiana*. Science Press. Beijing, China.
- Mabberley, D.J. (2008).** *The Plant Book. A Portable Dictionary of Plants, their Classification and Uses - 3rd Edition*. Cambridge University Press, Cambridge, 354pp.
- Maity, D. (2014).** A new species of *Gentiana* L. (Gentianaceae) from Sikkim Himalaya. *Edinburgh Journal of Botany* 71: 289–296; <http://doi.org/10.1017/S096042861400016X>
- Maximowicz, C.J. (1981).** Diagnosis plantarum novarum asiaticarum. 1V. Scripsit. *Bulletin de l'Academie imperial des Science de ST-Peterbourg, sér* 327(4): 500–501.
- Shabir, M., P. Agnihotri, D. Husain, J.K. Tiwari & T. Husain (2017a).** On the current status of the genus *Gentiana* L. (Gentianaceae) in India. *Pleione* 11(1): 16–24.
- Shabir, M., P. Agnihotri, D. Husain, J.K. Tiwari & T. Husain (2017b).** Lectotypification of the names of three species of *Gentiana* (Gentianaceae) occurring in India. *Phytotaxa* 324 (3): 293–297; <http://doi.org/10.11646/phytotaxa.324.3.6>
- Struwe, L. & A. Albert (2002).** *Gentianaceae: Systematics and Natural History*. Cambridge University Press, Cambridge, 227pp.





OPEN ACCESS



The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

August 2018 | Vol. 10 | No. 9 | Pages: 12147–12298

Date of Publication: 26 August 2018 (Online & Print)

DOI: 10.11609/jott.2018.10.9.12147-12298

www.threatenedtaxa.org

Article

Appearances are deceptive: molecular phylogeny recovers the Scaly Gecko *Hemidactylus scabriceps* (Reptilia: Squamata: Gekkonidae) as a member of a scansorial and rupicolous clade

-- Achyuthan N. Srikanthan, Gandla Chethan Kumar, Aishwarya J. Urs & Sumaithangi Rajagopalan Ganesh, Pp. 12147–12162

Communications

Foraging and roosting ecology of the Lesser Dog-faced Fruit Bat *Cynopterus brachyotis* (Mammalia: Chiroptera: Pteropodidae) in southern India

-- T. Karuppudurai & K. Sripathi, Pp. 12163–12172

Diversity and status of avifauna in man-made sacred ponds of Kurukshetra, India

-- Parmesh Kumar & Archana Sharma, Pp. 12173–12193

Diversity and distribution of freshwater turtles (Reptilia: Testudines) in Goa, India

-- Trupti D. Jadhav, Nitin S. Sawant & Soorambail K. Shyama, Pp. 12194–12202

Breeding behaviour of the Coromandel Damselfly *Ceriatagris coromandelianus* (Fabricius) (Zygoptera: Coenagrionidae) in central India: copulation

-- Nilesh R. Thakkar, Payal R. Verma & Raymond J. Andrew, Pp. 12203–12209

The status assessment of *Corynandra viscosa* subsp. *nagarjunakondensis* (Magnoliopsida: Cleomaceae), endemic to Nagarjunakonda, Andhra Pradesh, India

-- Veeravarapu Hanumantha Rao, Vaidyula Vasudeva Rao, Anuti Baleeshwar Reddy & Vatsavaya Satyanarayana Raju, Pp. 12210–12217

Short Communications

New records of termites (Blattodea: Termitidae: Syntermitinae) from Colombia

-- Olga Patricia Pinzón & Daniel Castro, Pp. 12218–12225

New reports of thrips (Thysanoptera: Terebrantia: Thripidae) from India

-- R.R. Rachana & R. Varatharajan, Pp. 12226–12229

New records of earthworm fauna (Oligochaeta: Glossoscolecidae and Megascolecidae) collected from Satkosia-Baisipalli Wildlife Sanctuary of Odisha, India

-- Rinku Goswami, Pp. 12230–12234

Diversity and endemism of butterflies of montane forests of Eravikulam National Park in the Western Ghats, India

-- E.R. Sreekumar, S. Nikhil, K.G. Ajay & P.O. Nameer, Pp. 12235–12246

Partners



Member



Publisher & Host

