OPEN ACCESS

The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

COMMUNICATION

NEW KISSING BUG (HEMIPTERA: REDUVIIDAE: TRIATOMINAE) RECORDS FROM NAPO AND MORONA-SANTIAGO PROVINCES WITH DISTRIBUTION UPDATES IN ECUADOR

Ana Soto-Vivas, Sandra Enríquez, Ernesto Villacrés, Jazzmin Arrivillaga, Martín Hinojosa & Jonathan Liria

26 October 2018 | Vol. 10 | No. 11 | Pages: 12515-12522
10.11609/jott.4345.10.11.12515-12522

For Focus, Scope, Aims, Policies and Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>
For Article Submission Guidelines visit <http://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>
For Policies against Scientific Misconduct visit <http://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>
For reprints contact info@threatenedtaxa.org

Partners

المحمد بن زايد
للمحافظة على
الكائنات الحية
The Mohamed bin Zayed
SPECIES CONSERVATION FUND

Member

Publisher & Host

NEW KISSING BUG (HEMIPTERA: REDUVIIDAE: TRIATOMINAE) RECORDS FROM NAPO AND MORONA-SANTIAGO PROVINCES WITH DISTRIBUTION UPDATES IN ECUADOR

Ana Soto-Vivas¹ , Sandra Enríquez² , Ernesto Villacrés³ , Jazzmin Arrivillaga⁴ ,
Martín Hinojosa⁵ & Jonathan Liria⁶

ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

OPEN ACCESS

¹ Carrera de Biología, Facultad de Ciencias Biológicas, Universidad Central del Ecuador, Quito 170129, Ecuador

^{2,3} Instituto de Investigación en Salud Pública y Zoonosis (CIZ), Universidad Central del Ecuador, Quito 170129, Ecuador

⁴ Área de Ambiente, Turismo Histórico Cultural, Facultad de Comunicación Social, Universidad Central del Ecuador, Quito 170129, Ecuador

⁵ Carrera Ingeniería en Ecosistemas, Universidad Regional Amazónica Ikiam, Vía Tena, Muyuna Kilómetro 7, Napo, Ecuador

⁶ Grupo de Investigación en Población y Ambiente, Universidad Regional Amazónica Ikiam, Vía Tena, Muyuna Kilómetro 7, Napo, Ecuador

¹ aysoto@uce.edu.ec, ² sandrabycid@gmail.com, ³ ernestovillacresguevara@yahoo.es, ⁴ jcarrivillaga@uce.edu.ec,

⁵ martin.hinojosa@ikiam.edu.ec, ⁶ jonathan.liria@gmail.com (corresponding author)

Abstract: Four species of Triatominae (Hemiptera: Reduviidae) are reported in four localities in Ecuadorian Amazonia. *Eratyrus mucronatus* Stål, 1859, *Rhodnius pictipes* Stål, 1872 & *Panstrongylus geniculatus* (Latreille, 1811) in Napo Province, and *Rhodnius robustus* Larrousse, 1927 in Morona-Santiago Province. Two specimens of *R. pictipes* were found in an urban area of Tena City (capital province). These findings can indicate a risk of Chagas disease transmission in urban and peri-urban areas of the Amazonia.

Keywords: Chagas disease, new records, Rhodniini, Triatomini.

Spanish Abstract: Se reportan cuatro especies de Triatominae (Hemiptera: Reduviidae) en cuatro localidades en la Amazonía ecuatoriana. *Eratyrus mucronatus* Stål, 1859, *Rhodnius pictipes* Stål, 1872 y *Panstrongylus geniculatus* (Latreille, 1811) en la provincia de Napo, y *Rhodnius robustus* Larrousse, 1927 en la Provincia de Morona-Santiago. Dos especímenes de *R. pictipes* se encontraron en el área urbana de la ciudad de Tena (provincia capital). Estos hallazgos pueden indicar un riesgo de transmisión de la enfermedad de Chagas en áreas urbanas y periurbanas de la Amazonía.

DOI: <https://doi.org/10.11609/jott.4345.10.11.12515-12522> | **ZooBank:** urn:lsid:zoobank.org:pub:0B7A9E1F-0043-4D56-925A-9A1F9D188BBC

Editor: Anonymity requested.

Date of publication: 26 October 2018 (online & print)

Manuscript details: Ms # 4345 | Received 27 June 2018 | Final received 20 September 2018 | Finally accepted 01 October 2018

Citation: Soto-Vivas, A., S. Enríquez, E. Villacrés, J. Arrivillaga, M. Hinojosa & J. Liria (2018). New kissing bug (Hemiptera: Reduviidae: Triatominae) records from Napo and Morona-Santiago provinces with distribution updates in Ecuador. *Journal of Threatened Taxa* 10(11): 12515–12522; <https://doi.org/10.11609/jott.4345.10.11.12515-12522>

Copyright: © Soto-Vivas et al. 2018. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use of this article in any medium, reproduction and distribution by providing adequate credit to the authors and the source of publication.

Funding: Facultad de Ciencias Biológicas - Universidad Central del Ecuador, Centro Internacional de Zoonosis - Universidad Central del Ecuador, Universidad Regional Amazónica Ikiam.

Competing interests: The authors declare no competing interests.

Author Details: see end of this article.

Author Contribution: ASV conducted the Triatominae identification and wrote the first manuscript draft. JL, EV and MH conducted the specimens collections. ASV, JA, SE and JL wrote the final manuscript. SE prepared the specimen photographs. JL elaborated the distribution maps.

Acknowledgements: The authors thank Pablo Araujo for the triatomine specimens photographs.

INTRODUCTION

The triatomine bugs are characterized by the hematophagous habit and morphological adaptations associated with the blood feeding. These insects are the main vectors of Chagas Disease or American Trypanosomiasis (Lent & Wygodzinski 1979; World Bank 1993). The Triatominae subfamily comprises five tribes, 17 genera, and 150 species, where *Triatoma* Laporte, 1832 and *Rhodnius* Stål 1859, are the most important vectors. Ecuador currently contains about 17 species such as the following: *Triatoma dimidiata* (Latreille, 1811), *Triatoma carrioni* Larrousse, 1926, *Triatoma venosa* (Stål, 1872), *Triatoma dispar* Lent, 1950, *Eratyrus mucronatus* Stål, 1859, *Eratyrus cuspidatus* Stål, 1859, *Cavernicola pilosa* Barber, 1937, *Panstrongylus geniculatus* (Latreille, 1811), *Panstrongylus rufotuberculatus* (Champion, 1899), *Panstrongylus howardi* (Neiva, 1911), *Panstrongylus chinai* (Del Ponte, 1929), *Panstrongylus lignarius* (Walker, 1873), *Pastrongylus herreri* (Wygodzinsky, 1948), *Rhodnius ecuadoriensis* Lent & León, 1958, *Rhodnius pictipes* Stål, 1872, *Rhodnius robustus* Larrousse, 1927, and was recently described *Rhodnius barretti* Abad-Franch et al. 2013 (Abad-Franch et al. 2001; Galvão et al. 2003; Abad-Franch et al. 2013; Vaca-Moncayo et al. 2017).

Triatoma dimidiata is the main vector in Ecuador associated with the Chagas disease transmission in Guayas and Manabí provinces; this species is distributed in Los Ríos, El Oro, Pichincha and Bolívar provinces. On the other hand, *R. ecuadoriensis* is associated with the transmission in Loja and El Oro provinces and is reported in Manabí, Guayas and Los Ríos too. Finally, *T. carrioni* is reported in Azuay, Cañar, Loja, El Oro, Zamora Chinchipe, and recently in Pichincha (Grijalva et al. 2003). Other species with less importance are *P. rufotuberculatus*, *P. chinai*, *P. geniculatus*, *P. lignarius*, *P. howardi*, *T. venosa*, *T. dispar*, *E. mucronatus*, *E. cuspidatus* and *C. pilosa*; however in the Amazonia *R. pictipes* and *R. robustus* are sylvatic vectors (Aguilar et al. 1999; Abad-Franch et al. 2001; Galvão et al. 2003; Vaca-Moncayo et al. 2017). Due to this, we reported new triatomine records in two Ecuadorian Amazonia Provinces with update distribution in Ecuador.

MATERIAL AND METHODS

This study is based on, one male specimen of *E. mucronatus*, three males specimens of *R. pictipes* and one female specimen of *P. geniculatus*, found dead

in the main entrance of the Universidad Regional Amazónica Ikiam (0.954°S & 77.862°W); more recently, two dead female specimens of *R. robustus* were found in the Universidad Regional Amazónica Ikiam (same coordinates). Later, two additional dead female specimens of *R. pictipes* were found in two houses of the urban area of Tena City (0.989°S & 77.827°W and 0.987°S & 77.812°W). The first specimen was found inside the house in the living room close to the kitchen, and the second specimen was found outside the house on the third floor in the department entrance. Houses are surrounded by secondary forest patches at 20–125 m and 200–278 m to the Tena River. All the insects were pinned, mounted and the identification was determined using taxonomic keys of Lent & Wygodzinsky (1979), Carcavallo et al. (1998a) and Soto-Vivas (2009). In *R. pictipes*, the male genitalia were dissected following the morphological description of Bérenger & Pluot-Sigwalt (2002). The triatomine specimens were deposited at the entomological collection of Instituto de Investigación en Salud Pública y Zoonosis in the Universidad Central del Ecuador: “Colección Nacional de Referencia de Artrópodos de Importancia en Zoonosis (CONRAZ)”. Finally, the four specimens of *R. robustus* collected in Yuwientza (2.067°S & 77.883°W), a Shuar community located in Morona Santiago Province was verified by CONRAZ.

RESULTS AND DISCUSION

The list of species with diagnostic characters, distribution and medical importance are given below:

Family Reduviidae

Subfamily Triatominae

Tribe Triatomini

Eratyrus mucronatus Stål, 1859 (Image 1)

Diagnostic characters: Rostrum with first and second segments larger (subequal in length), third segment short; anterior process of scutellum form a sharply pointed spine (the spine as long as the entire scutellum); fore lobe of pronotum with 1+1 strong discal spines; humeral angles distinctly spinose; subapical reddish spot of corium comparatively small, anteriorly not attaining level of m-cu cross-vein; free portion of vesica flattened apically in side view; abdomen dorsally with five prominent tubercles along each urotergites (Lent & Wygodzinsky 1979; Soto-Vivas 2009).

Material examined: EC-N-M-T25, 01.ix.2016, 1 male, Campus Universidad Regional Amazónica Ikiam (7km to Muyuna), Napo, 0.954°S & 77.862°W, 600m, coll. Conraz.

Distribution (Carcavallo et al. 1999; Galvão et al. 2003; Chávez 2006; Guhl et al. 2007; Bérenger et al. 2009; Morocoima et al. 2010; Meneguetti et al. 2011; Obara et al. 2013; Hiwat 2014; Galvão 2014; Ceccarelli et al. 2018): Bolivia (Beni, La Paz, Chuquisaca, Oruro, Potosí); Brazil (Acre, Amazonas, Goiás, Maranhão, Mato Grosso, Pará, Rondonia, Tocantins); Colombia (Antioquia, Boyacá, Casanare, Cundinamarca, Meta, Norte Santander, Vaupés); Ecuador (Esmeraldas, Loja, Napo, Orellana, Sucumbíos), Guatemala, Guiana, French Guiana (Cayenne, Grand Santi, Montsinéry-Tonnegrande, Mana, Régina, Roura, Saul, Kourou, Sinnamary); Panama, Perú (Junín, Madre de Dios, Loreto, Ucayali, San Martín), Suriname (Brokopondo, Paramaribo, Para, Sipaliwini); Trinidad (Tunapuna-Piarco, San Juan-Laventille); Venezuela (Anzoátegui, Aragua, Amazonas, Barinas, Carabobo, Cojedes, Guárico, Falcón, Lara, Mérida, Monagas, Portuguesa, Sucre, Táchira, Trujillo, Yaracuy, Zulia).

Medical importance: *Eratyrus mucronatus* has been found naturally infected with *Trypanosoma cruzi* (Chagas 1909); It was reported in small domestic colonies attracted to light in Andean areas of Bolivia. Also, it was recorded in domiciliation process, invading urban and rural areas near fragmented forests (Soto-Vivas et al. 2001; Abad-Franch et al. 2009).

***Panstrongylus geniculatus* (Latreille, 1811) (Image 2)**

Diagnostic characters: Length more than 20mm; head comparatively short and stout, in lateral view; posterior process of scutellum elongate subcylindrical, narrowly tapering apically; pronotum with extensive, conspicuous black markings; all connexival segments with light and dark marking; abdomen light colored ventrally, with longitudinal series of black spots (Lent & Wygodzinsky 1979; Soto-Vivas 2009).

Material examined: EC-N-M-T34, 22.viii.2018, 1 female, Campus Universidad Regional Amazónica Ikiam (7km to Muyuna), Napo, 0.954°S & 77.862°W, 600m, coll. Conraz.

Distribution (Carcavallo et al., 1999; Galvão et al. 2003; Chávez 2006; Guhl et al. 2007; Leite et al. 2007; Bérenger et al. 2009; Morocoima et al. 2010; Meneguetti et al. 2011; Obara et al. 2013; Hiwat 2014; Galvão 2014; Ceccarelli et al. 2018): Argentina (Chaco, Corrientes, Entre Ríos, Formosa, Misiones, Santa Fe, Santiago Del Estero); Bolivia (Beni, Cochabamba, Santa Cruz, Tarija); Brazil (Acre, Amapá, Amazonas, Bahia, Ceará, Distrito Federal, Espírito Santo, Goiás, Maranhão, Mato Grosso, Mato Grosso do Sul, Minas Gerais, Pará, Paraná, Piauí, Rio de Janeiro, Rondônia, Roraima, São Paulo, Tocantins); Colombia (Antioquia, Amazonas, Bolívar, Boyacá, Cauca, Casanare, Cesar, Cundinamarca, Magdalena, Huila, Meta, Norte de Santander, Putumayo, Santander, Sucre, Tolima, Valle del Cauca); Costa Rica (Alajuela, Cartago, Guanacaste, Heredia, Limón, Puntarenas, San José); Ecuador (Esmeraldas, Imbabura, Manabí, Napo, Orellana, Pastaza, Pichincha, Sucumbíos); French Guiana (Cayenne, Saint-Laurent-du-Maroni); Guatemala, Guyana (Cuyuni-Mazaruni); Mexico (Chiapas, Veracruz, Yucatán); Nicaragua (Atlántico Norte, Atlántico Sur, Managua, Río San Juan); Panama (Bocas del Toro, Colón, Los Santos, Panamá); Paraguay (Alto Paraná, Boquerón, Caaguazú, Concepción, Paraguari); Peru (Amazonas, Ayacucho, Cajamarca, Cusco, Huánuco, Junín, Madre de Dios, Loreto, Pasco, San Martín, Ucayali); Suriname (Brokopondo, Commewijne, Para, Paramaribo, Saramacca, Sipaliwini, Wanica); Trinidad & Tobago (Diego Martín, San Juan-Laventille, Sangre Grande, Siparia, Tunapuna-Piarco); Uruguay; Venezuela (Anzoátegui, Amazonas, Aragua, Barinas, Bolívar, Carabobo, Delta Amacuro, Distrito Capital, Falcón, Guárico, Lara, Mérida, Miranda, Monagas,

© Pablo Araujo & Sandra Enríquez

Image 1. Dorsal view of *Eratyrus mucronatus* (male) collected at Campus Universidad Regional Amazónica Ikiam, Napo.

Trujillo, Táchira, Yaracuy, Vargas, Zulia).

Medical importance: *Panstrongylus geniculatus* is a widely distributed species occupying natural habitats like burrows; this species shows association with *Dasytus novemcinctus* Linnaeus, 1758 and *Didelphis marsupialis* Linnaeus, 1758 (Abad-Franch et al. 2001) and rodents. It is believed that it is responsible for maintaining the enzootic cycle of *T. cruzi*.

Tribe Rhodniini

Rhodnius pictipes Stål, 1872 (Image 3)

Diagnostic characters: Head laterally behind the eyes with callosities setiferous tubercles; antenna inserted proximal to head; anteocular region of the head at least 2.7 times as long as postocular; pronotum very granulate, rugose; femora yellowish, mottled with dark brown; tibiae of all pairs of legs with dark submedian annulus; corium brownish, irregularly spotted with black; rectangular dark spots of dorsal connexival segments with conspicuous pointed projection posteriorly, at least on segments 3 to 5; process of pygophore bispinous with short base (Lent & Wygodzinsky 1979; Béranger & Pluot-Sigwalt 2002; Soto-Vivas 2009).

Material examined: EC-N-M-T26, EC-N-M-T27, EC-N-M-T28, 03.iii.2016, 3 males, Campus Universidad Regional Amazónica Ikiam (7km to Muyuna), Napo, 0.954°S & 77.862°W, 600m, coll. CONRAZ. EC-N-T-T29, 23.iii.2018, 1 female, Barrio El Dorado (Tena City), Napo, 0.987°S & 77.812°W, 500m, CONRAZ. EC-N-T-T30, 20.v.2018, 1 female Barrio San Antonio (Tena City), Napo, 0.989°S & 77.827°W, 517m, coll. CONRAZ.

Distribution (Carcavallo et al. 1999; Abad-Franch

et al. 2001; Galvão et al. 2003; Chávez 2006; Cortez et al. 2007; Guhl et al. 2007; Hiwat 2014; Galvão 2014; Ceccarelli et al. 2018): Bolivia (Cochabamba, Santa Cruz, Beni, Pando); Belize, Brazil (Amapá, Amazonas, Goiás, Maranhão, Mato Grosso, Pará, Piauí, Roraima, Tocantins); Colombia (Amazonas, Boyacá, Caquetá, Cundinamarca, Guaviare, Meta, Norte Santander, Putumayo, Vaupés); Ecuador (Azuay, Morona Santiago, Napo, Orellana, Sucumbíos); Guiana, French Guiana (Cayenne, Saint-Laurent du Maroni); Peru (Ayacucho, Cusco, Huánuco, Madre de Dios, Loreto, Ucayali, San Martín, Junín); Suriname (Brokopondo, Commewijne, Coronie, Marowijne, Paramaribo, Para, Saramacca, Sipaliwini, Wanica); Trinidad (Diego Martín, Rio Claro-Mayaro, Siparia, Tanapuma-Piarco); Venezuela (Anzoátegui, Amazonas, Apure, Aragua, Bolívar, Carabobo, Cojedes, Delta Amacuro, Falcón, Mérida, Miranda, Monagas, Portuguesa, Táchira, Trujillo, Sucre, Yaracuy, Zulia).

Medical importance: Occasionally attracted to light in human dwellings. It has been found naturally infected with *T. cruzi*. (Carcavallo et al. 1999; Feliciangeli et al. 2004a; Abad-Franch et al. 2009).

Rhodnius robustus Larrousse, 1927 (Image 4)

Diagnostic characters: Head laterally behind the eyes with callosities setiferous tubercles; antenna inserted proximal to the head apex; anteocular region about four times as long as postocular; distance between eyes dorsally smaller than, or equal to, width of eye; head distinctly longer than pronotum (1:0.65–0.80); median process of pygophore narrow, pointed apically and with narrow triangular base; larger species, length of males

Image 2. Dorsal view of *Panstrongylus geniculatus* (female) collected at Campus Universidad Regional Amazónica Ikiam, Napo.

© Pablo Araujo & Sandra Enríquez

20–23.5 mm, of females 23–26 mm (Lent & Wygodzinsky 1979; Soto-Vivas 2009).

Material examined: EC-MS-Y-T17, EC-MS-Y-T18, EC-MS-Y-T19, EC-MS-Y-T20, 14.xi.2009, 3 males and 1 female, Yuwientza “Shuar community”, Morona Santiago, 2.067°S & 77.883°W, 1,126m, coll. Conraz; EC-N-M-T32, 12.vii.2018, 1 female, Campus Universidad Regional Amazónica Ikiám (7km to Muyuna), Napo, 0.954°S & 77.862°W, 600m, coll. Conraz; EC-N-M-T33, 23.viii.2018, 1 female, Campus Universidad Regional Amazónica Ikiám (7km to Muyuna), Napo, 0.954°S & 77.862°W, 600m, coll. Conraz.

Distribution: (Carcavallo et al. 1999; Abad-Franch et al. 2001; Galvão et al. 2003; Chávez 2006; Cortez et al. 2007; Guhl et al. 2007; Bérenger et al. 2009; Hiwat, 2014; Galvão 2014; Ceccarelli et al. 2018): Bolivia (Cochabamba, Beni, La Paz, Santa Cruz, Pando); Brazil (Acre, Amapá, Amazonas, Goiás, Maranhão, Mato Grosso, Pará, Rondonia, Roraima, Tocantins); Colombia (Amazonas, Arauca, Bolívar, Cundinamarca, Norte Santander, Santander, Tolima); Ecuador (Napo, Morona Santiago, Sucumbíos, Orellana); French Guiana (Cayenne, Macouria, Matoury, Grand-Santi, Regina), Perú (Amazonas, Cajamarca, Junín, Madre de Dios, Loreto, San Martín, Ucayali); Suriname (Brokopondo, Marowijne, Para, Saramacca, Sipaliwini); Venezuela (Apure, Barinas, Bolívar, Cojedes, Falcón, Mérida, Monagas, Sucre, Táchira, Trujillo, Yaracuy).

Medical importance: This species is closely related to *R. prolixus*, and has been associated to sylvatic habits and infected with *T. cruzi*. In northern South America, they were found migrating from palms to the houses, occurring where *R. prolixus* was eliminated by Chagas disease control programs (Monteiro et al. 2003; Guhl et al. 2007; Longa & Scorza 2007).

Epidemiological significance

The Amazonian Chagas disease transmission has increased in recent years; the first reports were in Brazil by Coura et al. (1994, 1995, 2002) and Albajar et al. (2003); recently Santalla et al. (2011), stated a case from Bolivian Amazonia. In Ecuador, the first records were registered by Aguilar & Yépez (1995) in three Amazonian Provinces: Sucumbíos, Napo, and Pastaza. More recently, Amunárriz et al. (2010), reported a population infected by *T. cruzi* located between the margins of Napo and Aguarico rivers. These authors mentioned three triatomine species involved: *P. geniculatus*, *R. pictipes* and *R. robustus*. Abad-Franch & Monteiro (2007) stated that all Amazonian triatomine species comprises four tribes: Rhodniini, Bolbolderini, Cavernicolini, and Triatomini (with *Eratyrus*).

Eratyrus mucronatus is a sylvatic species responsible for the enzootic Chagas transmission (Morocoima et al. 2010) and reported occasionally in the peridomicile with a synanthropic trend (Noireau et al. 1995; Carcavallo et

© Pablo Araujo & Sandra Enríquez

Image 3. Dorsal view of *Rhodnius pictipes* (male) collected at Campus Universidad Regional Amazónica Ikiám, Napo.

Figure 1. New records from Napo and Morona Santiago Provinces and distribution in Ecuador based on Abad-Franch et al. (2001): A - *E. mucronatus* (m), *R. pictipes* (p) and *R. robustus* (r), and B - *P. geniculatus* (g).

Image 4. Dorsal view of *Rhodnius robustus* (male) collected at Yuwientza "Shuar community", Morona Santiago.

al. 1998b; Soto-Vivas et al. 2001).

Panstrongylus geniculatus is a sylvatic species associated with animal burrows and trees that provide microclimate conditions for their survival (Herrera & Urdaneta-Morales 1992; 1997). In the last decade, this species has been associated with domiciliated environments, frequently found in chicken coops and invading homes attracted by light or in search of food (Felicangeli et al. 2004b). Also, it has been found colonizing *Rattus rattus* Linnaeus, 1758 burrows in precarious dwellings (Reyes-Lugo 2009). Although it is true that in urban environments this species prefers to feed on chickens, dogs, cats, and synanthropic rodents, it is important to keep the entomological vigilance and its possible implication in the transmission of *T. cruzi*. (Herrera et al. 2003).

Sylvatic *Rhodnius* species distribution is related to the Arecaceae (Palms), and their feeding habits are associated with birds; this triatomine species are specialized to explore the Palm microhabitats (Lent & Wygodzinsky 1979; Abad-Franch et al. 2009). Also, several authors stated the association between *Rhodnius* and birds nest of *Phacellodomus rufifrons* (Wied-Neuwied, 1821) and other Furnariidae species (Lent & Jurberg 1975; Abad-Franch & Monteiro 2007).

Abad-Franch et al. (2001) studied the biogeography of Ecuadorian triatomine based on distribution maps for each species. We georeferenced all Orellana/Napo records for *E. mucronatus*, *R. pictipes* and *P. geniculatus*, and the entire records correspond only to Orellana based on the Provinces boundaries (Fig. 1 A,B); also, we checked all triatomine records found in the Zoology Museum (QCAZ) on-line data base at the Pontificia Universidad Católica del Ecuador (QCAZ 2018). Due to this it was recorded for the first time that three species in Napo, and also the *R. robustus* distribution has spread out to Morona Santiago Province.

Finally, the land use changes, the wild source feed availability and triatomine species competence, can be a risk of disease transmission in urban and peri-urban areas of the Ecuadorian Amazonia. Due to this it is relevant to implement programs for entomological vigilance for the Chagas disease.

REFERENCES

- Abad-Franch, F., A. Paucar, C. Carpio, C. Cuba, M. Aguilar & M. Miles (2001). Biogeography of Triatominae (Hemiptera: Reduviidae) in Ecuador: implications for the design of control strategies. *Memoria do Instituto Oswaldo Cruz* 96: 611–620; <https://doi.org/10.1590/S0074-02762001000500004>
- Abad-Franch, F. & F. Monteiro (2007). Biogeography and evolution of Amazonian triatomines (Heteroptera: Reduviidae): implications for Chagas disease surveillance in humid forest ecoregions. *Memoria do Instituto Oswaldo Cruz* 102: 57–70; <https://doi.org/10.1590/S0074-02762007005000108>
- Abad-Franch, F., F. Monteiro, N. Jaramillo, R. Gurgel-Gonçalves, F. Dias & L. Diotaiuti (2009). Ecology, evolution, and the long-term surveillance of vector-borne Chagas disease: A multi-scale appraisal of the tribe Rhodniini (Triatominae). *Acta Tropica* 110: 159–167; <https://doi.org/10.1016/j.actatropica.2008.06.005>
- Abad-Franch, F., M. Pavan, N. Jaramillo, F. Palomeque, C. Dale, D. Chaverra & F. Monteiro (2013). *Rhodnius barretti*, a new species of Triatominae (Hemiptera: Reduviidae) from western Amazonia. *Memoria do Instituto Oswaldo Cruz* 108: 92–99; <https://doi.org/10.1590/0074-0276130434>
- Aguilar, M. & R. Yépez (1995). Evolución epidemiológica de la Enfermedad de Chagas en el Ecuador. In: Schofield C.J., J.P. Dujardin & J. Jurberg (eds.). Proceedings of the International Workshop on Population genetics and control of Triatominae. Santo Domingo de Los Colorados, Ecuador, 24–28 September, 115pp.
- Aguilar, H., F. Abad-Franch & J. Racines (1999). Epidemiology of Chagas disease in Ecuador. A brief review. *Memoria do Instituto Oswaldo Cruz* 94: 387–393; <https://doi.org/10.1590/S0074-02761999000700076>
- Amunárriz, M., S. Quito, V. Tandazo & M. López (2010). Seroprevalencia de la enfermedad de Chagas en el cantón Aguarico, Amazonia ecuatoriana. *Revista Panamericana de Salud Publica* 28: 25–29; <https://doi.org/10.1590/S1020-49892010000700004>
- Bérenger, J. & D. Pluot-Sigwalt (2002). *Rhodnius amazonicus* Almeida, Santos & Sposina, 1973, Bona species, close to *R. pictipes* Stal, 1872 (Heteroptera, Reduviidae, Triatominae). *Memoria do Instituto Oswaldo Cruz* 97: 73–77; <https://doi.org/10.1590/S0074-02762002000100011>
- Bérenger, J., D. Pluot-Sigwalt, F. Pagès, D. Blanchetis & C. Aznar (2009). The triatominae species of French Guiana (Heteroptera: Reduviidae). *Memoria do Instituto Oswaldo Cruz* 104: 1111–1116; <https://doi.org/10.1590/S0074-02762009000800007>
- Ceccarelli, S., A. Balsalobre, P. Medone, M. Cano, R. Gurgel, D. Felicangeli, D. Vezzani, C. Wisnivesky-Colli, D. Gorla, G. Marti & J. Rabinovich (2018). DataTri, a database of American triatomine species occurrence. *Scientific Data* 5: 180071.
- Carcavallo R., I. Galíndez, J. Jurberg, C. Galvão & D. Canale (1998a). Pictorial Keys for tribes, genera and species of the subfamily Triatominae, pp. 107–244. In: Carcavallo, R., I. Galíndez-Girón, J. Jurberg & H. Lent (eds.). *Atlas of Chagas Disease Vectors in the Americas, Vol. III*. Editora Fiocruz, Rio de Janeiro, RJ, Brazil, ix+1217pp.
- Carcavallo R., F. Rodríguez, R. Salvatella, S. Curto de Casas, I. Sherlock, C. Galvão, S. de Rocha, I. Galíndez, M. Otero, A. Matinez, D. Canale, T. Farr & J. Barata (1998b). Habitats and related fauna. pp 561–660. In: Carcavallo, R., I. Galíndez-Girón, J. Jurberg & H. Lent (eds.). *Atlas of Chagas Disease Vectors in the Americas, Vol. III*. Editora Fiocruz, Rio de Janeiro, RJ, Brazil, ix+1217pp.
- Carcavallo, R., S. Curto de Casas, I. Sherlock, I. Galíndez, J. Jurberg, C. Galvão, C. Mena Francois N. (1999). Geographical distribution and alti-latitudinal dispersion pp. 747–838 In: Carcavallo, R., I. Galíndez-Girón, J. Jurberg & H. Lent (eds.). *Atlas of Chagas Disease Vectors in the Americas, Vol. III*. Editora Fiocruz, Rio de Janeiro, RJ, Brazil, ix+1217pp.
- Chávez, J. (2006). Contribución al estudio de los triatomíneos del Perú: Distribución geográfica, nomenclatura y notas taxonómicas. *Anales de la Facultad de Medicina* 67: 65–76; <https://doi.org/10.15381/anales.v67i1.1296>
- Coura, J.R., T.V. Barret & C.M. Naranjo (1994). Ataque de populações humanas por triatomíneos silvestres no Amazonas: uma nova forma de transmissão da infecção Chagásica? *Revista da Sociedade Brasileira de Medicina Tropical* 27: 251–253; <https://dx.doi.org/10.1590/S0037-86821994000400009>
- Coura, J., M. Naranjo & H. Willcox (1995). Chagas' disease in the Brazilian Amazon. II. A serological survey. *Revista do Instituto de Medicina Tropical de São Paulo* 37: 103–107; <https://doi.org/10.1590/S0036-46651995000200003>

- Coura, J., A. Junqueira, M. Boia, O. Fernandes, C. Bonfante, J. Campos, L. Santos & R. Devera (2002). Chagas disease in the Brazilian Amazon: IV. A new cross-sectional study. *Revista do Instituto de Medicina Tropical de São Paulo* 44: 159–165; <https://doi.org/10.1590/S0036-46652002000300009>
- Cortez, M., M. Avalos, V. Rocha & D. Gorla (2007). Distribución biogeográfica de los triatomíneos en Bolivia: Discriminación de la distribución de las especies en relación a variables ambientales, pp. 67–137. In: Cortez, M.R. (ed.). *Triatomíneos de Bolivia y la enfermedad de Chagas*. La Paz: Ministerio de Salud y Deportes. Programa Nacional de Chagas.
- Feliciangeli, M.D., J. Benítez, P. Reyes, C. Maldonado & E. Borges (2004a). ¿Hay Enfermedad de Chagas en la región Amazónica de Venezuela? *Boletín de la Dirección de Malariología y Saneamiento Ambiental* 44: 67–76.
- Feliciangeli M.D., H. Carrasco, J. S. Patterson, B. Suarez, C. Martínez & M. Medina (2004b). Mixed domestic infestation *Rhodnius prolixus* Stal, 1859 and *Panstrongylus geniculatus* Latreille, 1811, vector incrimination, and seroprevalence for *Trypanosoma cruzi* among inhabitants in El Guamito, Lara State, Venezuela. *American Journal of Tropical Medicine and Hygiene* 71: 501–505; <https://doi.org/10.4269/ajtmh.2004.71.501>
- Galvão, C., R. Carvallo, D. Da Silva & J. Jurberg (2003). A Checklist of the current valid species of subfamily Triatominae Jeannel, 919 (Hemiptera, Reduviidae) and their geographical distribution, with nomenclatural and taxonomic notes. *Zootaxa* 202: 1–36; <https://doi.org/10.11646/zootaxa.202.1.1>
- Grijalva, M., L. Escalante, R. Paredes, J. Costales, A. Padilla, E. Rowland & A. Racines (2003). Seroprevalence and risk factors for *Trypanosoma cruzi* Infection in the Amazon Region of Ecuador. *American Journal of Tropical Medicine and Hygiene* 69: 380–385.
- Guhl, F., G. Aguilera, N. Pinto & D. Vergara (2007). Actualización de la distribución geográfica y ecoepidemiología de la fauna de triatomíneos (Reduviidae: Triatominae) en Colombia. *Biomédica* 27: 143–162; <https://doi.org/10.7705/biomedica.v27i1.258>
- Herrera, L. & S. Urdaneta-Morales (1992). *Didelphis marsupialis*: a primary reservoir of *Trypanosoma cruzi* in urban areas of Caracas, Venezuela. *Annals of Tropical Medicine & Parasitology* 68: 607–612; <https://doi.org/10.1080/00034983.1992.11812716>
- Herrera, L. & S. Urdaneta-Morales. (1997). Synanthropic rodent reservoirs of *Trypanosoma (Schizotrypanum) cruzi* in the valley of Caracas, Venezuela. *Revista do Instituto de Medicina Tropical de São Paulo* 39: 279–282; <https://doi.org/10.1590/s0036-46651997000500006>
- Herrera, L., S. Urdaneta-Morales & H. Carrasco (2003). *Trypanosoma cruzi*: comportamiento de metatrimastigotes obtenidos de *Didelphis marsupialis* y *Panstrongylus geniculatus*. *Revista Científica FCV-LUZ* 4:307–11.
- Hiwat, H. (2014). Triatominae species of Suriname (Heteroptera: Reduviidae) and their role as vectors of Chagas disease. *Memoria do Instituto Oswaldo Cruz* 109: 452–458; <https://doi.org/10.1590/0074-0276130408>
- Leite, G., C.B. Dos Santos & A. Falqueto (2007). Insecta, Hemiptera, Reduviidae, *Panstrongylus geniculatus*: geographic distribution map. *Check List* 3: 147–152; <https://dx.doi.org/10.15560/3.2.147>
- Lent, H. & J. Jurberg (1975). O Genero *Panstrongylus* Berg, 1879, com um estudo sobre a genitalia externa das espécies (Hemiptera, Reduviidae, Triatominae). *Revista Brasileira de Biologia* 35: 379–438.
- Lent, H. & P. Wygodzinsky (1979). Revision of the Triatominae (Hemiptera, Reduviidae), and their significance as vectors of Chagas disease. *Bulletin of the American Museum of Natural History* 163: 123–520.
- Longa, A. & J.V. Scorza (2007). Migración de *Rhodnius robustus* (Hemiptera: Triatominae) desde *Acrocomia aculeata* (Palmae) hacia domicilios rurales en Venezuela. *Boletín de Malariología y Salud Ambiental* 47: 213–22.
- Meneguetti, D., O. Trevisan, R. Rosa & L. Camargo (2011). First report *Eratyrus mucronatus*, Stal, 1859, (Hemiptera, Reduviidae, Triatominae), in the State of Rondonia, Brazil. *Revista da Sociedade Brasileira de Medicina Tropical* 44: 511–512; <https://doi.org/10.1590/S0037-86822011000400022>
- Monteiro, F.A., T. Barrett, S. Fitzpatrick, C. Cordon-Rosales, M.D. Feliciangeli & C. Beard (2003). Molecular phylogeography of the Amazonian Chagas disease vectors *Rhodnius prolixus* and *R. robustus*. *Molecular Ecology* 12: 997–1006; <https://doi.org/10.1046/j.1365-294X.2003.01802.x>
- Morocoima, A., J. Chique, L. Herrera & S. Urdaneta-Morales (2010). *Eratyrus mucronatus* (Stal, 1859) (Hemiptera, Reduviidae, Triatominae): Primer registro para el estado Anzoátegui (Venezuela). *Boletín de Malariología y Salud Ambiental* 50: 307–309.
- Noireau, F., M. Bosseno, R. Carrasco, J. Telleria, F. Vargas, C. Camacho, N. Yaksic & S. Frederique Breniér (1995). Sylvatic Triatomines (Hemiptera: Reduviidae) in Bolivia: Trends Toward Domesticity and Possible Infection with *Trypanosoma cruzi* (Kinetoplastida: Trypanosomatidae). *Journal of Medical Entomology* 32: 594–598; <https://doi.org/10.1093/jmedent/32.5.59>
- Obara, M., A. Cardoso, M. Pinto, C. Souza, R. Silva & R. Gurgel-Gonçalves (2013). *Eratyrus mucronatus* Stål, 1859 (Hemiptera: Reduviidae: Triatominae): First report in the State of Acre, Brazil, and updated geographic distribution in South America. *Check List* 9: 851–854; <https://doi.org/10.15560/9.4.851>
- QCAZ (2018). Base de datos del Museo de Zoología (QCAZ). Pontificia Universidad Católica del Ecuador. Available in <http://bioweb.puce.edu.ec/QCAZ/baseDatos>
- Reyes-Lugo, M. (2009). *Panstrongylus geniculatus* Latreille 1811 (Hemiptera: Reduviidae: Triatominae), vector de la enfermedad de Chagas en el ambiente domiciliario del centro-norte de Venezuela. *Rev Biomed* 20: 180–205.
- Santalla, V., C. Oporto, E. Espinoza, T. Rios & L. Brutus (2011). Primer brote reportado de la enfermedad de chagas en la Amazonia Boliviana: reporte de 14 casos agudos por transmisión oral de *Trypanosoma cruzi* en Guayaramerín, Beni-Bolivia. *BIOFARBO* 19: 52–58.
- Soto-Vivas, A. (2009). Clave pictórica de triatomíneos (Hemiptera: Triatominae) de Venezuela. *Boletín de Malariología y Salud Ambiental* 49: 259–274.
- Soto-Vivas, A., H. Barazarte & D. Molina de F. (2001). Primer reporte de *Eratyrus mucronatus* Stal 1959 (Hemiptera: Reduviidae) en ambiente domiciliario. *Entomotropica* 16: 215–217.
- Vaca-Moncayo F., S. Enríquez, J. Arrivillaga-Henríquez., E. Villacrés-Guevara, P. Araujo & W. Benítez-Ortiz (2017). Actualización de la distribución geográfica de *Triatoma dispar* (Hemiptera: Reduviidae: Triatominae) en Ecuador. *Revista Colombiana de Entomología* 43: 255–261.

Author Details: ANA SOTO-VIVAS PhD, is a Lecturer-Researcher at the Central University of Ecuador. Her research interests are medical entomology, geometric morphometrics, and Triatominae systematics and biogeography. SANDRA ENRÍQUEZ MSc Biology, is a researcher at the Public Health and Zoonoses Research Institute (CIZ) of the Central University of Ecuador, Quito. Biologist and Entomologist. Her research interest is the taxonomy of arthropod vectors, biodiversity, taxonomy of aquatic insects and environmental impact. ERNESTO VILLACRES Biologist, is a researcher at CIZ. His research interests are Triatominae and others arthropod vectors. JAZZMIN ARRIVILLAGA PhD, is a Lecturer-Researcher at the Central University of Ecuador. Her research interests are medical entomology and Sandflies systematics and biogeography. MR. MARTÍN HINOJOSA is an undergraduate student of Ecosystem Engineering program of Ikiam. JONATHAN LIRIA PhD, is a Lecturer-Researcher at the Ikiam University. His research interests are medical entomology, geometric morphometrics, and Culicidae systematics and biogeography.

OPEN ACCESS

The Journal of Threatened Taxa is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use of articles in any medium, reproduction, and distribution by providing adequate credit to the authors and the source of publication.

ISSN 0974-7907 (Online); ISSN 0974-7893 (Print)

October 2018 | Vol. 10 | No. 11 | Pages: 12443–12618

Date of Publication: 26 October 2018 (Online & Print)

DOI: 10.11609/jott.2018.10.11.12443-12618

www.threatenedtaxa.org

Article

The terrestrial life of sea kraits: insights from a long-term study on two *Laticauda* species (Reptilia: Squamata: Elapidae) in the Andaman Islands, India
-- Zoya Tyabji, Nitya Prakash Mohanty, Erina Young & Tasneem Khan, Pp. 12443–12450

Communications

Fishing Cat *Prionailurus viverrinus* Bennett, 1833 (Carnivora: Felidae) distribution and habitat characteristics in Chitwan National Park, Nepal
-- Rama Mishra, Khadga Basnet, Rajan Amin & Babu Ram Lamichhane, Pp. 12451–12458

Status distribution and feeding habit of Wild Boar *Sus scrofa* (Mammalia: Artiodactyla: Suidae) in Pench Tiger Reserve, Madhya Pradesh, India
-- Shaheer Khan & Orus Ilyas, Pp. 12459–12463

The composition and status of waterbirds of Perur Lake in Tamil Nadu, India
-- G. Parameswaran & R. Sivashankar, Pp. 12464–12488

The herpetofauna of Jigme Singye Wangchuck National Park in central Bhutan: status, distribution and new records
-- Sangay Tshewang & Letro Letro, Pp. 12489–12498

The Odonata (Insecta) of Dhofar, southern Oman
-- Elaine M. Cowan & Peter J. Cowan, Pp. 12499–12514

New kissing bug (Hemiptera: Reduviidae: Triatominae) records from Napo and Morona-Santiago provinces with distribution updates in Ecuador
-- Ana Soto-Vivas, Sandra Enríquez, Ernesto Villacrés, Jazmin Arrivillaga, Martín Hinojosa & Jonathan Liria, Pp. 12515–12522

Orchid diversity in two community forests of Makawanpur District, central Nepal
-- Bijaya Pant, Mukti Ram Paudel, Mukesh Babu Chand, Shreeti Pradhan, Bijaya Bahadur Malla & Bhakta Bahadur Raskoti, Pp. 12523–12530

Habitat distribution modeling for reintroduction and conservation of *Aristolochia indica* L. - a threatened medicinal plant in Assam, India
-- Bhaskar Sarma, Prantik Sharma Baruah & Bhaben Tanti, Pp. 12531–12537

Pollination ecology of *Synedrella nodiflora* (L.) Gaertn. (Asteraceae)
-- B. Usharani & A.J. Solomon Raju, Pp. 12538–12551

Review

Status of studies on zooplankton fauna of Arunachal Pradesh, India
-- Bikramjit Sinha, Pp. 12552–12560

Short Communications

First record of the endangered Arabian Tahr *Arabitragus jayakari* (Thomas, 1894) in the Hatta Mountain Conservation Area, Dubai, United Arab Emirates
-- Jeruel Cabadonga Aguhob, Junid N. Shah, Esmat Elfaki Mohammed Elhassan, Aisha Almurr Al Muhery, Mohamed Mustafa Eltayeb Mohamed, Juma Abdulla Saeed Mohammad Al Omairi, Hamad Hashim Mohammed Khalaf Albedwawi, Obaid Mohammed Salem Mohammed Al Bedwawi, Hassan Zain Alsharif & Afra Mahmood Mohammad Ali Haji, Pp. 12561–12565

Partners

Member

Publisher & Host