



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

Journal of Threatened Taxa

Building evidence for conservation globally

www.threatenedtaxa.org

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

SHORT COMMUNICATION

FIRST RECORD OF THE SMALL BAMBOO BAT *TYLONYCTERIS FULVIDA* (PETERS, 1872) (MAMMALIA: CHIROPTERA: VESPERTILIONIDAE) FROM NEPAL

Basant Sharma, Anoj Subedi, Bandana Subedi, Shristee Panthee & Pushpa Raj Acharya

26 July 2019 | Vol. 11 | No. 9 | Pages: 14216–14219

DOI: 10.11609/jott.4502.11.9.14216-14219



For Focus, Scope, Aims, Policies, and Guidelines visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-0>

For Article Submission Guidelines, visit <https://threatenedtaxa.org/index.php/JoTT/about/submissions#onlineSubmissions>

For Policies against Scientific Misconduct, visit <https://threatenedtaxa.org/index.php/JoTT/about/editorialPolicies#custom-2>

For reprints, contact [<ravi@threatenedtaxa.org>](mailto:ravi@threatenedtaxa.org)

The opinions expressed by the authors do not reflect the views of the Journal of Threatened Taxa, Wildlife Information Liaison Development Society, Zoo Outreach Organization, or any of the partners. The journal, the publisher, the host, and the partners are not responsible for the accuracy of the political boundaries shown in the maps by the authors.

Partner



صندوق محمد بن زايد
للمحافظة على
الكائنات الحية

The Mohamed bin Zayed
SPECIES CONSERVATION FUND

Member



Publisher & Host





ISSN 0974-7907 (Online)
ISSN 0974-7893 (Print)

PLATINUM
OPEN ACCESS



FIRST RECORD OF THE SMALL BAMBOO BAT *TYLONYCTERIS FULVIDA* (PETERS, 1872) (MAMMALIA: CHIROPTERA: VESPERTILIONIDAE) FROM NEPAL

Basant Sharma¹ , Anoj Subedi² , Bandana Subedi³ , Shristee Panthee⁴ & Pushpa Raj Acharya⁵

^{1,4} Bat Friends Pokhara, Hariyokharka-15 Pokhara, Kaski, Nepal.

^{1,5} Nepal Bat Research and Conservation Union (NeBRCU), Batulechaur-16 Pokhara, Kaski, Nepal.

¹⁻⁴ Tribhuvan University, Institute of Forestry, Hariyokharka-15 Pokhara, Kaski, Nepal.

⁵ Central Campus of Science and Technology, Faculty of Science, Mid-western University, Birendranagar, Surkhet, Nepal.

¹b.s.sharma237@gmail.com (corresponding author), ²anojsubedi99@gmail.com, ³bandanasubedi84@gmail.com,

⁴shristeesharma3@gmail.com, ⁵armalepushpa@gmail.com

Abstract: A bamboo bat of the genus *Tylonycteris* was captured near Gupteshore Cave of Kushma in Parbat, Nepal. Traditionally, two species of *Tylonycteris* (of *T. pachypus* complex and *T. robustula* complex) are known from the Indian subcontinent. Due to inconsistency in taxonomic classification, several changes were recently made within the genus *Tylonycteris*—*T. pachypus* was corrected to *T. fulvida* and *T. robustula* to *T. malayana*. The occurrence of *Tylonycteris* from Nepal's diversified zoogeography, however, was never mentioned. This note provides a new record of *Tylonycteris* from Nepal. Based on morphological characteristics and species distribution range, this note confirms the captured species as *T. fulvida*.

Keywords: Gupteshore Cave, Kushma, new record, Parbat, *Tylonycteris*.

A total of 128 species of bats are reported from the Indian subcontinent, including 115 species of yangochiropterans and 13 species of yinpterochiropterans (Srinivasulu et al. 2010). Nepal records 53 species of bats within the families Pteropodidae (5), Rhinolophidae (9), Hipposideridae (4), Megadermatidae (1), Emballonuridae (1), Vespertilionidae (31), and Miniopteridae (2) (Acharya et

al. 2010). The reported number of species represents the bat diversity of about 5% of the world and over 40% of southern Asia. Among these, two species are categorized as Critically Endangered, one as Endangered, two as Vulnerable, four as Near Threatened, 25 as Least Concern, and 19 as Data Deficient in the National Red List (Jnawali et al. 2011). Vespertilionidae is the most species-rich family with 58.5% of bat species from Nepal within 15 genera. *Myotis* (7) is the most species-rich genus within this family, followed by *Murina* (3) and *Pipistrellus* (3). There has been, however, no previous evidence of the genus *Tylonycteris* in Nepal.

Traditionally, *Tylonycteris* was classified as containing only two species: *T. pachypus* (Temminck, 1840) and *T. robustula* (Thomas, 1915). Several other taxa were included as subspecies within these two species groups (Simmons 2005). Later, Feng et al. (2008) described a third species, *T. pygmaea* (Feng, Li & Wang 2008), which is smaller than its congeners. It is endemic to the Yunnan Province in southern China, while the former two

DOI: <https://doi.org/10.11609/jott.4502.11.9.14216-14219> | ZooBank: urn:lsid:zoobank.org:pub:70B837FC-1AE4-44E4-B783-302CF8283C8A

Editor: Paul Racey, University of Exeter, UK.

Date of publication: 26 July 2019 (online & print)

Manuscript details: #4502 | Received 22 August 2018 | Final received 30 June 2019 | Finally accepted 05 July 2019

Citation: Sharma, B., A. Subedi, B. Subedi, S. Panthee & P.R. Acharya (2019). First record of the Small Bamboo Bat *Tylonycteris fulvida* (Peters, 1872) (Mammalia: Chiroptera: Vespertilionidae) from Nepal. *Journal of Threatened Taxa* 11(9): 14216–14219. <https://doi.org/10.11609/jott.4502.11.9.14216-14219>

Copyright: © Sharma et al. 2019. Creative Commons Attribution 4.0 International License. JoTT allows unrestricted use, reproduction, and distribution of this article in any medium by adequate credit to the author(s) and the source of publication.

Funding: Rufford Foundation, UK.

Competing interests: The authors declare no competing interests.



Acknowledgements: We are grateful to the Rufford Foundation UK for proving grant support for the project. We would like to express our sincere thanks to Joe Chun-Chia Huang for confirming the identity of the species and for their valuable suggestions. We are also grateful to the Department of National Parks and Wildlife Conservation (DNPWC Nepal) and Department of Forests (DOF Nepal) for providing permission to carry out trapping survey in the area.




Figure 1. New record of *Tylonycteris fulvida* near Gupeteshore Cave of Kushma, Parbat, Nepal.

species have much more extensive geographic ranges that greatly overlap in southeastern Asia (Tu et al. 2017). Due to inconsistency in taxonomic classification, Tu et al. (2017) recently revalidated several changes within *Tylonycteris*—*T. pachypus* was corrected to *T. fulvida* (Blyth, 1859) and *T. robustula* to *T. malayana* (Chasen, 1940).

Both *T. fulvida* and *T. malayana* were previously recorded from the Indian subcontinent (Bates & Harrison 1997). *Tylonycteris fulvida* occurs in southern and northeastern South Asia, southern China, and much of southeastern Asia (Bates et al. 2008a). In southern Asia, this species is widely distributed in and known from India (Andaman Islands, Karnataka, Kerala, Manipur, Meghalaya, Mizoram, Sikkim, Tripura, and West Bengal) (Molur et al. 2002; Das 2003) and Bangladesh (Khan 2001; Srinivasulu & Srinivasulu 2005). *Tylonycteris malayana* ranges from northeastern India through parts of southern China to much of mainland and insular southeastern Asia (Bates et al. 2008b). In southern Asia, this species is only recorded from Mizoram and Andaman Islands in India (Molur et al. 2002; Srinivasulu et al. 2018). Both these species are listed as Least

Table 1. Morphometric measurements of *Tylonycteris* sp. captured Gupeteshore Cave at Kushma in Parbat, Nepal, compared with that of *T. fulvida* and *T. malayana*.

Parameters	Measurements (mm) (captured bat)	Bates & Harrison (1997)	
		<i>T. fulvida</i> (range)	<i>T. malayana</i> (range)
FA	26.9	26.1–29.0	26.6–28.1
HB	38.5	34.0–46.0	40.0–44.0
TL	25.8	26.0–33.0	26.0–31.0
HF	6.6	5.0–7.0	5.0–5.5
EL	9.3	9.0–10.0	8.5–10.5
TIB	11.6	-	-
3mt	26.4	23.8–27.0	25.8–26.4
1ph3mt	11.4	-	-
2ph3mt	14.6	-	-
4mt	26.6	23.8–26.9	25.4–26.0
1ph4mt	10.6	-	-
2ph4mt	7.1	-	-
5mt	26.1	23.2–26.0	24.8–25.6
1ph5mt	7.2	-	-
2ph5mt	3.5	-	-
BW (gm)	4	-	-



Image 1. Small Bamboo Bat *Tylonycteris fulvida* captured near Gupteshore Cave of Kushma, Parbat, Nepal: 1 - nostrils | 2 - shape of the head | 3 - ventral portion | 4 - lateral view | Blue circle - circular pads on thumbs | Red circle - pad on the sole of the hindfoot.

Concern in IUCN Red List (Bates et al. 2008a,b).

The study was conducted near the Gupeteshore Cave of Kushma (headquarters of Parbat District) during the field expedition of “Bats survey and conservation outreach programs along Kaligandaki Canyon of Nepal” funded by the Rufford Foundation (UK) in 2017. The cave is situated in steep slope pasture land with a small grove of trees forming a forest-like patch just above the cave structure, where trapping was conducted. The trapping area is dominated by *Dalbergia sissoo* and clusters of *Bambusa* sp.

Two mist nets (height 2.6m, lengths 4m & 6m, 38mm mesh) were deployed to capture the bats 30cm above

the ground. Mist nets were left open from 18.00h to 21.00h and continuously checked at 10-minute intervals to reduce entanglement of the trapped bats. External morphometric measurements of the trapped bats were taken using vernier callipers (0.01mm accuracy). The measurements taken include head and body length (HB), forearm length (FA), ear length (EL), tail length (TL), hind foot length (HF), tibia length (TIB), 3rd metacarpal and phalanges length (3mt, 1ph3mt, 2ph3mt), 4th metacarpal and phalanges length (4mt, 1ph4mt, 2ph4mt), and 5th metacarpal and phalanges length (5mt, 1ph5mt, 2ph5mt). Body fur and other special features were noted. Body weight (BW) was measured using a

Pesola spring balance (1gm accuracy).

A single male specimen of *Tylonycteris* sp. was caught in the mist net located at 28.226°N & 83.674°E at an elevation of 868m on 27 April 2018 at 19.30h (two hours after sunset). The morphometric measurements are given in Table 1. The recorded location of *Tylonycteris* sp. is given in Fig. 1. The bat was released after images were taken. It was identified by referring to Bates & Harrison (1997) and consultation with experts in the field.

We identified the bat on the basis of its morphological characteristics: 1) shape of head, 2) circular pads on the base of thumb, 3) pad on the sole of the hindfoot, 4) lengths of 3rd, 4th, and 5th metacarpal, and 5) pelage colouration. *Tylonycteris* is a minute bat. Its head was characteristically flattened with the nostrils projecting forward and slightly downwards (Image 1). The fleshy pads at the base of the thumb and on the sole of the hindfoot were the most striking features of *Tylonycteris* (Image 1). The wings were short with the 3rd, 4th, and 5th metacarpals about equal in length (Table 1). The dorsal pelage was thick, short, and golden-brown, except for the muzzle which was darker, while the ventral pelage was not so dense and was paler (Image 1).

The wide distribution of *T. fulvida* along southern Asia, recorded at the closest location from Nepal, i.e., in India at Sikkim and Darjeeling of West Bengal (near the border of Nepal and India), body size, and distinguishing pelage colouration strongly suggest the captured bat to be *T. fulvida* rather than *T. malayana*, as the latter has no further record from southern Asia except from Mizoram (at the border of India and Myanmar) and the Andaman Islands in India (Molur et al. 2002; Srinivasulu et al. 2018). Additionally, the pelage colouration in *T. malayana* is uniformly grey-brown dorsally (as opposed to that of the captured bat) and slightly paler ventrally (Bates & Harrison 1997; Srinivasulu et al. 2018). With this note, we confirm the presence of *T. fulvida* in Nepal, highlighting the first record for the country.


REFERENCES

- Acharya, P.R., H. Adhikari, S. Dahal, A. Thapa & S. Thapa (2010). Bats of Nepal: A Field Guide. Small Mammals Conservation and Research Foundation (SMCRF), Kantimarga, New Baneshwor, Kathmandu, Nepal, 116pp.
- Bates, P.J.J. & D.L. Harrison (1997). *Bats of the Indian Subcontinent*. Harrison Zoological Museum, Sevenoaks, England, 258pp.
- Bates, P., C. Francis, G. Rosell-Ambal & L. Heaney (2008a). *Tylonycteris robustula*. In: The IUCN Red List of Threatened Species: e.T22578A9378011. Accessed on 10 July 2018. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T22578A9378011.en>
- Bates, P., C. Francis, G. Rosell-Ambal, L. Heaney, S. Molur & C. Srinivasulu (2008b). *Tylonycteris pachypus*. In: The IUCN Red List of Threatened Species: e.T22577A9377719. Accessed on 10 July 2018. <https://doi.org/10.2305/IUCN.UK.2008.RLTS.T22577A9377719.en>
- Das, P.K. (2003). Studies on some Indian Chiroptera from West Bengal. *Records of the Zoological Survey of India, Occasional Paper* 217: 1–164.
- Feng, Q., S. Li & Y. Wang (2008). A new species of bamboo bat (Chiroptera: Vespertilionidae: *Tylonycteris*) from southwestern China. *Zoological Science* 25(2): 225–234. <https://doi.org/10.2108/zsj.25.225>
- Jnawali, S.R., H.S. Baral, S. Lee, N. Subedi, K.P. Acharya, G.P. Upadhyay, M. Pandey, R. Shrestha, D. Joshi, B.R. Lamichane, J. Griffiths, A. Khatriwoda & R. Amin (2011). The Status of Nepal's Mammals: The National Red List Series. Department of National Parks and Wildlife Conservation (DNPWC), Kathmandu, Nepal, 214pp.
- Khan, M.A.R. (2001). Status and distribution of bats in Bangladesh with notes on their ecology. *Zoos' Print Journal* 16(5): 479–483. <https://doi.org/10.11609/JoTT.ZPJ.16.5.479-83>
- Molur, S., G. Marimuthu, C. Srinivasulu, S. Mistry, A.M. Hutson, P.J.J. Bates, S. Walker, K. Padmapriya & A.R. Binupriya (2002). Status of South Asian Chiroptera: Conservation Assessment and Management Plan (C.A.M.P.). Workshop Report. Zoo Outreach Organisation, Conservation Breeding Specialist Group South Asia & Wildlife Information & Liaison Development Society, Coimbatore, India, 154pp.
- Simmons, N.B. (2005). Order Chiroptera, pp312–529. In: Wilson, D.E. & D.M. Reeder (eds.). *Mammal Species of the World: A Taxonomic and Geographic Reference*. Johns Hopkins University Press, London, 2142pp.
- Srinivasulu, C. & B. Srinivasulu (2005). A review of chiropteran diversity of Bangladesh. *Bat Net* 6(2): 5–11.
- Srinivasulu, C., P.A. Racey & S. Mistry (2010). A key to the bats (Mammalia: Chiroptera) of South Asia. *Journal of Threatened Taxa* 2(7): 1001–1076. <https://doi.org/10.11609/JoTT.o2352.1001-76>
- Srinivasulu, C., A. Srinivasulu, B. Srinivasulu & G. Jones (2018). A new subspecies of the Malayan Bamboo Bat (Chiroptera: Vespertilionidae: *Tylonycteris malayana eremtaga*) from the Andaman Islands, India. *Journal of Threatened Taxa* 10(1): 11210–11217. <https://doi.org/10.11609/jott.3906.10.1.11210-11217>
- Tu, V.T., G. Csorba, M. Ruedi, N.M. Furey, N.T. Son, V.D. Thong, C. Bonillo & A. Hassanin (2017). Comparative phylogeography of bamboo bats of the genus *Tylonycteris* (Chiroptera, Vespertilionidae) in southeast Asia. *European Journal of Taxonomy* 274: 1–38. <https://doi.org/10.5852/ejt.2017.274>





PLATINUM
OPEN ACCESS



The Journal of Threatened Taxa (JoTT) is dedicated to building evidence for conservation globally by publishing peer-reviewed articles online every month at a reasonably rapid rate at www.threatenedtaxa.org. All articles published in JoTT are registered under [Creative Commons Attribution 4.0 International License](https://creativecommons.org/licenses/by/4.0/) unless otherwise mentioned. JoTT allows unrestricted use, reproduction, and distribution of articles in any medium by providing adequate credit to the author(s) and the source of publication.

ISSN 0974-7907 (Online) | ISSN 0974-7893 (Print)

July 2019 | Vol. 11 | No. 9 | Pages: 14087–14246

Date of Publication: 26 July 2019 (Online & Print)

DOI: 10.11609/jott.2019.11.9.14087-14246

www.threatenedtaxa.org

Article

Species richness and abundance of monogonont rotifers in relation to environmental factors in the UNESCO Sakaerat Biosphere Reserve, Thailand
– Nattaporn Plangklang, Chaichat Boonyanusith & Sujeephon Athibai, Pp. 14087–14100

Communications

Distribution and habitats of *Paphiopedilum* Pfitzer (Orchidaceae) known to occur in Bhutan
– Dhan Bahadur Gurung, Nima Gyeltshen, Kezang Tobgay, Stig Dalström, Jangchu Wangdi, Bhakta Bahadur Ghalley, Lekey Chaida, Phuntsho, Ngawang Gyeltshen, Kelzang Dawa, Tandin Wangchuk, Rebecca Pradhan, Thomas Hoijer & Choki Gyeltshen, Pp. 14101–14111

Diurnal *Serianthes nelsonii* Merr. leaflet paraheliotropism reduces leaflet temperature, relieves photoinhibition, and alters nyctinastic behavior
– Thomas Edward Marler, Pp. 14112–14118

Pollination ecology of *Brownlowia tersa* (Malvaceae), a Near Threatened non-viviparous true mangrove shrub
– Aluri Jacob Solomon Raju, Pp. 14119–14127

A note on the taxonomy and natural history of the Summer Clicker *Lahugada dohertyi* (Distant, 1891) (Insecta: Hemiptera: Cicadidae) along with its distribution in northern West Bengal, India
– Vivek Sarkar, Pp. 14128–14136

Observations on nesting activity, life cycle, and brood ball morphometry of the Bordered Dung Beetle *Oniticellus cinctus* (Fabricius, 1775) (Coleoptera: Scarabaeidae) under laboratory conditions
– Amar Paul Singh, Kritish De, Shagun Mahajan, Ritwik Mondal & Virendra Prasad Uniyal, Pp. 14137–14143

Spiders of Odisha: a preliminary checklist
– Sudhir Ranjan Choudhury, Manju Siliwal & Sanjay Keshari Das, Pp. 14144–14157

Status of water birds in Haripura-Baur Reservoir, western Terai-Arc landscape, Uttarakhand, India
– Tanveer Ahmed, Harendra Singh Bargali, Deepa Bisht, Gajendra Singh Mehra & Afifullah Khan, Pp. 14158–14165

Bird diversity in the coastal talukas of Sindhudurg District, Maharashtra, India
– Golusu Babu Rao, Santhanakrishnan Babu, Goldin Quadros & Vijaykumar Anoop, Pp. 14166–14186

Greater One-horned Rhinoceros *Rhinoceros unicornis* (Mammalia: Perissodactyla: Rhinocerotidae) population census in the Rajiv Gandhi Orang National Park, Assam, India
– Deba Kumar Dutta & Parikshit Kakati, Pp. 14187–14193

Crowding, group size and population structure of the Blackbuck *Antelope cervicapra* (Linnaeus, 1758) (Mammalia: Cetartiodactyla: Bovidae) in the semi-arid habitat of Haryana, India
– Deepak Rai & Jyoti, Pp. 14194–14203

Short Communications

An updated checklist of Indian western Himalayan gymnosperms and lectotypification of three names
– Jibankumar Singh Khuraijam & Jaideep Mazumdar, Pp. 14204–14211

New record of Blue Perch *Badis badis* (Anabantiformes: Badidae) from Godavari River basin of Telangana State, India
– Kante Krishna Prasad & Chelmala Srinivasulu, Pp. 14212–14215

First record of the Small Bamboo Bat *Tylonycteris fulvida* (Peters, 1872) (Mammalia: Chiroptera: Vespertilionidae) from Nepal
– Basant Sharma, Anoj Subedi, Bandana Subedi, Shristee Panthee & Pushpa Raj Acharya, Pp. 14216–14219

Is canine distemper virus (CDV) a lurking threat to large carnivores? A case study from Ranthambhore landscape in Rajasthan, India
– Nadisha Sidhu, Jimmy Borah, Sunny Shah, Nidhi Rajput & Kajal Kumar Jadav, Pp. 14220–14223

Notes

Extended distribution of the vulnerable Cooper's Stone Flower *Corallodiscus cooperi* (Gesneriaceae) in India
– Vikas Kumar, Samiran Panday, Sudhansu Sekhar Dash, Bipin Kumar Sinha & Paramjit Singh, Pp. 14224–14227

Extended distribution record of two bellflower species of *Codonopsis* (Campanulaceae) from the Indian state of Arunachal Pradesh
– Khilendra Singh Kanwal, Umeshkumar Lalchand Tiwari, Lod Yama & Mahendra Singh Lodhi, Pp. 14228–14231

First record of the Blue-and-white Flycatcher *Cyanoptila cyanomelana* (Temminck, 1829) (Aves: Passeriformes: Muscicapidae) from Bhutan
– Kado Rinchen, Kinley Kinley, Chhimi Dorji & Dorji Wangmo, Pp. 14232–14234

Butterflies collected using malaise traps as useful bycatches for ecology and conservation
– Augusto Henrique Batista Rosa, Lucas Neves Perillo, Frederico Siqueira Neves, Danilo Bandini Ribeiro & André Victor Lucci Freitas, Pp. 14235–14237

Notes on the hairstreak butterflies *Eusaspa* Moore, 1884 (Lepidoptera: Lycaenidae) with new distribution records to the Indian eastern Himalaya
– Gaurab Nandi Das, Subrata Gayen, Motoki Saito & Kailash Chandra, Pp. 14238–14241

First report of the Australian gall midge *Actilasioptera tumidifolium* Gagné, 1999 (Diptera: Cecidomyiidae) from Andaman Islands, India
– Duraikannu Vasanthakumar & Radheshyam Murlidhar Sharma, Pp. 14242–14243

New record of Blanford's Fox *Vulpes cana* (Mammalia: Carnivora: Canidae) in central Oman: a connection between the northern and southern populations
– Taimur Alsaid, Abdulrahman Aluwaisi, Sultan Albalushi, Zahran Alabdulsalam, Said Alharsusi & Steven Ross, Pp. 14244–14246

Partner



صندوق محمد بن زايد
للمحافظة على
الكائنات الحية
The Mohamed bin Zayed
SPECIES CONSERVATION FUND

Member



Publisher & Host

